

A connection between quantum dot Dirac operators and $\bar{\partial}$ -Robin Laplacians

Joaquim Duran (CRM & UPC)
Joint work with Albert Mas (UPC & CRM)
and Tomás Sanz-Perela (UB & CRM)

Analytic and Algebraic Methods in Physics XXII (special session on Spectral Theory)

Prague, August 28, 2025



UNIVERSITAT POLITÈCNICA
DE CATALUNYA
BARCELONATECH



jduran@crm.cat

References

[D., 2025] J. Duran, *The $\bar{\partial}$ -Robin Laplacian*, arXiv:2507.16895, (2025).

[D., Mas, Sanz-Perela, 2025] J. Duran, A. Mas, T. Sanz-Perela, *A connection between quantum dot Dirac operators and $\bar{\partial}$ -Robin Laplacians in the context of shape optimization problems*, arXiv:2507.18698, (2025).


References

[D., 2025] J. Duran, *The $\bar{\partial}$ -Robin Laplacian*, arXiv:2507.16895, (2025).

[D., Mas, Sanz-Perela, 2025] J. Duran, A. Mas, T. Sanz-Perela, *A connection between quantum dot Dirac operators and $\bar{\partial}$ -Robin Laplacians in the context of shape optimization problems*, arXiv:2507.18698, (2025).


[Antunes, Benguria, Lotoreichik, Ourmières-Bonafos, 2021] P. R. S. Antunes, R. D. Benguria, V. Lotoreichik, T. Ourmières-Bonafos, *A Variational Formulation for Dirac Operators in Bounded Domains. Applications to Spectral Geometric Inequalities*, Communications in Mathematical Physics, 386 (2021), 781–818.

Graphene




Graphene quantum dot

$$\Omega \subset \mathbb{R}^2$$




Graphene quantum dot


$$\Omega \subset \mathbb{R}^2$$



Graphene quantum dot



Graphene quantum dot



Physical model

From now on, we shall assume that $\Omega \subset \mathbb{R}^2$ is a bounded \mathcal{C}^2 domain,

Physical model

From now on, we shall assume that $\Omega \subset \mathbb{R}^2$ is a bounded \mathcal{C}^2 domain, with unit outward normal vector (ν_1, ν_2) .

Physical model

From now on, we shall assume that $\Omega \subset \mathbb{R}^2$ is a bounded \mathcal{C}^2 domain, with unit outward normal vector (ν_1, ν_2) .


The energies of an electron of mass $m \geq 0$ in a graphene quantum dot Ω are modeled by the eigenvalues λ of

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } \Omega, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u & \text{on } \partial\Omega, \end{cases}$$

where $u, v : \Omega \rightarrow \mathbb{C}$ and $\theta \in [-\frac{\pi}{2}, \frac{3\pi}{2})$.


Qualitative behavior when the domain is a disk

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } D_R, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u & \text{on } \partial D_R. \end{cases} \quad \begin{matrix} m \geq 0 \\ \theta \in \left[-\frac{\pi}{2}, \frac{3\pi}{2}\right) \end{matrix}$$




Qualitative behavior when the domain is a disk

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \varphi = \lambda \varphi & \text{in } D_R, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u, \quad \varphi = (u, v)^T & \text{on } \partial D_R. \end{cases} \quad \begin{matrix} m \geq 0 \\ \theta \in \left[-\frac{\pi}{2}, \frac{3\pi}{2}\right) \end{matrix}$$




Qualitative behavior when the domain is a disk

$$\begin{cases} (-i(\sigma_1\partial_1 + \sigma_2\partial_2) + m\sigma_3)\varphi = \lambda\varphi & \text{in } D_R, \\ v = i\frac{1 - \sin\theta}{\cos\theta}(\nu_1 + i\nu_2)u, \quad \varphi = (u, v)^T & \text{on } \partial D_R. \end{cases} \quad \begin{matrix} m \geq 0 \\ \theta \in [-\frac{\pi}{2}, \frac{3\pi}{2}) \end{matrix}$$



Qualitative behavior when the domain is a disk

$$\begin{cases} (-i(\sigma_1\partial_1 + \sigma_2\partial_2) + m\sigma_3)\varphi = \lambda\varphi & \text{in } D_R, \\ \varphi = (i\cos\theta(\sigma_1\nu_1 + \sigma_2\nu_2) + \sin\theta)\sigma_3\varphi & \text{on } \partial D_R. \end{cases} \quad \begin{matrix} m \geq 0 \\ \theta \in [-\frac{\pi}{2}, \frac{3\pi}{2}) \end{matrix}$$



Quantum dot Dirac operators

For $\theta \in \left(-\frac{\pi}{2}, \frac{3\pi}{2}\right) \setminus \left\{\frac{\pi}{2}\right\}$ (Benguria, Fournais, Stockmeyer, Van Den Bosch, 2017):

$$\begin{aligned}\text{Dom}(\mathcal{D}_\theta) &:= \left\{ \varphi \in H^1(\Omega)^2 : \varphi = (i \cos \theta (\sigma_1 \nu_1 + \sigma_2 \nu_2) + \sin \theta) \sigma_3 \varphi \text{ in } H^{1/2}(\partial\Omega) \right\}, \\ \mathcal{D}_\theta \varphi &:= \left(-i(\sigma_1 \partial_1 + \sigma_2 \partial_2) + m \sigma_3 \right) \varphi \quad \text{for all } \varphi \in \text{Dom}(\mathcal{D}_\theta).\end{aligned}$$

Quantum dot Dirac operators

For $\theta \in \left(-\frac{\pi}{2}, \frac{3\pi}{2}\right) \setminus \left\{\frac{\pi}{2}\right\}$ (Benguria, Fournais, Stockmeyer, Van Den Bosch, 2017):

$$\begin{aligned}\text{Dom}(\mathcal{D}_\theta) &:= \left\{ \varphi \in H^1(\Omega)^2 : \varphi = (i \cos \theta (\sigma_1 \nu_1 + \sigma_2 \nu_2) + \sin \theta) \sigma_3 \varphi \text{ in } H^{1/2}(\partial\Omega) \right\}, \\ \mathcal{D}_\theta \varphi &:= \left(-i(\sigma_1 \partial_1 + \sigma_2 \partial_2) + m \sigma_3 \right) \varphi \quad \text{for all } \varphi \in \text{Dom}(\mathcal{D}_\theta).\end{aligned}$$

For $\theta \in \left\{-\frac{\pi}{2}, \frac{\pi}{2}\right\}$ (Schmidt, 1995):

$$\begin{aligned}\text{Dom}(\mathcal{D}_{-\frac{\pi}{2}}) &:= \left\{ \varphi = (u, v)^\top \in L^2(\Omega)^2 : (\sigma_1 \partial_1 + \sigma_2 \partial_2) \varphi \in L^2(\Omega)^2, u \in H_0^1(\Omega) \right\}, \\ \text{Dom}(\mathcal{D}_{+\frac{\pi}{2}}) &:= \left\{ \varphi = (u, v)^\top \in L^2(\Omega)^2 : (\sigma_1 \partial_1 + \sigma_2 \partial_2) \varphi \in L^2(\Omega)^2, v \in H_0^1(\Omega) \right\}, \\ \mathcal{D}_{\pm \frac{\pi}{2}} \varphi &:= \left(-i(\sigma_1 \partial_1 + \sigma_2 \partial_2) + m \sigma_3 \right) \varphi \quad \text{for all } \varphi \in \text{Dom}(\mathcal{D}_{\pm \frac{\pi}{2}}).\end{aligned}$$

Quantum dot Dirac operators

For $\theta \in \left(-\frac{\pi}{2}, \frac{3\pi}{2}\right) \setminus \left\{\frac{\pi}{2}\right\}$ (Benguria, Fournais, Stockmeyer, Van Den Bosch, 2017):

$$\begin{aligned}\text{Dom}(\mathcal{D}_\theta) &:= \left\{ \varphi \in H^1(\Omega)^2 : \varphi = (i \cos \theta (\sigma_1 \nu_1 + \sigma_2 \nu_2) + \sin \theta) \sigma_3 \varphi \text{ in } H^{1/2}(\partial\Omega) \right\}, \\ \mathcal{D}_\theta \varphi &:= \left(-i(\sigma_1 \partial_1 + \sigma_2 \partial_2) + m \sigma_3 \right) \varphi \quad \text{for all } \varphi \in \text{Dom}(\mathcal{D}_\theta).\end{aligned}$$


For $\theta \in \left\{-\frac{\pi}{2}, \frac{\pi}{2}\right\}$ (Schmidt, 1995):

$$\begin{aligned}\text{Dom}(\mathcal{D}_{-\frac{\pi}{2}}) &:= \left\{ \varphi = (u, v)^\top \in L^2(\Omega)^2 : (\sigma_1 \partial_1 + \sigma_2 \partial_2) \varphi \in L^2(\Omega)^2, u \in H_0^1(\Omega) \right\}, \\ \text{Dom}(\mathcal{D}_{+\frac{\pi}{2}}) &:= \left\{ \varphi = (u, v)^\top \in L^2(\Omega)^2 : (\sigma_1 \partial_1 + \sigma_2 \partial_2) \varphi \in L^2(\Omega)^2, v \in H_0^1(\Omega) \right\}, \\ \mathcal{D}_{\pm \frac{\pi}{2}} \varphi &:= \left(-i(\sigma_1 \partial_1 + \sigma_2 \partial_2) + m \sigma_3 \right) \varphi \quad \text{for all } \varphi \in \text{Dom}(\mathcal{D}_{\pm \frac{\pi}{2}}).\end{aligned}$$

$$\sigma(D_{\pm \frac{\pi}{2}}) = \{\pm m\} \cup \left\{ \pm \sqrt{\Lambda + m^2} : \Lambda \in \sigma(-\Delta_D) \right\}, \quad \text{with } \pm m \in \sigma_{\text{ess}}(D_{\pm \frac{\pi}{2}}).$$


Qualitative behavior when the domain is a disk

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } D_R, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u & \text{on } \partial D_R. \end{cases} \quad \begin{matrix} m \geq 0 \\ \theta \in \left[-\frac{\pi}{2}, \frac{3\pi}{2}\right) \end{matrix}$$




Invariances of quantum dot Dirac operators

$$\left\{ \begin{array}{l} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} \quad \text{in } D_R, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u \quad \text{on } \partial D_R. \end{array} \right. \quad \begin{array}{l} m \leq 0 \\ \theta \in \left[-\frac{\pi}{2}, \frac{3\pi}{2}\right) \end{array}$$




Invariances of quantum dot Dirac operators

$$\left\{ \begin{array}{l} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} \quad \text{in } D_R, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u \quad \text{on } \partial D_R. \end{array} \right. \quad \begin{array}{l} m \in \mathbb{R} \\ \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \end{array}$$



Invariances of quantum dot Dirac operators

$$\left\{ \begin{array}{l} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} \quad \text{in } D_R, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u \quad \text{on } \partial D_R. \end{array} \right. \quad \begin{array}{l} m \in \mathbb{R} \\ \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \end{array}$$



The fundamental energy


For $m \in \mathbb{R}$ and $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, the fundamental energy is

$$\lambda_{\Omega}(\theta) := \min \left(\sigma(\mathcal{D}_{\theta}) \cap [0, +\infty) \right).$$

The fundamental energy

For $m \in \mathbb{R}$ and $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$, the fundamental energy is


$$\lambda_{\Omega}(\theta) := \min \left(\sigma(\mathcal{D}_{\theta}) \cap [0, +\infty) \right).$$



The fundamental energy

For $m \in \mathbb{R}$ and $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, the fundamental energy is


$$\lambda_{\Omega}(\theta) := \min \left(\sigma(\mathcal{D}_{\theta}) \cap [0, +\infty) \right).$$



A shape optimization problem

Conjecture

Assume that $m \geq 0$. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary and let $D \subset \mathbb{R}^2$ be a disk with the same area as Ω . If Ω is not a disk, then $\lambda_\Omega(\theta) > \lambda_D(\theta)$ for all $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$.



Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } \Omega, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u \\ \end{cases} \quad \text{in } \Omega,$$

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = i(\lambda + m)\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = i(\lambda + m)\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Assume $\lambda > m$ \downarrow $(\theta, \lambda) \xrightarrow{T} (a, \mu) := \left((\lambda + m)\frac{1 - \sin \theta}{\cos \theta}, \lambda^2 - m^2 \right), \mu, a > 0$

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = i(\lambda + m)\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Assume $\lambda > m$ $\left| \begin{array}{l} (\theta, \lambda) \xrightarrow{T} (a, \mu) := \left((\lambda + m)\frac{1 - \sin \theta}{\cos \theta}, \lambda^2 - m^2 \right), \mu, a > 0 \end{array} \right.$

$$\begin{cases} -\Delta u = \mu u & \text{in } \Omega, \\ (\nu_1 - i\nu_2)(\partial_1 + i\partial_2)u + au = 0 & \text{on } \partial\Omega. \end{cases}$$

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = i(\lambda + m)\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Assume $\lambda > m$ \downarrow $(\theta, \lambda) \xrightarrow{T} (a, \mu) := \left((\lambda + m)\frac{1 - \sin \theta}{\cos \theta}, \lambda^2 - m^2 \right)$, $\mu, a > 0$

$$\begin{cases} -\Delta u = \mu u & \text{in } \Omega, \\ (\nu_1 - i\nu_2)(\partial_1 + i\partial_2)u + au = 0 & \text{on } \partial\Omega. \end{cases} \longrightarrow \text{Associated operator?}$$


The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$

The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$

- Ω is a bounded \mathcal{C}^2 domain in \mathbb{R}^2 ,




The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$

- Ω is a bounded \mathcal{C}^2 domain in \mathbb{R}^2 ,


- $f \in L^2(\Omega)$ is complex valued,



The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$


- Ω is a bounded \mathcal{C}^2 domain in \mathbb{R}^2 ,
- $f \in L^2(\Omega)$ is complex valued,
- $a > 0$, and



The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$


- Ω is a bounded \mathcal{C}^2 domain in \mathbb{R}^2 ,
- $f \in L^2(\Omega)$ is complex valued,
- $a > 0$, and



The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$

- Ω is a bounded \mathcal{C}^2 domain in \mathbb{R}^2 ,
- $f \in L^2(\Omega)$ is complex valued,
- $a > 0$, and




We identify $\mathbb{R}^2 \equiv \mathbb{C}$

$$\begin{aligned} \nu &= \nu_1 + i\nu_2 \\ \bar{\nu} &= \nu_1 - i\nu_2 \end{aligned}$$

The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$

- Ω is a bounded C^2 domain in \mathbb{R}^2 ,
- $f \in L^2(\Omega)$ is complex valued,
- $a > 0$, and



We identify $\mathbb{R}^2 \equiv \mathbb{C}$

$$\begin{aligned} \nu &= \nu_1 + i\nu_2 \\ \bar{\nu} &= \nu_1 - i\nu_2 \end{aligned}$$

$$\nabla = (\partial_1, \partial_2)$$


$$\partial_z = \frac{1}{2}(\partial_1 - i\partial_2)$$

$$\partial_{\bar{z}} = \frac{1}{2}(\partial_1 + i\partial_2)$$

The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$

- Ω is a bounded C^2 domain in \mathbb{R}^2 ,
- $f \in L^2(\Omega)$ is complex valued,
- $a > 0$, and



We identify $\mathbb{R}^2 \equiv \mathbb{C}$

$$\begin{aligned} \nu &= \nu_1 + i\nu_2 \\ \bar{\nu} &= \nu_1 - i\nu_2 \end{aligned}$$

$$\nabla = (\partial_1, \partial_2)$$

$$\partial_z = \frac{1}{2}(\partial_1 - i\partial_2)$$


$$\partial_{\bar{z}} = \frac{1}{2}(\partial_1 + i\partial_2)$$

$$2\bar{\nu}\partial_{\bar{z}}u = (\nu_1 - i\nu_2)(\partial_1 + i\partial_2)u = \partial_{\nu}u + i\partial_{\tau}u$$

The auxiliary problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases} \quad \text{where}$$

- Ω is a bounded C^2 domain in \mathbb{R}^2 ,
- $f \in L^2(\Omega)$ is complex valued,
- $a > 0$, and



We identify $\mathbb{R}^2 \equiv \mathbb{C}$

$$\begin{aligned} \nu &= \nu_1 + i\nu_2 \\ \bar{\nu} &= \nu_1 - i\nu_2 \end{aligned}$$

$$\nabla = (\partial_1, \partial_2)$$

$$\partial_z = \frac{1}{2}(\partial_1 - i\partial_2)$$

$$\partial_{\bar{z}} = \frac{1}{2}(\partial_1 + i\partial_2)$$

$$2\bar{\nu}\partial_{\bar{z}}u = (\nu_1 - i\nu_2)(\partial_1 + i\partial_2)u = \partial_{\nu}u + i\partial_{\tau}u$$

Comparison with the Robin Laplacian

$$\left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{array} \right. \quad \text{v.s.} \quad \left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ \partial_{\nu}u + au = 0 & \text{on } \partial\Omega. \end{array} \right.$$

Comparison with the Robin Laplacian

$$\left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{array} \right. \quad \text{v.s.} \quad \left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ \partial_{\nu}u + au = 0 & \text{on } \partial\Omega. \end{array} \right.$$

The weak formulation of the Robin problem is obtained from

$$\int_{\Omega} f \bar{v} = \int_{\Omega} -\Delta u \bar{v} \quad \text{for all } v \in \mathcal{C}^{\infty}(\bar{\Omega}),$$

Comparison with the Robin Laplacian

$$\left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{array} \right. \quad \text{v.s.} \quad \left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ \partial_{\nu}u + au = 0 & \text{on } \partial\Omega. \end{array} \right.$$

The weak formulation of the Robin problem is obtained from

$$\int_{\Omega} f \bar{v} = \int_{\Omega} -\Delta u \bar{v} \quad \text{for all } v \in C^{\infty}(\bar{\Omega}),$$

decomposing $\Delta = \operatorname{div} \nabla$, integrating by parts, and applying the B.C.:

$$\int_{\Omega} f \bar{v} = \int_{\Omega} -\operatorname{div}(\nabla u) \bar{v} = \int_{\Omega} \nabla u \cdot \bar{\nabla} \bar{v} - \int_{\partial\Omega} \partial_{\nu} u \bar{v} = \int_{\Omega} \nabla u \cdot \bar{\nabla} \bar{v} + a \int_{\partial\Omega} u \bar{v}.$$

Comparison with the Robin Laplacian

$$\left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{array} \right. \quad \text{v.s.} \quad \left\{ \begin{array}{ll} -\Delta u = f & \text{in } \Omega, \\ \partial_{\nu}u + au = 0 & \text{on } \partial\Omega. \end{array} \right.$$

The weak formulation of the Robin problem is obtained from

$$\int_{\Omega} f \bar{v} = \int_{\Omega} -\Delta u \bar{v} \quad \text{for all } v \in C^{\infty}(\bar{\Omega}),$$

decomposing $\Delta = \operatorname{div} \nabla$, integrating by parts, and applying the B.C.:


$$\int_{\Omega} f \bar{v} = \int_{\Omega} -\operatorname{div}(\nabla u) \bar{v} = \int_{\Omega} \nabla u \cdot \bar{\nabla} \bar{v} - \int_{\partial\Omega} \partial_{\nu} u \bar{v} = \int_{\Omega} \nabla u \cdot \bar{\nabla} \bar{v} + a \int_{\partial\Omega} u \bar{v}.$$

Decomposing, instead, $\Delta = 4\partial_z\partial_{\bar{z}}$ and integrating by parts, we get

$$\int_{\Omega} f \bar{v} = -4 \int_{\Omega} \partial_z \partial_{\bar{z}} u \bar{v} = 4 \int_{\Omega} \partial_{\bar{z}} u \overline{\partial_z v} - \int_{\partial\Omega} 2\bar{\nu} \partial_{\bar{z}} u \bar{v}.$$

Qualitative behavior when the domain is a disk

$$\begin{cases} -\Delta u = \mu u & \text{in } D_R, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial D_R. \end{cases}$$



The $\bar{\partial}$ -Robin Laplacian

Theorem (D., 2025)

For every $a > 0$, the operator

$$\text{Dom}(\mathcal{R}_a) := \left\{ u \in H^1(\Omega) : \partial_{\bar{z}} u \in H^1(\Omega), 2\bar{\nu} \partial_{\bar{z}} u + au = 0 \text{ in } H^{1/2}(\partial\Omega) \right\},$$
$$\mathcal{R}_a u := -\Delta u \quad \text{for all } u \in \text{Dom}(\mathcal{R}_a),$$

is the unique self-adjoint operator such that, for every $f \in L^2(\Omega)$, there exists a unique $u \in \text{Dom}(\mathcal{R}_a)$ solving

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ 2\bar{\nu} \partial_{\bar{z}} u + au = 0 & \text{on } \partial\Omega, \end{cases}$$

in the weak sense.

The $\bar{\partial}$ -Robin Laplacian

Theorem (D., 2025)

For every $a > 0$, the operator

$$\text{Dom}(\mathcal{R}_a) := \left\{ u \in H^1(\Omega) : \partial_{\bar{z}} u \in H^1(\Omega), 2\bar{v} \partial_{\bar{z}} u + au = 0 \text{ in } H^{1/2}(\partial\Omega) \right\},$$
$$\mathcal{R}_a u := -\Delta u \quad \text{for all } u \in \text{Dom}(\mathcal{R}_a),$$

is the unique self-adjoint operator such that, for every $f \in L^2(\Omega)$, there exists a unique $u \in \text{Dom}(\mathcal{R}_a)$ solving

$$4 \int_{\Omega} \partial_{\bar{z}} u \overline{\partial_{\bar{z}} v} + a \int_{\partial\Omega} u \bar{v} = \int_{\Omega} f \bar{v} \quad \text{for all } v \in C^\infty(\bar{\Omega}).$$

The $\bar{\partial}$ -Robin Laplacian

Theorem (D., 2025)

For every $a > 0$, the operator

$$\text{Dom}(\mathcal{R}_a) := \left\{ u \in H^1(\Omega) : \partial_{\bar{z}} u \in H^1(\Omega), 2\bar{v} \partial_{\bar{z}} u + au = 0 \text{ in } H^{1/2}(\partial\Omega) \right\},$$
$$\mathcal{R}_a u := -\Delta u \quad \text{for all } u \in \text{Dom}(\mathcal{R}_a),$$

is the unique self-adjoint operator such that, for every $f \in L^2(\Omega)$, there exists a unique $u \in \text{Dom}(\mathcal{R}_a)$ solving

$$4 \int_{\Omega} \partial_{\bar{z}} u \overline{\partial_{\bar{z}} v} + a \int_{\partial\Omega} u \bar{v} = \int_{\Omega} f \bar{v} \quad \text{for all } v \in C^\infty(\bar{\Omega}).$$

Moreover, $\sigma(\mathcal{R}_a)$ is purely discrete and contained in $(0, +\infty)$.

The $\bar{\partial}$ -Robin Laplacian

Theorem (D., 2025)

For every $a > 0$, the operator

$$\text{Dom}(\mathcal{R}_a) := \left\{ u \in H^1(\Omega) : \partial_{\bar{z}} u \in H^1(\Omega), 2\bar{v} \partial_{\bar{z}} u + au = 0 \text{ in } H^{1/2}(\partial\Omega) \right\},$$
$$\mathcal{R}_a u := -\Delta u \quad \text{for all } u \in \text{Dom}(\mathcal{R}_a),$$


is the unique self-adjoint operator such that, for every $f \in L^2(\Omega)$, there exists a unique $u \in \text{Dom}(\mathcal{R}_a)$ solving

$$4 \int_{\Omega} \partial_{\bar{z}} u \overline{\partial_{\bar{z}} v} + a \int_{\partial\Omega} u \bar{v} = \int_{\Omega} f \bar{v} \quad \text{for all } v \in E(\Omega).$$

Moreover, $\sigma(\mathcal{R}_a)$ is purely discrete and contained in $(0, +\infty)$.

The spectrum of $\overline{\partial}$ -Robin Laplacians

$$\begin{cases} -\Delta u = \mu u & \text{in } D_R, \\ 2\overline{\nu}\partial_{\overline{z}}u + au = 0 & \text{on } \partial D_R. \end{cases}$$



The first eigenvalue of $\overline{\partial}$ -Robin Laplacians

For $a > 0$, the first eigenvalue of \mathcal{R}_a is

$$\mu_{\Omega}(a) := \min (\sigma(\mathcal{R}_a)).$$

The first eigenvalue of $\bar{\partial}$ -Robin Laplacians

For $a > 0$, the first eigenvalue of \mathcal{R}_a is

$$\mu_{\Omega}(a) := \min (\sigma(\mathcal{R}_a)).$$

Theorem (D., 2025)

The following hold:

- Let $E(\Omega) = \{u \in L^2(\Omega) : \partial_{\bar{z}}u \in L^2(\Omega) \text{ and } u \in L^2(\partial\Omega)\}$. Then,

$$\mu_{\Omega}(a) = \inf_{u \in E(\Omega) \setminus \{0\}} \frac{4 \int_{\Omega} |\partial_{\bar{z}}u|^2 + a \int_{\partial\Omega} |u|^2}{\int_{\Omega} |u|^2}.$$

The first eigenvalue of $\bar{\partial}$ -Robin Laplacians

For $a > 0$, the first eigenvalue of \mathcal{R}_a is

$$\mu_{\Omega}(a) := \min (\sigma(\mathcal{R}_a)).$$

Theorem (D., 2025)

The following hold:

- Let $E(\Omega) = \{u \in L^2(\Omega) : \partial_{\bar{z}}u \in L^2(\Omega) \text{ and } u \in L^2(\partial\Omega)\}$. Then,

$$\mu_{\Omega}(a) = \inf_{u \in E(\Omega) \setminus \{0\}} \frac{4 \int_{\Omega} |\partial_{\bar{z}}u|^2 + a \int_{\partial\Omega} |u|^2}{\int_{\Omega} |u|^2}.$$

- The function $a \mapsto \mu_{\Omega}(a)$ is continuous, strictly increasing, and bijective from $(0, +\infty)$ to $(0, \Lambda_{\Omega})$, where Λ_{Ω} is the first eigenvalue of the Dirichlet Laplacian in Ω .

Relation with quantum dot Dirac operators

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = i(\lambda + m)\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Assume $\lambda > m$ \downarrow $(\theta, \lambda) \xrightarrow{T} (a, \mu) := \left((\lambda + m)\frac{1 - \sin \theta}{\cos \theta}, \lambda^2 - m^2 \right)$, $\mu, a > 0$

$$\begin{cases} -\Delta u = \mu u & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega. \end{cases} \longrightarrow \bar{\partial}\text{-Robin Laplacian } \mathcal{R}_a$$

Fundamental energies are mapped to each other

Proposition (D., Mas, Sanz-Perela, 2025)

Let $(\theta, \lambda) \in (-\frac{\pi}{2}, \frac{\pi}{2}) \times (m, +\infty)$ and $(a, \mu) \in (0, +\infty) \times (0, +\infty)$ be such that $T(\theta, \lambda) = (a, \mu)$. Then, $\lambda = \lambda_{\Omega}(\theta)$ if and only if $\mu = \mu_{\Omega}(a)$.

Fundamental energies are mapped to each other

Proposition (D., Mas, Sanz-Perela, 2025)

Let $(\theta, \lambda) \in (-\frac{\pi}{2}, \frac{\pi}{2}) \times (m, +\infty)$ and $(a, \mu) \in (0, +\infty) \times (0, +\infty)$ be such that $T(\theta, \lambda) = (a, \mu)$. Then, $\lambda = \lambda_\Omega(\theta)$ if and only if $\mu = \mu_\Omega(a)$.

Conjecture (QD)

Assume that $m \geq 0$. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary and let $D \subset \mathbb{R}^2$ be a disk with the same area as Ω . If Ω is not a disk, then $\lambda_\Omega(\theta) > \lambda_D(\theta)$ for all $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$.

Fundamental energies are mapped to each other

Proposition (D., Mas, Sanz-Perela, 2025)

Let $(\theta, \lambda) \in (-\frac{\pi}{2}, \frac{\pi}{2}) \times (m, +\infty)$ and $(a, \mu) \in (0, +\infty) \times (0, +\infty)$ be such that $T(\theta, \lambda) = (a, \mu)$. Then, $\lambda = \lambda_\Omega(\theta)$ if and only if $\mu = \mu_\Omega(a)$.

Conjecture (QD)

Assume that $m \geq 0$. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary and let $D \subset \mathbb{R}^2$ be a disk with the same area as Ω . If Ω is not a disk, then $\lambda_\Omega(\theta) > \lambda_D(\theta)$ for all $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$.

Conjecture ($\bar{\partial}$ -R)

Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary and let $D \subset \mathbb{R}^2$ be a disk with the same area as Ω . If Ω is not a disk, then $\mu_\Omega(a) > \mu_D(a)$ for all $a > 0$.

Equivalence of conjectures

Conjectures (QD) and $(\bar{\partial}\text{-R})$ are equivalent. More specifically:

Equivalence of conjectures

Conjectures (QD) and $(\bar{\partial}\text{-R})$ are equivalent. More specifically:

Theorem (D., Mas, Sanz-Perela, 2025)

- Given $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, set

$$a := \mu_{\Omega}^{-1}(\lambda_D(\theta)^2 - m^2) > 0.$$

Equivalence of conjectures

Conjectures (QD) and $(\bar{\partial}\text{-R})$ are equivalent. More specifically:

Theorem (D., Mas, Sanz-Perela, 2025)

- Given $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, set

$$a := \mu_{\Omega}^{-1}(\lambda_D(\theta)^2 - m^2) > 0.$$

If $\mu_{\Omega}(a) > \mu_D(a)$ then $\lambda_{\Omega}(\theta) > \lambda_D(\theta)$.

Equivalence of conjectures

Conjectures (QD) and $(\bar{\partial}\text{-R})$ are equivalent. More specifically:

Theorem (D., Mas, Sanz-Perela, 2025)

- Given $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, set

$$a := \mu_{\Omega}^{-1}(\lambda_D(\theta)^2 - m^2) > 0.$$

If $\mu_{\Omega}(a) > \mu_D(a)$ then $\lambda_{\Omega}(\theta) > \lambda_D(\theta)$.

- Given $a > 0$, set

$$\theta := \lambda_{\Omega}^{-1}(\sqrt{\mu_D(a) + m^2}) \in (-\frac{\pi}{2}, \frac{\pi}{2}).$$

Equivalence of conjectures

Conjectures (QD) and $(\bar{\partial}\text{-R})$ are equivalent. More specifically:

Theorem (D., Mas, Sanz-Perela, 2025)

- Given $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, set

$$a := \mu_{\Omega}^{-1}(\lambda_D(\theta)^2 - m^2) > 0.$$

If $\mu_{\Omega}(a) > \mu_D(a)$ then $\lambda_{\Omega}(\theta) > \lambda_D(\theta)$.

- Given $a > 0$, set

$$\theta := \lambda_{\Omega}^{-1}(\sqrt{\mu_D(a) + m^2}) \in (-\frac{\pi}{2}, \frac{\pi}{2}).$$

If $\lambda_{\Omega}(\theta) > \lambda_D(\theta)$ then $\mu_{\Omega}(a) > \mu_D(a)$.

The strategy

Instead of tackling the conjecture for the quantum dot Dirac operators \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$:

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda_\Omega(\theta) - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda_\Omega(\theta) + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega, \end{cases}$$

The strategy

Instead of tackling the conjecture for the quantum dot Dirac operators \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$:

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda_\Omega(\theta) - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda_\Omega(\theta) + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega, \end{cases}$$

we tackle the conjecture for the $\bar{\partial}$ -Robin Laplacians \mathcal{R}_a , $a > 0$:

$$\begin{cases} -\Delta u = \mu_\Omega(a)u & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases}$$

The strategy

Instead of tackling the conjecture for the quantum dot Dirac operators \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$:

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda_\Omega(\theta) - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda_\Omega(\theta) + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega, \end{cases}$$

we tackle the conjecture for the $\bar{\partial}$ -Robin Laplacians \mathcal{R}_a , $a > 0$:

$$\begin{cases} -\Delta u = \mu_\Omega(a)u & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega, \end{cases}$$


for which

$$\mu_\Omega(a) = \inf_{u \in E(\Omega) \setminus \{0\}} \frac{4 \int_\Omega |\partial_{\bar{z}}u|^2 + a \int_{\partial\Omega} |u|^2}{\int_\Omega |u|^2}.$$

Validity of the $(\bar{\partial}$ -R) conjecture in the regime $a \rightarrow +\infty$


Theorem (D., Mas, Sanz-Perela, 2025)

Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary and let $D \subset \mathbb{R}^2$ be a disk with the same area as Ω . If Ω is not a disk, then there exists $a_0 > 0$ (depending on Ω) such that $\mu_\Omega(a) > \mu_D(a)$ for all $a \in (a_0, +\infty)$.



Intuition for the regime $a \rightarrow 0^+$

$$\begin{aligned} \lim_{a \rightarrow 0^+} \frac{\mu_\Omega(a)}{a} &= \lim_{a \rightarrow 0^+} \inf_{u \in E(\Omega) \setminus \{0\}} \frac{\frac{4}{a} \int_\Omega |\partial_{\bar{z}} u|^2 + \int_{\partial\Omega} |u|^2}{\int_\Omega |u|^2} \\ &= \inf_{u \in E(\Omega) \setminus \{0\}: \partial_{\bar{z}} u = 0 \text{ in } \Omega} \frac{\int_{\partial\Omega} |u|^2}{\int_\Omega |u|^2} =: S_\Omega \end{aligned}$$




Validity of the $(\bar{\partial}$ -R) conjecture in the regime $a \rightarrow 0^+$

Theorem (D., Mas, Sanz-Perela, 2025)

Let $\Omega \subset \mathbb{R}^2$ be a \mathcal{C}^2 , bounded, simply connected domain, and let $D \subset \mathbb{R}^2$ be a disk with the same area as Ω . Then,

$$\lim_{a \rightarrow 0^+} \frac{\mu_\Omega(a)}{a} = \inf_{u \in E(\Omega) \setminus \{0\}: \partial_{\bar{z}} u = 0 \text{ in } \Omega} \frac{\int_{\partial\Omega} |u|^2}{\int_\Omega |u|^2} =: S_\Omega \geq S_D = 2\sqrt{\frac{\pi}{|\Omega|}},$$

with equality if and only if Ω is a disk. As a consequence, there exists $a_1 > 0$ (depending on Ω) such that $\mu_\Omega(a) > \mu_D(a)$ for all $a \in (0, a_1)$.



Translation to the quantum dot Dirac operators

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m \geq 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = i(\lambda + m)\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

Assume $\lambda > m$ \downarrow $(\theta, \lambda) \xrightarrow{T} (a, \mu) := \left((\lambda + m)\frac{1 - \sin \theta}{\cos \theta}, \lambda^2 - m^2 \right)$, $\mu, a > 0$

$$\begin{cases} -\Delta u = \mu u & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega. \end{cases} \longrightarrow \text{Conjecture } (\bar{\partial}\text{-R}) \text{ valid for } a \rightarrow 0^+, +\infty.$$


Theorem (D., Mas, Sanz-Perela, 2025)

Assume that $m \geq 0$. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary and let $D \subset \mathbb{R}^2$ be a disk with the same area as Ω . If Ω is not a disk, then:

- There exists $\theta_0 \in (-\frac{\pi}{2}, \frac{\pi}{2})$ (depending on Ω) such that $\lambda_\Omega(\theta) > \lambda_D(\theta)$ for all $\theta \in (-\frac{\pi}{2}, \theta_0)$.
- If in addition Ω is simply connected, then there exists $\theta_1 \in (-\frac{\pi}{2}, \frac{\pi}{2})$ (depending on Ω) such that $\lambda_\Omega(\theta) > \lambda_D(\theta)$ for all $\theta \in (\theta_1, \frac{\pi}{2})$.

The case of negative mass

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } D_R, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u & \text{on } \partial D_R. \end{cases} \quad \begin{matrix} m < 0 \\ \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \end{matrix}$$



Same construction

Quantum dot Dirac operator \mathcal{D}_θ , $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $m < 0$.

$$\begin{cases} -i(\partial_1 - i\partial_2)v = (\lambda - m)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = (\lambda + m)v & \text{in } \Omega, \\ v = i\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$



$$\begin{cases} -\Delta u = (\lambda^2 - m^2)u & \text{in } \Omega, \\ -i(\partial_1 + i\partial_2)u = i(\lambda + m)\frac{1 - \sin \theta}{\cos \theta}(\nu_1 + i\nu_2)u & \text{on } \partial\Omega. \end{cases}$$

If $\lambda \rightarrow |m|^+$ \downarrow $(\theta, \lambda) \xrightarrow{T} (a, \mu) := \left((\lambda + m)\frac{1 - \sin \theta}{\cos \theta}, \lambda^2 - m^2 \right) \rightarrow (0^+, 0^+)$

$$\begin{cases} -\Delta u = \mu u & \text{in } \Omega, \\ 2\bar{\nu}\partial_{\bar{z}}u + au = 0 & \text{on } \partial\Omega. \end{cases}$$

\longrightarrow Conjecture ($\bar{\partial}$ -R) valid for $a \rightarrow 0^+$.

Another shape optimization problem

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = |m| \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } \Omega, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u & \text{on } \partial\Omega. \end{cases} \quad (1)$$

Theorem (D., Mas, Sanz-Perela, 2025)

Assume that $m < 0$, and set $\vartheta(\theta) := \frac{1 - \sin \theta}{\cos \theta}$. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary. Then:

Another shape optimization problem

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = |m| \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } \Omega, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u & \text{on } \partial\Omega. \end{cases} \quad (1)$$

Theorem (D., Mas, Sanz-Perela, 2025)

Assume that $m < 0$, and set $\vartheta(\theta) := \frac{1 - \sin \theta}{\cos \theta}$. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary. Then:

- $\min \left\{ \theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) : (1) \text{ has a nonzero solution} \right\} = \vartheta^{-1} \left(\frac{2|m|}{S_\Omega} \right).$

Another shape optimization problem

$$\begin{cases} \begin{pmatrix} m & -i(\partial_1 - i\partial_2) \\ -i(\partial_1 + i\partial_2) & -m \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = |m| \begin{pmatrix} u \\ v \end{pmatrix} & \text{in } \Omega, \\ v = i \frac{1 - \sin \theta}{\cos \theta} (\nu_1 + i\nu_2) u & \text{on } \partial\Omega. \end{cases} \quad (1)$$

Theorem (D., Mas, Sanz-Perela, 2025)

Assume that $m < 0$, and set $\vartheta(\theta) := \frac{1 - \sin \theta}{\cos \theta}$. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 boundary. Then:

- $\min \left\{ \theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) : (1) \text{ has a nonzero solution} \right\} = \vartheta^{-1} \left(\frac{2|m|}{S_\Omega} \right).$
- If in addition Ω is simply connected, then

$$\min \left\{ \theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) : (1) \text{ has a nonzero solution} \right\} \geq \vartheta^{-1} \left(|m| \sqrt{\frac{|\Omega|}{\pi}} \right)$$

and the equality holds if and only if Ω is a disk of the same area.

Acknowledgements



UNIVERSITAT POLITÈCNICA
DE CATALUNYA
BARCELONATECH




**Generalitat
de Catalunya**



**Institut
d'Estudis
Catalans**



**Nuclear
Physics
Institute**



References

[D., 2025] J. Duran, *The $\bar{\partial}$ -Robin Laplacian*, arXiv:2507.16895, (2025).



[D., Mas, Sanz-Perela, 2025] J. Duran, A. Mas, T. Sanz-Perela, *A connection between quantum dot Dirac operators and $\bar{\partial}$ -Robin Laplacians in the context of shape optimization problems*, arXiv:2507.18698, (2025).

jduran@crm.cat