

Eigenvalue bounds in the linear stability of the Ekman spiral

Borbala Gerhat

Institute of Science and Technology Austria

28 August 2025



B. Gerhat, O.O. Ibrogimov and P. Siegl

(Comm. Math. Phys. 2022)

On the point spectrum in the Ekman boundary layer problem



B. Gerhat

(in preparation)

The number of eigenvalues in the Ekman boundary layer problem

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No. 101034413.



Ekman boundary layer


[Ekman (1905)]

- Navier–Stokes in rotating frame

$$\left. \begin{aligned} \partial_t u - \nu \Delta u + \Omega e_3 \times u + (u \cdot \nabla) u + \nabla p &= 0 \\ \operatorname{div} u &= 0 \end{aligned} \right\}$$

- unknown **velocity** u and **pressure** p on **half space**
- surface and bottom **boundary conditions**

$$\left. \begin{aligned} u(t, x, y, 0) &= (0, 0, 0) \\ \lim_{z \rightarrow \infty} u(t, x, y, z) &= (u_\infty, 0, 0) \end{aligned} \right\}$$



Ekman boundary layer


[Ekman (1905)]

- Navier–Stokes in rotating frame

$$\left. \begin{aligned} \partial_t u - \nu \Delta u + \Omega e_3 \times u + (u \cdot \nabla) u + \nabla p &= 0 \\ \operatorname{div} u &= 0 \end{aligned} \right\}$$

- unknown **velocity** u and **pressure** p on **half space**
- surface and bottom **boundary conditions**

$$\left. \begin{aligned} u(t, x, y, 0) &= (0, 0, 0) \\ \lim_{z \rightarrow \infty} u(t, x, y, z) &= (u_\infty, 0, 0) \end{aligned} \right\}$$



Ekman spiral

$$\delta = (2\nu/\Omega)^{\frac{1}{2}}$$

$$u_E(z) = u_\infty \left(1 - e^{-\frac{z}{\delta}} \cos\left(\frac{z}{\delta}\right), e^{-\frac{z}{\delta}} \sin\left(\frac{z}{\delta}\right), 0 \right), \quad p_E(y) = -\Omega u_\infty y$$

Spectral problem

Non-selfadjoint operator matrix family

$$R = u_\infty \delta / \nu \geq 0, \quad \alpha > 0$$

$$\mathcal{T}(\lambda) = \begin{pmatrix} (-\partial_x^2 + \alpha^2)(-\partial_x^2 + \alpha^2 - \lambda) & 2\partial_x \\ 2\partial_x & -\partial_x^2 + \alpha^2 - \lambda \end{pmatrix} + i\alpha R \begin{pmatrix} V(-\partial_x^2 + \alpha^2) + V'' & 0 \\ U' & V \end{pmatrix}$$

- constant **domain** in ambient **Hilbert space**

$$\text{Dom } \mathcal{T} = \left\{ (f, g) \in H^4(\mathbb{R}_+) \times H^2(\mathbb{R}_+) \mid f(0) = f'(0) = g(0) = 0 \right\} \subseteq L^2(\mathbb{R}_+) \oplus L^2(\mathbb{R}_+)$$

Spectral problem

Non-selfadjoint operator matrix family

$$R = u_\infty \delta / \nu \geq 0, \quad \alpha > 0$$

$$\mathcal{T}(\lambda) = \begin{pmatrix} (-\partial_x^2 + \alpha^2)(-\partial_x^2 + \alpha^2 - \lambda) & 2\partial_x \\ 2\partial_x & -\partial_x^2 + \alpha^2 - \lambda \end{pmatrix} + i\alpha R \begin{pmatrix} V(-\partial_x^2 + \alpha^2) + V'' & 0 \\ U' & V \end{pmatrix}$$

- constant **domain** in ambient **Hilbert space**

$$\text{Dom } \mathcal{T} = \left\{ (f, g) \in H^4(\mathbb{R}_+) \times H^2(\mathbb{R}_+) \mid f(0) = f'(0) = g(0) = 0 \right\} \subseteq L^2(\mathbb{R}_+) \oplus L^2(\mathbb{R}_+)$$

- generalised **coefficients**

$$u_E \rightarrow (U, V, 0)$$

$$U', V, V', V'' \in L^1(\mathbb{R}_+) \cap L^\infty(\mathbb{R}_+)$$

Spectral problem

Non-selfadjoint operator matrix family

$$R = u_\infty \delta / \nu \geq 0, \quad \alpha > 0$$

$$\mathcal{T}(\lambda) = \begin{pmatrix} (-\partial_x^2 + \alpha^2)(-\partial_x^2 + \alpha^2 - \lambda) & 2\partial_x \\ 2\partial_x & -\partial_x^2 + \alpha^2 - \lambda \end{pmatrix} + i\alpha R \begin{pmatrix} V(-\partial_x^2 + \alpha^2) + V'' & 0 \\ U' & V \end{pmatrix}$$

- constant **domain** in ambient **Hilbert space**

$$\text{Dom } \mathcal{T} = \left\{ (f, g) \in H^4(\mathbb{R}_+) \times H^2(\mathbb{R}_+) \mid f(0) = f'(0) = g(0) = 0 \right\} \subseteq L^2(\mathbb{R}_+) \oplus L^2(\mathbb{R}_+)$$

- generalised **coefficients**

$$u_E \rightarrow (U, V, 0)$$

$$U', V, V', V'' \in L^1(\mathbb{R}_+) \cap L^\infty(\mathbb{R}_+)$$

- structure / location of spectrum**

$$\sigma(\mathcal{T}) = \left\{ \lambda \in \mathbb{C} \mid 0 \in \sigma(\mathcal{T}(\lambda)) \right\},$$

$$\forall \alpha > 0 : \text{Re } \sigma(\mathcal{T}) \geq 0 ?$$

$$\text{dependence on } R ?$$

Previous results

- **experiments**, non-rigorous **numerical stability** analysis [Faller (1963), Lilly (1966), Spooner (1982)]
- local **existence** and **uniqueness** [Giga–Inui–Mahalov–Matsui (2007), Giga–Saal (2013)]
- non-linear **stability analysis** [Giga–Saal (2015), Hess–Hieber–Mahalov–Saal (2010), Fischer–Saal (2013)]

Previous results

- **experiments**, non-rigorous **numerical stability** analysis [Faller (1963), Lilly (1966), Spooner (1982)]
- local **existence** and **uniqueness** [Giga–Inui–Mahalov–Matsui (2007), Giga–Saal (2013)]
- non-linear **stability analysis** [Giga–Saal (2015), Hess–Hieber–Mahalov–Saal (2010), Fischer–Saal (2013)]

Spectral theory

[Greenberg–Marletta (2004), Marletta–Tretter (2007)]


$$\sigma_{\text{ess}}(\mathcal{T}) = \left\{ \lambda \in \mathbb{C} \mid \exists \xi \in \mathbb{R} : (\xi^2 + \alpha^2)(\xi^2 + \alpha^2 - \lambda)^2 + 4\xi^2 = 0 \right\}$$

- **numerical range** enclosure

$$\sigma(\mathcal{T}) \subseteq \left\{ \lambda \in \mathbb{C} \mid \operatorname{Re} \lambda \geq \gamma, |\operatorname{Im} \lambda| \leq \eta \right\}$$

- only **discrete eigenvalues** in **exterior** of $\sigma_{\text{ess}}(\mathcal{T})$

$$\# \sigma(\mathcal{T}) \setminus \sigma_{\text{ess}}(\mathcal{T}) = ?$$



$$\Sigma(\mathcal{T}) = \sigma(\mathcal{T}) \setminus \sigma_{\text{ess}}(\mathcal{T}) \subseteq \sigma_{\text{p}}(\mathcal{T})$$

Eigenvalue enclosure / counting

$$\Sigma(\mathcal{T}) = \sigma(\mathcal{T}) \setminus \sigma_{\text{ess}}(\mathcal{T}) \subseteq \sigma_{\text{p}}(\mathcal{T})$$

Theorem

[G–Ibrogimov–Siegl (2022)]

$$\Sigma(\mathcal{T}) \subseteq \left\{ \lambda \in \mathbb{C} \setminus \sigma_{\text{ess}}(\mathcal{T}) \mid r(\mathcal{Q}(\lambda)) \geq 1 \right\}$$


is **bounded** and **discrete** with

$$r(\mathcal{Q}(\lambda)) = \mathcal{O}\left(|\lambda|^{-\frac{1}{2}}\right), \quad \lambda \rightarrow \infty$$

Theorem

[G (prep)]

$$U', V, V'' \text{ decay exponentially} \implies \#\Sigma(\mathcal{T}) < \infty$$



Eigenvalue enclosure / counting

$$\Sigma(\mathcal{T}) = \sigma(\mathcal{T}) \setminus \sigma_{\text{ess}}(\mathcal{T}) \subseteq \sigma_{\text{p}}(\mathcal{T})$$

Theorem

[G–Ibrogimov–Siegl (2022)]

$$\Sigma(\mathcal{T}) \subseteq \left\{ \lambda \in \mathbb{C} \setminus \sigma_{\text{ess}}(\mathcal{T}) \mid r(\mathcal{Q}(\lambda)) \geq 1 \right\}$$


is **bounded** and **discrete** with

$$r(\mathcal{Q}(\lambda)) = \mathcal{O}\left(|\lambda|^{-\frac{1}{2}}\right), \quad \lambda \rightarrow \infty$$

Theorem

[G (prep)]

$$U', V, V'' \text{ decay exponentially} \implies \#\Sigma(\mathcal{T}) < \infty$$



Conjecture / work in progress

$$\forall \varepsilon > 0 : \#\Sigma(\mathcal{T}) \leq R^2 C_\varepsilon(\alpha) \left(\int_0^\infty e^{\varepsilon x} W(x) dx \right)^2, \quad W := \max\{|V|, |V''|, |U'|\}$$

Strategy

$$\mathcal{T}(\lambda) = \underbrace{\begin{pmatrix} (-\partial_x^2 + \alpha^2)(-\partial_x^2 + \alpha^2 - \lambda) & 2\partial_x \\ 2\partial_x & -\partial_x^2 + \alpha^2 - \lambda \end{pmatrix}}_{=\mathcal{L}(\lambda)} + i\alpha R \underbrace{\begin{pmatrix} V(-\partial_x^2 + \alpha^2) + V'' & 0 \\ U' & V \end{pmatrix}}_{=\mathcal{V}_2\mathcal{V}_1}$$

- **methods** from analysis of **Schrödinger** operators

- ⚠ operator **family** with **matrix structure**
- ⚠ free operator is **non-selfadjoint** and **4th order**
- ⚠ **perturbation** is **differential** operator

Strategy

$$\mathcal{T}(\lambda) = \underbrace{\begin{pmatrix} (-\partial_x^2 + \alpha^2)(-\partial_x^2 + \alpha^2 - \lambda) & 2\partial_x \\ 2\partial_x & -\partial_x^2 + \alpha^2 - \lambda \end{pmatrix}}_{=\mathcal{L}(\lambda)} + i\alpha R \underbrace{\begin{pmatrix} V(-\partial_x^2 + \alpha^2) + V'' & 0 \\ U' & V \end{pmatrix}}_{=\mathcal{V}_2\mathcal{V}_1}$$

- **methods** from analysis of **Schrödinger** operators



operator **family** with **matrix structure**



free operator is **non-selfadjoint** and **4th order**



perturbation is **differential** operator

- **holomorphic** family of **HS** integral operators

$$\mathbb{C} \setminus \sigma_{\text{ess}}(\mathcal{T}) \ni \lambda \quad \longmapsto \quad \mathcal{Q}(\lambda) = i\alpha R \mathcal{V}_1 \mathcal{L}(\lambda)^{-1} \mathcal{V}_2 \in \mathcal{S}_2$$

$$\Omega := \mathbb{C} \setminus \sigma_{\text{ess}}(\mathcal{T}) \ni \lambda \longmapsto \mathcal{Q}(\lambda) = i\alpha R \mathcal{V}_1 \mathcal{L}(\lambda)^{-1} \mathcal{V}_2$$

[Birman, Schwinger, Simon, Abramov–Aslanyan–Davies (2001), ...]

- **Birman–Schwinger** principle

$$\lambda \in \Sigma(\mathcal{T}) \iff \lambda \in \sigma(I + \mathcal{Q})$$

$$\Omega := \mathbb{C} \setminus \sigma_{\text{ess}}(\mathcal{T}) \ni \lambda \longmapsto \mathcal{Q}(\lambda) = i\alpha R \mathcal{V}_1 \mathcal{L}(\lambda)^{-1} \mathcal{V}_2$$

[Birman, Schwinger, Simon, Abramov–Aslanyan–Davies (2001), ...]

- **Birman–Schwinger principle**

$$\lambda \in \Sigma(\mathcal{T}) \iff \lambda \in \sigma(I + \mathcal{Q})$$


- bounded **spectral enclosure**

$$(\mathcal{Q} \rightarrow \mathcal{U}^{-1} \mathcal{Q} \mathcal{U})$$

$$\|\mathcal{Q}(\lambda)\| \leq \|\mathcal{Q}(\lambda)\|_{\text{HS}} = \mathcal{O}\left(|\lambda|^{-\frac{1}{2}}\right), \quad \lambda \rightarrow \infty$$

- both \mathcal{T} and $I + \mathcal{Q}$ have **spectrum of finite type** in Ω

$$m_{\text{alg}}(\lambda, \mathcal{T}) = m_{\text{alg}}(\lambda, I + \mathcal{Q})$$



No EV accumulation

Relate to holomorphic function

[Dunford–Schwarz, Gohberg–Krein]

$$\lambda \in \Sigma(\mathcal{T}) \iff \lambda \in \sigma(I + \mathcal{Q}) \iff \det_2(1 + \mathcal{Q}(\lambda)) = 0$$

- preserves **algebraic multiplicities**

[Frank (2018)]


- consider **two** connected **components**

$$\Omega = \Omega_+ \cup \Omega_-, \quad (\alpha^2, \infty) \subseteq \Omega_+$$

- **extend** holomorphically to small **neighbourhood**

$$f_{\pm} : (\Omega_{\pm})_{\varepsilon} \rightarrow \mathbb{C}, \quad f_{\pm}(\lambda) = \det_2(1 + \mathcal{Q}(\lambda)), \quad \lambda \in \Omega_{\pm}$$

- roots **do not accumulate** in **interior** of domain !



Counting zeros - work in progress ...

$$\lambda \in \Sigma(\mathcal{T}) \iff \det_2(1 + \mathcal{Q}(\lambda)) = 0$$

(including multiplicities)

Counting zeros - work in progress ...

$$\lambda \in \Sigma(\mathcal{T}) \iff \det_2(1 + \mathcal{Q}(\lambda)) = 0$$

(including multiplicities)

- bound in **Schrödinger** case

[Frank–Laptev–Safronov (2016)]

$$\forall \varepsilon > 0 : \# \sigma_{\text{disc}}(-\Delta_{\mathbb{D}}^{\mathbb{R}_+} + q) \leq \frac{1}{\varepsilon^2} \left(\int_0^\infty e^{\varepsilon x} |q(x)| dx \right)^2$$

Counting zeros - work in progress ...

$$\lambda \in \Sigma(\mathcal{T}) \iff \det_2(1 + \mathcal{Q}(\lambda)) = 0$$

(including multiplicities)

- bound in **Schrödinger** case

[Frank–Laptev–Safronov (2016)]

$$\forall \varepsilon > 0 : \# \sigma_{\text{disc}}(-\Delta_{\mathbb{D}^+} + q) \leq \frac{1}{\varepsilon^2} \left(\int_0^\infty e^{\varepsilon x} |q(x)| dx \right)^2$$

- more **complicated geometry**

$$\xi^2 - \lambda \leftrightarrow (\xi^2 + \alpha^2)(\xi^2 + \alpha^2 - \lambda)^2 + 4\xi^2$$

- suitable **biholomorphic** transformation with **controlled derivative**

$$" \sqrt{\Omega_{\pm}} \rightarrow \{ \operatorname{Im} z > 0 \} "$$

Thank you !

- [1] Abramov, A. A., Aslanyan, A., and Davies, E. B. (2001). Bounds on complex eigenvalues and resonances. *J. Phys. A Math. Gen.*, 34(1):57–72.
- [2] Frank, R. L. (2018). Eigenvalue bounds for Schrödinger operators with complex potentials. III. *Trans. Amer. Math. Soc.*, 370(1):219–240.
- [3] Frank, R. L., Laptev, A., and Safronov, O. (2016). On the number of eigenvalues of Schrödinger operators with complex potentials. *J. London Math. Soc. (2)*, 94(2):377–390.
- [4] Gerhat, B., Ibrogimov, O. O., and Siegl, P. (2022). On the point spectrum in the Ekman boundary layer problem. *Comm. Math. Phys.*, 392(2):377–397.
- [5] Greenberg, L. and Marletta, M. (2004). The Ekman flow and related problems: spectral theory and numerical analysis. *Math. Proc. Camb. Philos. Soc.*, 136(3):719–764.
- [6] Marletta, M. and Tretter, C. (2007). Essential spectra of coupled systems of differential equations and applications in hydrodynamics. *J. Differ. Equ.*, 243(1):36–69.