

Local perturbations of potential well arrays

David Spitzkopf, in collaboration with Pavel Exner

AAMP25, 26th of August, 2025

Introduction

Question: What happens if one locally perturbs a periodic Schrödinger operator?

Introduction

Question: What happens if one locally perturbs a periodic Schrödinger operator?

In 1D this was studied extensively since the 1970s (e.g. [Zh71], [Fa89], [GS93]).

Introduction

Question: What happens if one locally perturbs a periodic Schrödinger operator?

In 1D this was studied extensively since the 1970s (e.g. [Zh71], [Fa89], [GS93]).

Higher-dimensional fully periodic settings were analyzed later ([DH86], [FK98], [ADH89]).

Introduction

Question: What happens if one locally perturbs a periodic Schrödinger operator?

In 1D this was studied extensively since the 1970s (e.g. [Zh71], [Fa89], [GS93]).

Higher-dimensional fully periodic settings were analyzed later ([DH86], [FK98], [ADH89]).

Related mechanism: “conspiracy of wells” — nearby attractive wells can bind even if each alone is subcritical. ([KS79])

Introduction

Question: What happens if one locally perturbs a periodic Schrödinger operator?

In 1D this was studied extensively since the 1970s (e.g. [Zh71], [Fa89], [GS93]).

Higher-dimensional fully periodic settings were analyzed later ([DH86], [FK98], [ADH89]).

Related mechanism: “conspiracy of wells” — nearby attractive wells can bind even if each alone is subcritical. ([KS79])

Recently: geometric perturbations of soft waveguides [EV24], and curved arrays of symmetric wells [Ex24], also produce discrete spectrum.

Introduction

Question: What happens if one locally perturbs a periodic Schrödinger operator?

In 1D this was studied extensively since the 1970s (e.g. [Zh71], [Fa89], [GS93]).

Higher-dimensional fully periodic settings were analyzed later ([DH86], [FK98], [ADH89]).

Related mechanism: “conspiracy of wells” — nearby attractive wells can bind even if each alone is subcritical. ([KS79])

Recently: geometric perturbations of soft waveguides [EV24], and curved arrays of symmetric wells [Ex24], also produce discrete spectrum.

Our question here: can *much weaker* local shifts (of finitely many wells) already produce eigenvalues at the band bottom?

Introduction

Introduction


We consider an infinite array of disjoint potential wells in any dimension $\nu \geq 2$ and without any symmetry.

Introduction

We consider an infinite array of disjoint potential wells in any dimension $\nu \geq 2$ and without any symmetry. Local perturbations consist of shifting finitely many of the wells (preserving disjointness)

Introduction

We consider an infinite array of disjoint potential wells in any dimension $\nu \geq 2$ and without any symmetry. Local perturbations consist of shifting finitely many of the wells (preserving disjointness)



The Model

Formally, we consider

The Model

Formally, we consider

$$H_{V,Y} = -\Delta + \sum_j V_j(x), \text{ on } H^2(\mathbb{R}^\nu),$$

The Model

Formally, we consider

$$H_{V,Y} = -\Delta + \sum_j V_j(x), \text{ on } H^2(\mathbb{R}^\nu),$$

where the set Y encodes the positions of the centers of the wells.

The Model

Formally, we consider

$$H_{V,Y} = -\Delta + \sum_j V_j(x), \text{ on } H^2(\mathbb{R}^\nu),$$

where the set Y encodes the positions of the centers of the wells.

The wells are: real-valued nonnegative $V \in L^p(\Sigma_{\rho,R}(0))$, where $p = 2$ for $\nu = 2, 3$ and $p > \frac{1}{2}\nu$ for $\nu \geq 4$

The Model

Formally, we consider

$$H_{V,Y} = -\Delta + \sum_j V_j(x), \text{ on } H^2(\mathbb{R}^\nu),$$

where the set Y encodes the positions of the centers of the wells.

The wells are: real-valued nonnegative $V \in L^p(\Sigma_{\rho,R}(0))$, where $p = 2$ for $\nu = 2, 3$ and $p > \frac{1}{2}\nu$ for $\nu \geq 4$ and $\Sigma_{\rho,R}(0) = (-\rho, \rho) \times B_R(0)$.

The Model

Formally, we consider

$$H_{V,Y} = -\Delta + \sum_j V_j(x), \text{ on } H^2(\mathbb{R}^\nu),$$

where the set Y encodes the positions of the centers of the wells.

The wells are: real-valued nonnegative $V \in L^p(\Sigma_{\rho,R}(0))$, where $p = 2$ for $\nu = 2, 3$ and $p > \frac{1}{2}\nu$ for $\nu \geq 4$ and $\Sigma_{\rho,R}(0) = (-\rho, \rho) \times B_R(0)$.

The array is constructed by shifting V

The Model

Formally, we consider

$$H_{V,Y} = -\Delta + \sum_j V_j(x), \text{ on } H^2(\mathbb{R}^\nu),$$

where the set Y encodes the positions of the centers of the wells.

The wells are: real-valued nonnegative $V \in L^p(\Sigma_{\rho,R}(0))$, where $p = 2$ for $\nu = 2, 3$ and $p > \frac{1}{2}\nu$ for $\nu \geq 4$ and $\Sigma_{\rho,R}(0) = (-\rho, \rho) \times B_R(0)$.

The array is constructed by shifting V , i.e.

$$V_j : x \mapsto V(x - y_j), \text{ with } \Sigma_{\rho,R}(y_i) := \Sigma_{\rho,R}(0) + y_i$$

Properties of the unperturbed H_{V,Y_0}

H_{V,Y_0} is self-adjoint, bounded from below

Properties of the unperturbed H_{V,Y_0}

H_{V,Y_0} is self-adjoint, bounded from below and with
 $\sigma(H_{V,Y_0}) = \sigma_{\text{ess}}(H_{V,Y_0})$

Properties of the unperturbed H_{V,Y_0}

H_{V,Y_0} is self-adjoint, bounded from below and with $\sigma(H_{V,Y_0}) = \sigma_{\text{ess}}(H_{V,Y_0})$, where $[0, \infty) \subset \sigma(H_{V,Y_0})$ and $\inf \sigma(H_{V,Y_0}) < 0$.

Properties of the unperturbed H_{V,Y_0}

H_{V,Y_0} is self-adjoint, bounded from below and with $\sigma(H_{V,Y_0}) = \sigma_{ess}(H_{V,Y_0})$, where $[0, \infty) \subset \sigma(H_{V,Y_0})$ and $\inf \sigma(H_{V,Y_0}) < 0$.

In the negative part there may or may not be gaps, their number is finite and does not exceed $\#\sigma_{disc}(H_V)$.

Properties of the perturbed $H_{V,\gamma}$

Properties of the perturbed $H_{V,Y}$

Proposition (3.1.)

$H_{V,Y}$ is self-adjoint on $H^2(\mathbb{R}^\nu)$ for any local perturbation and $\sigma_{\text{ess}}(H_{V,Y}) = \sigma_{\text{ess}}(H_{V,Y_0})$.

Properties of the perturbed $H_{V,Y}$

Proposition (3.1.)

$H_{V,Y}$ is self-adjoint on $H^2(\mathbb{R}^\nu)$ for any local perturbation and $\sigma_{\text{ess}}(H_{V,Y}) = \sigma_{\text{ess}}(H_{V,Y_0})$.

- The perturbation is well-defined
- Its essential spectrum is stable

Main result

Eigenvalues at the bottom of the spectrum:

Main result

Eigenvalues at the bottom of the spectrum: If we shift purely in the horizontal direction (or in higher dimension orthogonal to the axis of Y)

Main result

Eigenvalues at the bottom of the spectrum: If we shift purely in the horizontal direction (or in higher dimension orthogonal to the axis of Y) the spectral threshold is preserved.

Main result

If we shift purely in the longitudinal direction

Main result

If we shift purely in the longitudinal direction we always* generate discrete eigenvalues below the threshold.

Main result

If we shift purely in the longitudinal direction we always* generate discrete eigenvalues below the threshold.

*provided $R \leq \rho\sqrt{\nu-1}$.

Main result


What about combined movement?

Main result

What about combined movement? Numerics

Main result

What about combined movement? Numerics



Proof Outline

Proof Outline

Let κ_0 be $-\kappa_0^2 = \inf \sigma(H_{V, \gamma_0})$

Proof Outline

Let κ_0 be $-\kappa_0^2 = \inf \sigma(H_{V,Y_0})$ it is then enough to check that there is a $\kappa > \kappa_0$ s.t. $1 \in \sigma(K_{V,Y}(-\kappa^2))$

Proof Outline

Let κ_0 be $-\kappa_0^2 = \inf \sigma(H_{V,Y_0})$ it is then enough to check that there is a $\kappa > \kappa_0$ s.t. $1 \in \sigma(K_{V,Y}(-\kappa^2))$, which happens precisely when $\sup \sigma(K_{V,Y}(-\kappa_0^2)) > 1$.

Proof Outline

Let κ_0 be $-\kappa_0^2 = \inf \sigma(H_{V,Y_0})$ it is then enough to check that there is a $\kappa > \kappa_0$ s.t. $1 \in \sigma(K_{V,Y}(-\kappa^2))$, which happens precisely when $\sup \sigma(K_{V,Y}(-\kappa_0^2)) > 1$.

The goal:

Proof Outline

Let κ_0 be $-\kappa_0^2 = \inf \sigma(H_{V,Y_0})$ it is then enough to check that there is a $\kappa > \kappa_0$ s.t. $1 \in \sigma(K_{V,Y}(-\kappa^2))$, which happens precisely when $\sup \sigma(K_{V,Y}(-\kappa_0^2)) > 1$.

The goal: Find a trial function $\phi \in L^2(\text{supp } V_Y)$ s.t.

Proof Outline

Let κ_0 be $-\kappa_0^2 = \inf \sigma(H_{V,Y_0})$ it is then enough to check that there is a $\kappa > \kappa_0$ s.t. $1 \in \sigma(K_{V,Y}(-\kappa^2))$, which happens precisely when $\sup \sigma(K_{V,Y}(-\kappa_0^2)) > 1$.

The goal: Find a trial function $\phi \in L^2(\text{supp } V_Y)$ s.t.

$$(\phi, K_{V,Y}(-\kappa_0^2)\phi) > \|\phi\|^2$$

Proof Outline

Trial function construction:

Proof Outline

Trial function construction:

Take the generalized eigenfunction ψ_0 corresponding to the bottom of the essential spectrum of H_{V, γ_0} .

Proof Outline

Trial function construction:

Take the generalized eigenfunction ψ_0 corresponding to the bottom of the essential spectrum of H_{V,Y_0} . The corresponding gen. eigenfunction for K_{V,Y_0} is $\phi_0 = V_{Y_0}^{1/2} \psi_0$.

Proof Outline

Trial function construction:

Take the generalized eigenfunction ψ_0 corresponding to the bottom of the essential spectrum of H_{V,Y_0} . The corresponding gen. eigenfunction for K_{V,Y_0} is $\phi_0 = V_{Y_0}^{1/2} \psi_0$. We proceed with $\phi_{0,j}(\xi + y_j) := \phi_0(\xi)$, where $\{\phi_{0,j} \in \sum_{j \in \mathbb{Z}}^{\oplus} L^2(\Sigma_{\rho,R}(y_j))\}$.

Proof Outline

Trial function construction:

Take the generalized eigenfunction ψ_0 corresponding to the bottom of the essential spectrum of H_{V,Y_0} . The corresponding gen. eigenfunction for K_{V,Y_0} is $\phi_0 = V_{Y_0}^{1/2} \psi_0$. We proceed with $\phi_{0,j}(\xi + y_j) := \phi_0(\xi)$, where $\{\phi_{0,j} \in \sum_{j \in \mathbb{Z}}^{\oplus} L^2(\Sigma_{\rho,R}(y_j))\}$.

But now ϕ_0 is not yet in $L^2(\text{supp } V_Y)$.

Proof Outline

Trial function construction:

Take the generalized eigenfunction ψ_0 corresponding to the bottom of the essential spectrum of H_{V,Y_0} . The corresponding gen. eigenfunction for K_{V,Y_0} is $\phi_0 = V_{Y_0}^{1/2} \psi_0$. We proceed with $\phi_{0,j}(\xi + y_j) := \phi_0(\xi)$, where $\{\phi_{0,j} \in \sum_{j \in \mathbb{Z}}^{\oplus} L^2(\Sigma_{\rho,R}(y_j))\}$.

But now ϕ_0 is not yet in $L^2(\text{supp } V_Y)$.

Mollifier: $h_n(x) = \sum_i h_{n,i} \chi_{\Sigma_{\rho,R}(y_i)}(x)$, $h_{n,i} = \frac{n^2}{n^2 + i^2}$.

Proof Outline

Trial function construction:

Take the generalized eigenfunction ψ_0 corresponding to the bottom of the essential spectrum of H_{V,Y_0} . The corresponding gen. eigenfunction for K_{V,Y_0} is $\phi_0 = V_{Y_0}^{1/2} \psi_0$. We proceed with $\phi_{0,j}(\xi + y_j) := \phi_0(\xi)$, where $\{\phi_{0,j} \in \sum_{j \in \mathbb{Z}}^{\oplus} L^2(\Sigma_{\rho,R}(y_j))\}$.

But now ϕ_0 is not yet in $L^2(\text{supp } V_Y)$.

Mollifier: $h_n(x) = \sum_i h_{n,i} \chi_{\Sigma_{\rho,R}(y_i)}(x)$, $h_{n,i} = \frac{n^2}{n^2 + i^2}$.

The effect of the mollifier is arbitrarily small:

Proof Outline

Trial function construction:

Take the generalized eigenfunction ψ_0 corresponding to the bottom of the essential spectrum of H_{V,Y_0} . The corresponding gen. eigenfunction for K_{V,Y_0} is $\phi_0 = V_{Y_0}^{1/2} \psi_0$. We proceed with $\phi_{0,j}(\xi + y_j) := \phi_0(\xi)$, where $\{\phi_{0,j} \in \sum_{j \in \mathbb{Z}}^{\oplus} L^2(\Sigma_{\rho,R}(y_j))\}$.

But now ϕ_0 is not yet in $L^2(\text{supp } V_Y)$.

Mollifier: $h_n(x) = \sum_i h_{n,i} \chi_{\Sigma_{\rho,R}(y_i)}(x)$, $h_{n,i} = \frac{n^2}{n^2 + i^2}$.

The effect of the mollifier is arbitrarily small:

Lemma

$$\langle h_n \phi_0, K_{V,Y_0}(-\kappa_0^2) h_n \phi_0 \rangle - \|h_n \phi_0\|^2 = O(n^{-2}) \rightarrow 0.$$

Proof Outline

Now we have:

Proof Outline

Now we have:

$$(h_n \phi_0, K_{V,Y}(-\kappa_0^2) h_n \phi_0) - \|h_n \phi_0\|^2 > 0,$$

holds for n large enough.

Proof Outline

Now we have:

$$(h_n\phi_0, K_{V,Y}(-\kappa_0^2)h_n\phi_0) - \|h_n\phi_0\|^2 > 0,$$

holds for n large enough. We can add and subtract $(h_n\phi_0, K_{V,Y_0}(-\kappa_0^2)h_n\phi_0)$ to the LHS and use the lemma to get

Proof Outline

Now we have:

$$(h_n\phi_0, K_{V,Y}(-\kappa_0^2)h_n\phi_0) - \|h_n\phi_0\|^2 > 0,$$

holds for n large enough. We can add and subtract $(h_n\phi_0, K_{V,Y_0}(-\kappa_0^2)h_n\phi_0)$ to the LHS and use the lemma to get

$$\lim_{n \rightarrow \infty} (h_n\phi_0, K_{V,Y}(-\kappa_0^2)h_n\phi_0) - (h_n\phi_0, K_{V,Y_0}(-\kappa_0^2)h_n\phi_0) > 0.$$

Proof Outline

Now we have:

$$(h_n \phi_0, K_{V,Y}(-\kappa_0^2) h_n \phi_0) - \|h_n \phi_0\|^2 > 0,$$

holds for n large enough. We can add and subtract

$(h_n \phi_0, K_{V,Y_0}(-\kappa_0^2) h_n \phi_0)$ to the LHS and use the lemma to get

$$\lim_{n \rightarrow \infty} (h_n \phi_0, K_{V,Y}(-\kappa_0^2) h_n \phi_0) - (h_n \phi_0, K_{V,Y_0}(-\kappa_0^2) h_n \phi_0) > 0.$$

Due to the local character of the perturbation and some technicalities we have

$$\sum_{i,j \in \mathbb{Z}} \left(\phi_0, \left[K_{V,Y}^{(i,j)}(-\kappa_0^2) - K_{V,Y_0}^{(i,j)}(-\kappa_0^2) \right] \phi_0 \right).$$

Proof Outline

We have the perturbation shift along the first axis

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Further the relative shifts of the neighbouring points $\delta_j = (y_{j+1} - y_j)_1 - a$;

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Further the relative shifts of the neighbouring points $\delta_j = (y_{j+1} - y_j)_1 - a$;
Then

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Further the relative shifts of the neighbouring points $\delta_j = (y_{j+1} - y_j)_1 - a$;
Then

$$\eta_{ij} = \sum_{r=\min(i,j)}^{\max(i,j)-1} \delta_r$$

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Further the relative shifts of the neighbouring points $\delta_j = (y_{j+1} - y_j)_1 - a$;
Then

$$\eta_{ij} = \sum_{r=\min(i,j)}^{\max(i,j)-1} \delta_r$$

This quantity has zero mean

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Further the relative shifts of the neighbouring points $\delta_j = (y_{j+1} - y_j)_1 - a$;
Then

$$\eta_{ij} = \sum_{r=\min(i,j)}^{\max(i,j)-1} \delta_r$$

This quantity has zero mean

$$\sum_{i,j \in \mathbb{Z}} \eta_{i,j} = 0.$$

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Further the relative shifts of the neighbouring points $\delta_j = (y_{j+1} - y_j)_1 - a$;
Then

$$\eta_{ij} = \sum_{r=\min(i,j)}^{\max(i,j)-1} \delta_r$$

This quantity has zero mean

$$\sum_{i,j \in \mathbb{Z}} \eta_{i,j} = 0.$$

This suggests to employ Jensen inequality!

Proof Outline

We have the perturbation shift along the first axis

$$\eta_{ij} = (y_i - y_i^{(0)} - y_j + y_j^{(0)})_1$$

Further the relative shifts of the neighbouring points $\delta_j = (y_{j+1} - y_j)_1 - a$;
Then

$$\eta_{ij} = \sum_{r=\min(i,j)}^{\max(i,j)-1} \delta_r$$

This quantity has zero mean

$$\sum_{i,j \in \mathbb{Z}} \eta_{i,j} = 0.$$

This suggests to employ Jensen inequality! But, we cannot use the convexity of the resolvent directly, since the perturbation enters its argument through Euclidian distance.

Proof Outline

Under the *non-optimal* condition $R \leq \rho\sqrt{\nu - 1}$, we have strict convexity.

Under the *non-optimal* condition $R \leq \rho\sqrt{\nu - 1}$, we have strict convexity.

We can thus use, again after some technicalities, the Jensen inequality

Under the *non-optimal* condition $R \leq \rho\sqrt{\nu-1}$, we have strict convexity.

We can thus use, again after some technicalities, the Jensen inequality and by assumption for at least some i, j , $\eta_{ij} \neq 0$

Under the *non-optimal* condition $R \leq \rho\sqrt{\nu-1}$, we have strict convexity.

We can thus use, again after some technicalities, the Jensen inequality and by assumption for at least some i, j , $\eta_{ij} \neq 0$, thus we have positivity.

Conclusion

- Studied local perturbations of a straight, equidistant array of disjoint wells in \mathbb{R}^ν ($\nu \geq 2$).

Conclusion

- Studied local perturbations of a straight, equidistant array of disjoint wells in \mathbb{R}^ν ($\nu \geq 2$).
- Main result: any local perturbation creates discrete eigenvalues below the threshold

Conclusion

- Studied local perturbations of a straight, equidistant array of disjoint wells in \mathbb{R}^ν ($\nu \geq 2$).
- Main result: any local perturbation creates discrete eigenvalues below the threshold
- The ratio condition $R \leq \rho\sqrt{\nu-1}$ used to preserve convexity.

Conclusion

- Studied local perturbations of a straight, equidistant array of disjoint wells in \mathbb{R}^ν ($\nu \geq 2$).
- Main result: any local perturbation creates discrete eigenvalues below the threshold
- The ratio condition $R \leq \rho\sqrt{\nu-1}$ used to preserve convexity.
- This condition is *sufficient but not necessary* (likely proof artifact; estimates are rough).

Conclusion

- Studied local perturbations of a straight, equidistant array of disjoint wells in \mathbb{R}^ν ($\nu \geq 2$).
- Main result: any local perturbation creates discrete eigenvalues below the threshold
- The ratio condition $R \leq \rho\sqrt{\nu-1}$ used to preserve convexity.
- This condition is *sufficient but not necessary* (likely proof artifact; estimates are rough).
- Possible goals are to relax aspect-ratio bound (find optimal bound for convexity); allow decaying (non-compact) shifts

References I



P. Exner, D. Spitzkopf: Local perturbations of potential well arrays, to appear in *Journal of Spectral Theory*



V.A. Zheludev: Perturbation of the spectrum of the one-dimensional Schrödinger operator with a periodic potential, in *Topics in Mathematical Physics, vol. 4* (M.Sh. Birman, ed.), Consultants Bureau, New York 1971; pp. 55–75.



S. Fassari: Spectral properties of the Kronig-Penney Hamiltonian with a localised impurity, *J. Math. Phys.* **30** (1989), 1385–1392.



F. Gesztesy, B. Simon: A short proof of Zheludev's theorem, *Trans. Am. Math. Soc.* **335** (1991), 329–340.



P.A. Deift, R. Hempel: On the existence of eigenvalues of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$, *Commun. Math. Phys.* **103** (1986), 461–490.



S. Fassari, M. Klaus: Coupling constant thresholds of perturbed periodic Hamiltonians, *J. Math. Phys.* **39** (1998), 4369–4416.



S. Alama, P.A. Deift, R. Hempel: Eigenvalue branches of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$, *Commun. Math. Phys.* **121** (1989), 291–321.

References II



M. Klaus, B. Simon: Binding of Schrödinger particles through conspiracy of potential wells, *Ann. Inst. H. Poincaré*, **A30** 1979, 83–87.



P. Exner: Geometry effects in quantum dot families, *Pure Appl. Funct. Anal.* **9** (2024), 1065–1080.



P. Exner, S. Vugalter: Bound states in bent soft waveguides, *J. Spect. Theory* **14** (2024), 427–457.