

# On the $C^*$ -algebraic rigidity of Heisenberg groups

Daniel Belitiță  
Institute of Mathematics of the Romanian Academy  
\*\*\*

Analytic and Algebraic Methods in Physics XXII

Prague, 27.08.2025



# References



Ingrid Beltiță, D. B.,  *$C^*$ -rigidity of the Heisenberg group*. Publications of the Research Institute for Mathematical Sciences RIMS 61 (2025), no. 3, 487-502.



Ingrid Beltiță, D. B., *Dual rigidity of the Heisenberg groups*. Preprint arXiv: 2508.08904 [math.RT].

$$[X_j, X_k] = 0, \quad [P_j, P_k] = 0, \quad [X_j, P_k] = i\delta_{jk}$$

$$[X_j, X_k] = 0, [P_j, P_k] = 0, [X_j, P_k] = i\delta_{jk}$$

- *Heisenberg(-Weyl) Lie algebra/group:*

$$\mathfrak{h}_n = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}, [(q, p, t), (q', p', t')] = [(0, 0, \langle p, q' \rangle - \langle p', q \rangle)]$$

$$H_n = (\mathfrak{h}_n, \cdot), x \cdot y := x + y + \frac{1}{2}[x, y]$$

$$[X_j, X_k] = 0, [P_j, P_k] = 0, [X_j, P_k] = i\delta_{jk}$$

- *Heisenberg(-Weyl) Lie algebra/group:*

$$\mathfrak{h}_n = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}, [(q, p, t), (q', p', t')] = [(0, 0, \langle p, q' \rangle - \langle p', q \rangle)]$$

$$H_n = (\mathfrak{h}_n, \cdot), x \cdot y := x + y + \frac{1}{2}[x, y]$$

- irreducible unitary representation  $\pi_\lambda: H_n \rightarrow U(L^2(\mathbb{R}^n))$  for  $\lambda \in \mathbb{R} \setminus \{0\}$ ,

$$\pi_\lambda(q, p, t)f(x) = e^{i\lambda(\langle p, x \rangle + \frac{1}{2}\langle p, q \rangle + t)}f(q + x)$$

$$[X_j, X_k] = 0, [P_j, P_k] = 0, [X_j, P_k] = i\delta_{jk}$$

- *Heisenberg(-Weyl) Lie algebra/group:*

$$\mathfrak{h}_n = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}, [(q, p, t), (q', p', t')] = [(0, 0, \langle p, q' \rangle - \langle p', q \rangle)]$$

$$H_n = (\mathfrak{h}_n, \cdot), x \cdot y := x + y + \frac{1}{2}[x, y]$$

- irreducible unitary representation  $\pi_\lambda: H_n \rightarrow U(L^2(\mathbb{R}^n))$  for  $\lambda \in \mathbb{R} \setminus \{0\}$ ,

$$\pi_\lambda(q, p, t)f(x) = e^{i\lambda(\langle p, x \rangle + \frac{1}{2}\langle p, q \rangle + t)}f(q + x)$$

$$\leadsto \frac{1}{i}d\pi_\lambda(1_k, 0, 0)f(x) = \underbrace{\frac{1}{i}\frac{\partial f}{\partial q_k}}_{P_k f}, \quad \frac{1}{i}d\pi_\lambda(0, 1_j, 0)f(q) = \underbrace{\lambda q_j f(q)}_{X_j f}$$

$$[X_j, X_k] = 0, [P_j, P_k] = 0, [X_j, P_k] = i\delta_{jk}$$

- *Heisenberg(-Weyl) Lie algebra/group:*

$$\mathfrak{h}_n = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}, [(q, p, t), (q', p', t')] = [(0, 0, \langle p, q' \rangle - \langle p', q \rangle)]$$

$$H_n = (\mathfrak{h}_n, \cdot), x \cdot y := x + y + \frac{1}{2}[x, y]$$

- irreducible unitary representation  $\pi_\lambda: H_n \rightarrow U(L^2(\mathbb{R}^n))$  for  $\lambda \in \mathbb{R} \setminus \{0\}$ ,

$$\pi_\lambda(q, p, t)f(x) = e^{i\lambda(\langle p, x \rangle + \frac{1}{2}\langle p, q \rangle + t)}f(q + x)$$

$$\leadsto \frac{1}{i}d\pi_\lambda(1_k, 0, 0)f(x) = \underbrace{\frac{1}{i}\frac{\partial f}{\partial q_k}}_{P_k f}, \quad \frac{1}{i}d\pi_\lambda(0, 1_j, 0)f(q) = \underbrace{\lambda q_j f(q)}_{X_j f}$$

## Problem on the Canonical Commutation Relations (CCR)

Is the “canonical host group” in some sense **unique**?

# The unitary dual space of the Heisenberg group $H_n$



# The unitary dual space of the Heisenberg group $H_n$



The CCR (resp., with  $[\cdot, \cdot]$  replaced by a Poisson bracket) arise in representations (resp., on coadjoint orbits) of many Lie groups. For the uniqueness problem we need *all* unitary irreducible representations, along with the topology on their set of equivalence classes.

# The unitary dual space of the Heisenberg group $H_n$



The CCR (resp., with  $[\cdot, \cdot]$  replaced by a Poisson bracket) arise in representations (resp., on coadjoint orbits) of many Lie groups. For the uniqueness problem we need *all* unitary irreducible representations, along with the topology on their set of equivalence classes.

- Besides the  $\infty$ -dim. representations  $\pi_\lambda: H_n \rightarrow U(L^2(\mathbb{R}^n))$ , there are the 1-dim. representations  $\chi_{a,b}: H_n \rightarrow U(1)$  for  $(a, b) \in \mathbb{R}^n \times \mathbb{R}^n$ ,  
$$\chi_{a,b}(q, p, t) := e^{i(a \cdot p + b \cdot q)}$$

# The unitary dual space of the Heisenberg group $H_n$



The CCR (resp., with  $[\cdot, \cdot]$  replaced by a Poisson bracket) arise in representations (resp., on coadjoint orbits) of many Lie groups. For the uniqueness problem we need *all* unitary irreducible representations, along with the topology on their set of equivalence classes.

- Besides the  $\infty$ -dim. representations  $\pi_\lambda: H_n \rightarrow U(L^2(\mathbb{R}^n))$ , there are the 1-dim. representations  $\chi_{a,b}: H_n \rightarrow U(1)$  for  $(a, b) \in \mathbb{R}^n \times \mathbb{R}^n$ ,

$$\chi_{a,b}(q, p, t) := e^{i(a \cdot p + b \cdot q)}$$

- $\widehat{H}_n = \{[\pi_\lambda] \mid \lambda \in \mathbb{R} \setminus \{0\}\} \cup \{\chi_{a,b} \mid (a, b) \in \mathbb{R}^n \times \mathbb{R}^n\} \simeq (\mathbb{R} \setminus \{0\}) \sqcup \mathbb{R}^{2n}$  endowed with the (non-separated!) topology given by the condition:

If  $\lambda_n \xrightarrow{n \rightarrow \infty} 0$  in  $\mathbb{R} \setminus \{0\}$ , then  $[\pi_{\lambda_n}] \xrightarrow{n \rightarrow \infty} \chi_{a,b}$  in  $\widehat{H}_n$  for all  $(a, b) \in \mathbb{R}^{2n}$

# Direct and inverse problems in representation theory

# Direct and inverse problems in representation theory

- $G$  is a Lie group (i.e., group with a compatible smooth structure)

**Recall:** The Lie groups form a full subcategory of **topological** groups, since every continuous morphism of Lie groups is smooth.

# Direct and inverse problems in representation theory

- $G$  is a Lie group (i.e., group with a compatible smooth structure)

**Recall:** The Lie groups form a full subcategory of **topological** groups, since every continuous morphism of Lie groups is smooth.

- $\widehat{G}$  = equivalence classes  $[\pi]$  of irreducible unitary repres.  $\pi: G \rightarrow U(\mathcal{H}_\pi)$

**Recall:**  $G \times \mathcal{H}_\pi \rightarrow \mathcal{H}_\pi$ ,  $(g, v) \mapsto \pi(g)v$  is a continuous group action; no nontrivial invariant subspaces

# Direct and inverse problems in representation theory

- $G$  is a Lie group (i.e., group with a compatible smooth structure)

**Recall:** The Lie groups form a full subcategory of **topological** groups, since every continuous morphism of Lie groups is smooth.

- $\widehat{G}$  = equivalence classes  $[\pi]$  of irreducible unitary repres.  $\pi: G \rightarrow U(\mathcal{H}_\pi)$

**Recall:**  $G \times \mathcal{H}_\pi \rightarrow \mathcal{H}_\pi$ ,  $(g, v) \mapsto \pi(g)v$  is a continuous group action; no nontrivial invariant subspaces

- Topology of  $\widehat{G}$  is defined by convergence of representation coefficients  $c_{vu} := \langle v, \pi(\cdot)u \rangle \in \mathcal{C}(G, \mathbb{C})$ ,  $u, v \in \mathcal{H}_\pi$ , on the compact subsets of  $G$ .

**E.g.**, if  $\mathcal{H}_\pi = \mathbb{C}^n$ , then  $\pi(\cdot) = (c_{jk}(\cdot))_{j,k}: G \rightarrow U(n)$ .

# Direct and inverse problems in representation theory

- $G$  is a Lie group (i.e., group with a compatible smooth structure)

**Recall:** The Lie groups form a full subcategory of **topological** groups, since every continuous morphism of Lie groups is smooth.

- $\hat{G}$  = equivalence classes  $[\pi]$  of irreducible unitary reps.  $\pi: G \rightarrow U(\mathcal{H}_\pi)$

**Recall:**  $G \times \mathcal{H}_\pi \rightarrow \mathcal{H}_\pi$ ,  $(g, v) \mapsto \pi(g)v$  is a continuous group action; no nontrivial invariant subspaces

- Topology of  $\hat{G}$  is defined by convergence of representation coefficients  $c_{vu} := \langle v, \pi(\cdot)u \rangle \in \mathcal{C}(G, \mathbb{C})$ ,  $u, v \in \mathcal{H}_\pi$ , on the compact subsets of  $G$ .

**E.g.**, if  $\mathcal{H}_\pi = \mathbb{C}^n$ , then  $\pi(\cdot) = (c_{jk}(\cdot))_{j,k}: G \rightarrow U(n)$ .

**Direct Problem** (  $G \rightsquigarrow \hat{G}$  ). Describe  $\hat{G}$  in terms of  $G$  itself.



# Direct and inverse problems in representation theory

- $G$  is a Lie group (i.e., group with a compatible smooth structure)

**Recall:** The Lie groups form a full subcategory of **topological** groups, since every continuous morphism of Lie groups is smooth.

- $\widehat{G}$  = equivalence classes  $[\pi]$  of irreducible unitary repres.  $\pi: G \rightarrow U(\mathcal{H}_\pi)$

**Recall:**  $G \times \mathcal{H}_\pi \rightarrow \mathcal{H}_\pi$ ,  $(g, v) \mapsto \pi(g)v$  is a continuous group action; no nontrivial invariant subspaces

- Topology of  $\widehat{G}$  is defined by convergence of representation coefficients  $c_{vu} := \langle v, \pi(\cdot)u \rangle \in \mathcal{C}(G, \mathbb{C})$ ,  $u, v \in \mathcal{H}_\pi$ , on the compact subsets of  $G$ .

**E.g.**, if  $\mathcal{H}_\pi = \mathbb{C}^n$ , then  $\pi(\cdot) = (c_{jk}(\cdot))_{j,k}: G \rightarrow U(n)$ .

**Direct Problem** (  $G \rightsquigarrow \widehat{G}$  ). Describe  $\widehat{G}$  in terms of  $G$  itself.

**Inverse Problem** (  $G \longleftarrow \widehat{G}$  ). Can  $G$  be recovered from  $\widehat{G}$ ?

# The Inverse Problem ( $G \overset{\sim}{\leftarrow} \widehat{G}$ ) for commutative groups

# The Inverse Problem ( $G \rightsquigarrow \widehat{G}$ ) for commutative groups

If  $G$  is **commutative**, then  $\widehat{G} \simeq \text{Hom}(G, \text{U}(1))$  (again a **group**)

# The Inverse Problem ( $G \rightsquigarrow \widehat{G}$ ) for commutative groups

If  $G$  is **commutative**, then  $\widehat{G} \simeq \text{Hom}(G, \text{U}(1))$  (again a **group**) and  $G \xrightarrow{\sim} \widehat{\widehat{G}}$  as topological **groups** (Pontryagin duality).

# The Inverse Problem ( $G \xleftarrow{\sim} \widehat{\widehat{G}}$ ) for commutative groups

If  $G$  is **commutative**, then  $\widehat{G} \simeq \text{Hom}(G, \text{U}(1))$  (again a **group**) and  $G \xrightarrow{\sim} \widehat{\widehat{G}}$  as topological **groups** (Pontryagin duality).

If  $G_n := (\mathbb{R}^n, +)$  then  $(\mathbb{R}^n, +) \xrightarrow{\sim} (\widehat{G_n}, \cdot)$ ,  $\lambda \mapsto e^{i\langle \lambda, \bullet \rangle}$  hence

# The Inverse Problem ( $G \rightsquigarrow \widehat{G}$ ) for commutative groups

If  $G$  is **commutative**, then  $\widehat{G} \simeq \text{Hom}(G, \text{U}(1))$  (again a **group**) and  $G \xrightarrow{\sim} \widehat{\widehat{G}}$  as topological **groups** (Pontryagin duality).

If  $G_n := (\mathbb{R}^n, +)$  then  $(\mathbb{R}^n, +) \xrightarrow{\sim} (\widehat{G}_n, \cdot)$ ,  $\lambda \mapsto e^{i\langle \lambda, \bullet \rangle}$  hence  $G_n \simeq G_m$  as Lie groups (i.e.,  $n = m$ )

$\Updownarrow$


$\widehat{G}_n \simeq \widehat{G}_m$  as locally compact **topological spaces** 

# The Inverse Problem ( $G \xleftarrow{\quad} \widehat{G}$ ) for commutative groups

If  $G$  is **commutative**, then  $\widehat{G} \simeq \text{Hom}(G, \text{U}(1))$  (again a **group**) and  $G \xrightarrow{\sim} \widehat{\widehat{G}}$  as topological **groups** (Pontryagin duality).

If  $G_n := (\mathbb{R}^n, +)$  then  $(\mathbb{R}^n, +) \xrightarrow{\sim} (\widehat{G}_n, \cdot)$ ,  $\lambda \mapsto e^{i\langle \lambda, \cdot \rangle}$  hence  
 $G_n \simeq G_m$  as Lie groups (i.e.,  $n = m$ )



$\widehat{G}_n \simeq \widehat{G}_m$  as locally compact **topological spaces** 




$\mathcal{C}_0(\widehat{G}_n) \simeq \mathcal{C}_0(\widehat{G}_m)$  as **commutative**  $C^*$ -algebras

# The Inverse Problem ( $G \rightsquigarrow \widehat{G}$ ) for commutative groups

If  $G$  is **commutative**, then  $\widehat{G} \simeq \text{Hom}(G, \text{U}(1))$  (again a **group**) and  $G \xrightarrow{\sim} \widehat{\widehat{G}}$  as topological **groups** (Pontryagin duality).

If  $G_n := (\mathbb{R}^n, +)$  then  $(\mathbb{R}^n, +) \xrightarrow{\sim} (\widehat{G}_n, \cdot)$ ,  $\lambda \mapsto e^{i\langle \lambda, \bullet \rangle}$  hence  $G_n \simeq G_m$  as Lie groups (i.e.,  $n = m$ )

$\widehat{G}_n \simeq \widehat{G}_m$  as locally compact **topological spaces** 

$\mathcal{C}_0(\widehat{G}_n) \simeq \mathcal{C}_0(\widehat{G}_m)$  as **commutative**  $C^*$ -algebras

This suggests that **IP** for noncommutative Lie groups may be approached via **noncommutative topology**, that is,  **$C^*$ -algebras**.



# Group $C^*$ -algebras

# Group $C^*$ -algebras

- $C^*$ -algebra  $\mathcal{A}$ : associative Banach  $*$ -algebra/ $\mathbb{C}$ ,  $\|a^*a\| = \|a\|^2$  for  $a \in \mathcal{A}$   
Equivalently: subalgebra  $\mathcal{A} = \overline{\mathcal{A}}^{\|\cdot\|} \subseteq \mathcal{B}(\mathcal{H})$  with  $a \in \mathcal{A} \Leftrightarrow a^* \in \mathcal{A}$
- $\widehat{\mathcal{A}}$  the set of all equivalence classes  $[\pi]$  of irred.  $*$ -repres.  $\pi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H}_\pi)$

# Group $C^*$ -algebras

- $C^*$ -algebra  $\mathcal{A}$ : associative Banach  $*$ -algebra/ $\mathbb{C}$ ,  $\|a^*a\| = \|a\|^2$  for  $a \in \mathcal{A}$   
Equivalently: subalgebra  $\mathcal{A} = \overline{\mathcal{A}^{\|\cdot\|}} \subseteq \mathcal{B}(\mathcal{H})$  with  $a \in \mathcal{A} \Leftrightarrow a^* \in \mathcal{A}$
- $\widehat{\mathcal{A}}$  the set of all equivalence classes  $[\pi]$  of irred.  $*$ -repres.  $\pi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H}_\pi)$
- $G$  Lie group

# Group $C^*$ -algebras

- $C^*$ -algebra  $\mathcal{A}$ : associative Banach  $*$ -algebra/ $\mathbb{C}$ ,  $\|a^*a\| = \|a\|^2$  for  $a \in \mathcal{A}$   
Equivalently: subalgebra  $\mathcal{A} = \overline{\mathcal{A}^{\|\cdot\|}} \subseteq \mathcal{B}(\mathcal{H})$  with  $a \in \mathcal{A} \Leftrightarrow a^* \in \mathcal{A}$
- $\widehat{\mathcal{A}}$  the set of all equivalence classes  $[\pi]$  of irred.  $*$ -repres.  $\pi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H}_\pi)$
- $G$  Lie group
- $\pi: G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$  repres.  $\leadsto \pi: \mathcal{C}_c(G) \rightarrow \mathcal{B}(\mathcal{H}_\pi)$ ,  $\pi(f) = \int_G f(x)\pi(x)dx$

Here  $\pi(f)\pi(h) = \pi(f * h)$  and  $\pi(f)^* = \pi(f^*)$  with  $f * h, f^* \in \mathcal{C}_c(G)$  independent on  $\pi$ .

# Group $C^*$ -algebras

- $C^*$ -algebra  $\mathcal{A}$ : associative Banach  $*$ -algebra/ $\mathbb{C}$ ,  $\|a^*a\| = \|a\|^2$  for  $a \in \mathcal{A}$   
Equivalently: subalgebra  $\mathcal{A} = \overline{\mathcal{A}^{\|\cdot\|}} \subseteq \mathcal{B}(\mathcal{H})$  with  $a \in \mathcal{A} \Leftrightarrow a^* \in \mathcal{A}$
- $\widehat{\mathcal{A}}$  the set of all equivalence classes  $[\pi]$  of irred.  $*$ -repres.  $\pi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H}_\pi)$
- $G$  Lie group
- $\pi: G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$  repres.  $\leadsto \pi: \mathcal{C}_c(G) \rightarrow \mathcal{B}(\mathcal{H}_\pi)$ ,  $\pi(f) = \int_G f(x)\pi(x)dx$

Here  $\pi(f)\pi(h) = \pi(f * h)$  and  $\pi(f)^* = \pi(f^*)$  with  $f * h, f^* \in \mathcal{C}_c(G)$  independent on  $\pi$ .

- $C^*(G) := \overline{\mathcal{C}_c(G)}^{\|\cdot\|}$ , where  $\|f\| := \sup_{[\pi] \in \widehat{G}} \|\pi(f)\| \leq \int_G |f(x)|dx$

# Group $C^*$ -algebras

- $C^*$ -algebra  $\mathcal{A}$ : associative Banach  $*$ -algebra/ $\mathbb{C}$ ,  $\|a^*a\| = \|a\|^2$  for  $a \in \mathcal{A}$   
Equivalently: subalgebra  $\mathcal{A} = \overline{\mathcal{A}^{\|\cdot\|}} \subseteq \mathcal{B}(\mathcal{H})$  with  $a \in \mathcal{A} \Leftrightarrow a^* \in \mathcal{A}$
- $\widehat{\mathcal{A}}$  the set of all equivalence classes  $[\pi]$  of irred.  $*$ -repres.  $\pi: \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H}_\pi)$
- $G$  Lie group
- $\pi: G \rightarrow \mathcal{U}(\mathcal{H}_\pi)$  repres.  $\leadsto \pi: \mathcal{C}_c(G) \rightarrow \mathcal{B}(\mathcal{H}_\pi)$ ,  $\pi(f) = \int_G f(x)\pi(x)dx$

Here  $\pi(f)\pi(h) = \pi(f * h)$  and  $\pi(f)^* = \pi(f^*)$  with  $f * h, f^* \in \mathcal{C}_c(G)$  independent on  $\pi$ .

- $C^*(G) := \overline{\mathcal{C}_c(G)}^{\|\cdot\|}$ , where  $\|f\| := \sup_{[\pi] \in \widehat{G}} \|\pi(f)\| \leq \int_G |f(x)|dx$

**Fact:**  $\widehat{G} \simeq \widehat{C^*(G)}$  so the unitary dual  $\widehat{G}$  can be recovered from  $C^*(G)$

# $C^*$ -rigidity of Lie groups

# $C^*$ -rigidity of Lie groups

Given a class of Lie groups  $\mathcal{G}$  and  $G_0$  in  $\mathcal{G}$ .

$G_0$  is called  **$C^*$ -rigid within  $\mathcal{G}$**  if  $C^*(G_0) \simeq C^*(G) \Leftrightarrow G_0 \simeq G$  for all  $G$  in  $\mathcal{G}$



# $C^*$ -rigidity of Lie groups

Given a class of Lie groups  $\mathcal{G}$  and  $G_0$  in  $\mathcal{G}$ .

$G_0$  is called  **$C^*$ -rigid within  $\mathcal{G}$**  if  $C^*(G_0) \simeq C^*(G) \Leftrightarrow G_0 \simeq G$  for all  $G$  in  $\mathcal{G}$

Which groups are  $C^*$ -rigid within the class of **1-connected** Lie groups?

(i.e., connected and simply connected)

# $C^*$ -rigidity of Lie groups

Given a class of Lie groups  $\mathcal{G}$  and  $G_0$  in  $\mathcal{G}$ .

$G_0$  is called  **$C^*$ -rigid within  $\mathcal{G}$**  if  $C^*(G_0) \simeq C^*(G) \Leftrightarrow G_0 \simeq G$  for all  $G$  in  $\mathcal{G}$

Which groups are  $C^*$ -rigid within the class of **1-connected** Lie groups?

(i.e., connected and simply connected)

**Example** (commutative groups):

Let  $G_0, G$  be 1-connected Lie groups with  $G_0$  commutative,  
hence  $G_0 \simeq (\mathbb{R}^n, +)$ .

# $C^*$ -rigidity of Lie groups

Given a class of Lie groups  $\mathcal{G}$  and  $G_0$  in  $\mathcal{G}$ .

$G_0$  is called  **$C^*$ -rigid within  $\mathcal{G}$**  if  $C^*(G_0) \simeq C^*(G) \Leftrightarrow G_0 \simeq G$  for all  $G$  in  $\mathcal{G}$

Which groups are  $C^*$ -rigid within the class of **1-connected** Lie groups?

(i.e., connected and simply connected)

**Example** (commutative groups):

Let  $G_0, G$  be 1-connected Lie groups with  $G_0$  commutative, hence  $G_0 \simeq (\mathbb{R}^n, +)$ .

Step 1:  $C^*(G_0) \simeq C^*(G) \implies C^*(G) \text{ commutative} \implies G \text{ commutative}$

# $C^*$ -rigidity of Lie groups

Given a class of Lie groups  $\mathcal{G}$  and  $G_0$  in  $\mathcal{G}$ .

$G_0$  is called  **$C^*$ -rigid within  $\mathcal{G}$**  if  $C^*(G_0) \simeq C^*(G) \Leftrightarrow G_0 \simeq G$  for all  $G$  in  $\mathcal{G}$

Which groups are  $C^*$ -rigid within the class of **1-connected** Lie groups?

(i.e., connected and simply connected)

**Example** (commutative groups):

Let  $G_0, G$  be 1-connected Lie groups with  $G_0$  commutative, hence  $G_0 \simeq (\mathbb{R}^n, +)$ .

Step 1:  $C^*(G_0) \simeq C^*(G) \implies C^*(G) \text{ commutative} \implies G \text{ commutative}$

Step 2:  $\mathcal{C}_0(\widehat{G_0}) \simeq \mathcal{C}_0(\widehat{G}) \implies G_0 \simeq G$  as above, by Brouwer's theorem

# Counterexamples

# Counterexamples

**Example 1** (compact groups). For the torus  $U(1)^n$  we have  $\mathbb{Z}^n \xrightarrow{\sim} \widehat{U(1)^n}$ , via  $\alpha \mapsto \chi_\alpha$ , where  $\chi_\alpha(z) = z^\alpha$ .

# Counterexamples

**Example 1** (compact groups). For the torus  $U(1)^n$  we have  $\mathbb{Z}^n \xrightarrow{\sim} \widehat{U(1)^n}$ , via  $\alpha \mapsto \chi_\alpha$ , where  $\chi_\alpha(z) = z^\alpha$ .

$\xrightarrow{\sim} U(1)^n \not\cong U(1)^m$  as Lie groups if  $n \neq m$

# Counterexamples

**Example 1** (compact groups). For the torus  $U(1)^n$  we have  $\mathbb{Z}^n \xrightarrow{\sim} \widehat{U(1)^n}$ , via  $\alpha \mapsto \chi_\alpha$ , where  $\chi_\alpha(z) = z^\alpha$ .

$\xrightarrow{\sim} U(1)^n \not\simeq U(1)^m$  as Lie groups if  $n \neq m$ , yet  $\widehat{U(1)^n} \simeq \widehat{U(1)^m}$  since  $\mathbb{Z}^n \simeq \mathbb{Z}^m$  as topological spaces, hence  $C^*(U(1)^n) \simeq C^*(U(1)^m)$ .



# Counterexamples

**Example 1** (compact groups). For the torus  $U(1)^n$  we have  $\mathbb{Z}^n \xrightarrow{\sim} \widehat{U(1)^n}$ , via  $\alpha \mapsto \chi_\alpha$ , where  $\chi_\alpha(z) = z^\alpha$ .

$\leadsto U(1)^n \not\simeq U(1)^m$  as Lie groups if  $n \neq m$ , yet  $\widehat{U(1)^n} \simeq \widehat{U(1)^m}$  since  $\mathbb{Z}^n \simeq \mathbb{Z}^m$  as topological spaces, hence  $C^*(U(1)^n) \simeq C^*(U(1)^m)$ .

$\leadsto$  We stick to *1-connected* Lie groups.

# Counterexamples

**Example 1** (compact groups). For the torus  $U(1)^n$  we have  $\mathbb{Z}^n \xrightarrow{\sim} \widehat{U(1)^n}$ , via  $\alpha \mapsto \chi_\alpha$ , where  $\chi_\alpha(z) = z^\alpha$ .

$\leadsto U(1)^n \not\simeq U(1)^m$  as Lie groups if  $n \neq m$ , yet  $\widehat{U(1)^n} \simeq \widehat{U(1)^m}$  since  $\mathbb{Z}^n \simeq \mathbb{Z}^m$  as topological spaces, hence  $C^*(U(1)^n) \simeq C^*(U(1)^m)$ .

$\leadsto$  We stick to *1-connected* Lie groups.

**Example 2** (1-connected solvable Lie groups). For  $D \in M_n(\mathbb{R})$  let

$G_D := \mathbb{R}^n \rtimes_D \mathbb{R}$  with  $(b_1, t_1) \cdot (b_2, t_2) = (b_1 + e^{t_1 D} b_2, t_1 + t_2)$ .

Also let  $n_\pm^D$  be the number of eigenvalues  $z \in \sigma(D)$  with  $\pm \operatorname{Re} z > 0$ .

# Counterexamples

**Example 1** (compact groups). For the torus  $U(1)^n$  we have  $\mathbb{Z}^n \xrightarrow{\sim} \widehat{U(1)^n}$ , via  $\alpha \mapsto \chi_\alpha$ , where  $\chi_\alpha(z) = z^\alpha$ .

$\leadsto U(1)^n \not\cong U(1)^m$  as Lie groups if  $n \neq m$ , yet  $\widehat{U(1)^n} \simeq \widehat{U(1)^m}$  since  $\mathbb{Z}^n \simeq \mathbb{Z}^m$  as topological spaces, hence  $C^*(U(1)^n) \simeq C^*(U(1)^m)$ .

$\leadsto$  We stick to *1-connected* Lie groups.

**Example 2** (1-connected solvable Lie groups). For  $D \in M_n(\mathbb{R})$  let

$$G_D := \mathbb{R}^n \rtimes_D \mathbb{R} \text{ with } (b_1, t_1) \cdot (b_2, t_2) = (b_1 + e^{t_1 D} b_2, t_1 + t_2).$$

Also let  $n_\pm^D$  be the number of eigenvalues  $z \in \sigma(D)$  with  $\pm \operatorname{Re} z > 0$ .

If  $D_j \in M_n(\mathbb{R})$  and  $n_+^{D_j} + n_-^{D_j} = n$  for  $j = 1, 2$ , then

- ▶  $G_{D_1} \simeq G_{D_2}$  as Lie groups iff  $\sigma(D_1) = r \cdot \sigma(D_2)$  for some  $r \in \mathbb{R} \setminus \{0\}$

# Counterexamples

**Example 1** (compact groups). For the torus  $U(1)^n$  we have  $\mathbb{Z}^n \xrightarrow{\sim} \widehat{U(1)^n}$ , via  $\alpha \mapsto \chi_\alpha$ , where  $\chi_\alpha(z) = z^\alpha$ .

$\leadsto U(1)^n \not\simeq U(1)^m$  as Lie groups if  $n \neq m$ , yet  $\widehat{U(1)^n} \simeq \widehat{U(1)^m}$  since  $\mathbb{Z}^n \simeq \mathbb{Z}^m$  as topological spaces, hence  $C^*(U(1)^n) \simeq C^*(U(1)^m)$ .

$\leadsto$  We stick to *1-connected* Lie groups.

**Example 2** (1-connected solvable Lie groups). For  $D \in M_n(\mathbb{R})$  let

$$G_D := \mathbb{R}^n \rtimes_D \mathbb{R} \text{ with } (b_1, t_1) \cdot (b_2, t_2) = (b_1 + e^{t_1 D} b_2, t_1 + t_2).$$

Also let  $n_\pm^D$  be the number of eigenvalues  $z \in \sigma(D)$  with  $\pm \operatorname{Re} z > 0$ .

If  $D_j \in M_n(\mathbb{R})$  and  $n_+^{D_j} + n_-^{D_j} = n$  for  $j = 1, 2$ , then

- ▶  $G_{D_1} \simeq G_{D_2}$  as Lie groups iff  $\sigma(D_1) = r \cdot \sigma(D_2)$  for some  $r \in \mathbb{R} \setminus \{0\}$
- ▶  $C^*(G_{D_1}) \simeq C^*(G_{D_2})$  iff  $\{n_+^{D_1}, n_-^{D_1}\} = \{n_+^{D_2}, n_-^{D_2}\}$ .

# The main result

# The main result

## Theorem

*Heisenberg groups are  $C^*$ -rigid within the class of 1-connected Lie groups.*

# The main result

## Theorem

*Heisenberg groups are  $C^*$ -rigid within the class of 1-connected Lie groups.*

That is, for every 1-connected Lie group,

$$H_n \simeq G \iff C^*(H_n) \simeq C^*(G)$$

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step 1}} G \text{ solvable}$$



# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step 1}} G \text{ solvable} \xRightarrow{\text{Step 2}} G \text{ nilpotent}$$

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step 1}} G \text{ solvable} \xRightarrow{\text{Step 2}} G \text{ nilpotent} \xRightarrow{\text{Step 3}} G \simeq H_n.$$

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step1}} G \text{ solvable} \xRightarrow{\text{Step2}} G \text{ nilpotent} \xRightarrow{\text{Step3}} G \simeq H_n.$$

## Step 1:

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  special topological features of  $\widehat{G}$ : finite subsets are closed; every limit set is either a singleton or  $\mathbb{R}^{2n}$

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step 1}} G \text{ solvable} \xRightarrow{\text{Step 2}} G \text{ nilpotent} \xRightarrow{\text{Step 3}} G \simeq H_n.$$

## Step 1:

- (a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  special topological features of  $\widehat{G}$ : finite subsets are closed; every limit set is either a singleton or  $\mathbb{R}^{2n}$
- (b)  $G = G_1 \times (S \rtimes K)$  by structure theory of Lie groups,

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step1}} G \text{ solvable} \xRightarrow{\text{Step2}} G \text{ nilpotent} \xRightarrow{\text{Step3}} G \simeq H_n.$$

## Step 1:

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  special topological features of  $\widehat{G}$ : finite subsets are closed; every limit set is either a singleton or  $\mathbb{R}^{2n}$

(b)  $G = G_1 \times (S \rtimes K)$  by structure theory of Lie groups,

where  $G_1$  is semisimple,  $K$  is compact semisimple, and  $S$  is solvable.

$\leadsto C^*(H_n) \simeq C^*(G) = C^*(G_1) \otimes C^*(S \rtimes K)$  and  $\widehat{H_n} \simeq \widehat{G_1} \times \widehat{S \rtimes K}$

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step1}} G \text{ solvable} \xRightarrow{\text{Step2}} G \text{ nilpotent} \xRightarrow{\text{Step3}} G \simeq H_n.$$

## Step 1:

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  special topological features of  $\widehat{G}$ : finite subsets are closed; every limit set is either a singleton or  $\mathbb{R}^{2n}$

(b)  $G = G_1 \times (S \rtimes K)$  by structure theory of Lie groups,

where  $G_1$  is semisimple,  $K$  is compact semisimple, and  $S$  is solvable.

$\leadsto C^*(H_n) \simeq C^*(G) = C^*(G_1) \otimes C^*(S \rtimes K)$  and  $\widehat{H_n} \simeq \widehat{G_1} \times \widehat{S \rtimes K}$

(c)  $H_n$  has no irred. representations of finite dim.  $> 1 \leadsto K = \{\mathbf{1}\}$

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step 1}} G \text{ solvable} \xRightarrow{\text{Step 2}} G \text{ nilpotent} \xRightarrow{\text{Step 3}} G \simeq H_n.$$

## Step 1:

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  special topological features of  $\widehat{G}$ : finite subsets are closed; every limit set is either a singleton or  $\mathbb{R}^{2n}$

(b)  $G = G_1 \times (S \rtimes K)$  by structure theory of Lie groups,

where  $G_1$  is semisimple,  $K$  is compact semisimple, and  $S$  is solvable.

$\leadsto C^*(H_n) \simeq C^*(G) = C^*(G_1) \otimes C^*(S \rtimes K)$  and  $\widehat{H_n} \simeq \widehat{G_1} \times \widehat{S \rtimes K}$

(c)  $H_n$  has no irred. representations of finite dim.  $> 1 \leadsto K = \{\mathbf{1}\}$

(d) Every limit set in  $\widehat{G_1}$  is at most countable  $\leadsto G_1 = \{\mathbf{1}\}$ .

# A breakdown of the proof

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then

$$G \text{ 1-connected} \xRightarrow{\text{Step 1}} G \text{ solvable} \xRightarrow{\text{Step 2}} G \text{ nilpotent} \xRightarrow{\text{Step 3}} G \simeq H_n.$$

## Step 1:

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  special topological features of  $\widehat{G}$ : finite subsets are closed; every limit set is either a singleton or  $\mathbb{R}^{2n}$

(b)  $G = G_1 \times (S \rtimes K)$  by structure theory of Lie groups,

where  $G_1$  is semisimple,  $K$  is compact semisimple, and  $S$  is solvable.

$\leadsto C^*(H_n) \simeq C^*(G) = C^*(G_1) \otimes C^*(S \rtimes K)$  and  $\widehat{H_n} \simeq \widehat{G_1} \times \widehat{S \rtimes K}$

(c)  $H_n$  has no irred. representations of finite dim.  $> 1 \leadsto K = \{\mathbf{1}\}$

(d) Every limit set in  $\widehat{G_1}$  is at most countable  $\leadsto G_1 = \{\mathbf{1}\}$ .

Thus  $G = S$  is a solvable Lie group.



# A breakdown of the proof (2)

## A breakdown of the proof (2)

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then  $G$  is solvable by Step 1.

Next:

$G$  solvable

## A breakdown of the proof (2)

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then  $G$  is solvable by Step 1.

Next:

$$G \text{ solvable} \xRightarrow{\text{Step 2}} G \text{ nilpotent}$$

## A breakdown of the proof (2)

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then  $G$  is solvable by Step 1.  
Next:

$$G \text{ solvable} \xrightarrow{\text{Step 2}} G \text{ nilpotent} \xrightarrow{\text{Step 3}} G \simeq H_n.$$

**Step 2:** The orbits of the coadjoint action of  $H_n$  on the dual of its Lie algebra are closed and 1-connected.

## A breakdown of the proof (2)

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then  $G$  is solvable by Step 1.

Next:

$$G \text{ solvable} \xrightarrow{\text{Step 2}} G \text{ nilpotent} \xrightarrow{\text{Step 3}} G \simeq H_n.$$

**Step 2:** The orbits of the coadjoint action of  $H_n$  on the dual of its Lie algebra are closed and 1-connected.

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  The coadjoint orbits of  $G$  are closed and 1-connected, too.

## A breakdown of the proof (2)

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then  $G$  is solvable by Step 1.

Next:

$$G \text{ solvable} \xrightarrow{\text{Step 2}} G \text{ nilpotent} \xrightarrow{\text{Step 3}} G \simeq H_n.$$

**Step 2:** The orbits of the coadjoint action of  $H_n$  on the dual of its Lie algebra are closed and 1-connected.

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  The coadjoint orbits of  $G$  are closed and 1-connected, too.

(b) The above properties implies that  $G$  is nilpotent.

## A breakdown of the proof (2)

If  $G$  is a Lie group and  $C^*(G) \simeq C^*(H_n)$ , then  $G$  is solvable by Step 1.

Next:

$$G \text{ solvable} \xrightarrow{\text{Step 2}} G \text{ nilpotent} \xrightarrow{\text{Step 3}} G \simeq H_n.$$

**Step 2:** The orbits of the coadjoint action of  $H_n$  on the dual of its Lie algebra are closed and 1-connected.

(a)  $\widehat{G} \simeq \widehat{H_n} \leadsto$  The coadjoint orbits of  $G$  are closed and 1-connected, too.

(b) The above properties implies that  $G$  is nilpotent.

**Step 3:**  $H_n$  is  $C^*$ -rigid within the class of nilpotent Lie groups (earlier result).

Q.E.D.

Refinement of  $C^*(G) \simeq C^*(H_n) \implies G \simeq H_n$



Refinement of  $C^*(G) \simeq C^*(H_n) \implies G \simeq H_n$

Recall: For any Lie groups  $G$  and  $H$ ,  $C^*(G) \simeq C^*(H) \implies \widehat{G} \simeq \widehat{H}$

Refinement of  $C^*(G) \simeq C^*(H_n) \implies G \simeq H_n$

Recall: For any Lie groups  $G$  and  $H$ ,  $C^*(G) \simeq C^*(H) \implies \widehat{G} \simeq \widehat{H}$

**Question:**  $\widehat{G} \simeq \widehat{H} \stackrel{?}{\implies} G \simeq H$

Refinement of  $C^*(G) \simeq C^*(H_n) \implies G \simeq H_n$

Recall: For any Lie groups  $G$  and  $H$ ,  $C^*(G) \simeq C^*(H) \implies \widehat{G} \simeq \widehat{H}$

**Question:**  $\widehat{G} \simeq \widehat{H} \stackrel{?}{\implies} G \simeq H$

**Answer:** YES if  $H = H_n$ .

Refinement of  $C^*(G) \simeq C^*(H_n) \implies G \simeq H_n$

Recall: For any Lie groups  $G$  and  $H$ ,  $C^*(G) \simeq C^*(H) \implies \widehat{G} \simeq \widehat{H}$

**Question:**  $\widehat{G} \simeq \widehat{H} \stackrel{?}{\implies} G \simeq H$

**Answer:** YES if  $H = H_n$ .

**Note:**  $\widehat{G}$  carries less information than  $C^*(G)$ .

Refinement of  $C^*(G) \simeq C^*(H_n) \implies G \simeq H_n$

Recall: For any Lie groups  $G$  and  $H$ ,  $C^*(G) \simeq C^*(H) \implies \widehat{G} \simeq \widehat{H}$

**Question:**  $\widehat{G} \simeq \widehat{H} \stackrel{?}{\implies} G \simeq H$

**Answer:** YES if  $H = H_n$ .

**Note:**  $\widehat{G}$  carries less information than  $C^*(G)$ .

E.g., the compact Lie groups  $SU(2)$  and  $SU(3)$  are 1-connected and  $\widehat{SU(2)} \simeq \widehat{SU(3)}$  but  $C^*(SU(2)) \not\simeq C^*(SU(3))$ .