

# Toward stationary solutions for cubic NLS with linear band crossing of Dirac type

---

Elena Danesi (Politecnico di Torino)

jw William Borrelli, Simone Dovetta and Lorenzo Tentarelli

Analytic and algebraic methods in physics XXII, Prague

27 August 2025

The nonlinear Dirac equation has been widely used to build *relativistic* models of extended particles

- Existence and properties of solutions: Cazenave-Vazquez '86, Balabane-Cazenave-Vazquez '90, Coti Zelati-Nolasco '25, Esteban-Séré '95,...
- Dirac-Maxwell, Dirac-Maxwell-Einstein eqs: Esteban-Georgiev-Séré '96, Nolasco '21, Paturel '00, Rota Nodari '10, Comech-Stuart '18,...
- Mean-field models for nucleons and nuclei: Esteban-Rota Nodari '12, Le Treust-Rota Nodari '13, Rota Nodari '12,...

# Introduction

The nonlinear Dirac equation has been widely used to build *relativistic* models of extended particles

- Existence and properties of solutions: Cazenave-Vazquez '86, Balabane-Cazenave-Vazquez '90, Coti Zelati-Nolasco '25, Esteban-Séré '95,...
- Dirac-Maxwell, Dirac-Maxwell-Einstein eqs: Esteban-Georgiev-Séré '96, Nolasco '21, Paturel '00, Rota Nodari '10, Comech-Stuart '18,...
- Mean-field models for nucleons and nuclei: Esteban-Rota Nodari '12, Le Treust-Rota Nodari '13, Rota Nodari '12,...

Recently, nonlinear Dirac equations appeared as effective models in solid state physics/optical systems, where particles have *linear* dispersion relation, in some regime.

# Introduction

We consider cubic focusing NLS

$$i\partial_t v = (-\Delta + V)v - |v|^2 v, \quad \text{on } \mathbb{R}_t \times \mathbb{R}_x^d, \quad d = 1, 2, \quad (1)$$

where  $V \in C^\infty(\mathbb{R}^d; \mathbb{R})$  is periodic w.r.t. some  $d$ -dimensional lattice  $\Lambda$ , describing an underlying periodic structure.

## The spectrum of $-\Delta + V$

In what follows,  $\mathcal{B}$  is the Brillouin zone of the dual lattice  $\Lambda^*$ .

## The spectrum of $-\Delta + V$

In what follows,  $\mathcal{B}$  is the Brillouin zone of the dual lattice  $\Lambda^*$ .

The spectrum  $\sigma(-\Delta + V) \subset \mathbb{R}$  is given by a union of **spectral bands**.

## The spectrum of $-\Delta + V$

In what follows,  $\mathcal{B}$  is the Brillouin zone of the dual lattice  $\Lambda^*$ . The spectrum  $\sigma(-\Delta + V) \subset \mathbb{R}$  is given by a union of **spectral bands**. They can be obtained by solving, for each  $k \in \mathcal{B}$  the **Floquet-Bloch eigenvalue problem**:

$$\begin{cases} (-\Delta + V)\Phi(x, k) = \mu(k)\Phi(x, k), & x \in \mathbb{R}^d \\ \Phi(x + v, k) = e^{ik \cdot v}\Phi(x, k), & v \in \Lambda \end{cases} \quad (2)$$

This yields a countable sequence of real-valued eigenvalues which are ordered, including multiplicity, such that

$$\mu_0(k) \leq \mu_1(k) \leq \mu_2(k) \leq \dots$$

# Floquet-Bloch decomposition

The corresponding eigenfunctions  $\Phi_j(\cdot, k)$  are called **Bloch waves**.

There holds

$$\sigma(-\Delta + V) = \bigcup_{j \in \mathbb{N}} \{\mu_j(k) : k \in \mathcal{B}\}$$

and the *Bloch functions*  $(\Phi_j(\cdot, k))_{j \in \mathbb{N}, k \in \mathcal{B}}$  form a complete set in  $L^2(\mathbb{R}^d)$  :

$$f(x) = \sum_{j \in \mathbb{N}} \int_{\mathcal{B}} \langle \Phi_j(\cdot, k), f(\cdot) \rangle_{L^2(\mathbb{R}^2)} \Phi_j(x, k) dk ,$$

$$e^{-it(-\Delta + V)} f(x) = \sum_{j \in \mathbb{N}} \int_{\mathcal{B}} e^{-i\mu_j(k)t} \langle \Phi_j(\cdot, k), f(\cdot) \rangle_{L^2(\mathbb{R}^2)} \Phi_j(x, k) dk .$$





# Dirac points

We say that  $-\Delta + V$  has a **linear/conical crossing** at a **Dirac point**  $k_* \in \mathcal{B}$ , if there are two bands  $\mu_n(k), \mu_{n+1}(k)$  such that

- $\mu(k_*) = \mu_n(k_*) = \mu_{n+1}(k_*)$  is a double eigenvalue of (2) ;
- there exist  $E_{\pm} : B_{\delta} \rightarrow \mathbb{R}$ , Lipschitz, s.t. if  $|k - k_*| < \delta$

$$\begin{aligned}\mu_{n+1}(k) - \mu(k_*) &= |\lambda| |k - k_*| (1 + E_+(k - k_*)) \\ \mu_n(k) - \mu(k_*) &= -|\lambda| |k - k_*| (1 + E_-(k - k_*))\end{aligned}\tag{3}$$

where  $E_{\pm}(\kappa) = O(\kappa)$ , as  $\kappa \rightarrow 0$ , and  $\lambda \in \mathbb{C}$  depends on  $V$ .



## Main goal

- We assume that the linear part has a *linear/conical band crossing*.
- This leads to some effective Dirac dynamics.
- The latter can be justified for special initial data, on suitable timescales.
- We are interested in the corresponding stationary NLD.
- Our aim is to construct *stationary solutions* to (1), using Dirac solitons.

## Linear Dirac dynamics in $2d$

Consider a wave packet spectrally concentrated around  $k_*$ :

$$u_0^\delta(x) = \delta(\psi_{1,0}(\delta x)\Phi_1(x) + \psi_{2,0}(\delta x)\Phi_2(x)), \quad \delta > 0, \quad (4)$$

the  $\Phi_{1,2}$  are the Bloch functions at  $k_*$ .

## Linear Dirac dynamics in $2d$

Consider a **wave packet spectrally concentrated around  $k_*$** :

$$u_0^\delta(x) = \delta(\psi_{1,0}(\delta x)\Phi_1(x) + \psi_{2,0}(\delta x)\Phi_2(x)), \quad \delta > 0, \quad (4)$$

the  $\Phi_{1,2}$  are the Bloch functions at  $k_*$ . The problem

$$\begin{cases} i\partial_t u = (-\Delta + V)u \\ u^\delta(0, x) = u_0^\delta(x) \end{cases}$$

has a unique solution of the form

$$u^\delta(t, x) = e^{-i\mu_* t} \left( \sum_{j=1}^2 \delta\psi_j(\delta t, \delta x)\Phi_j(x) + \eta^\delta(t, x) \right) \quad (5)$$

with  $\eta^\delta(0, x) = 0$  and  $\sup_{0 \leq t \leq \delta^{-2+}} \|\eta^\delta(t, \cdot)\|_{H_x^s} = O(\delta^\tau)$  for any  $s > 0$ , for some  $\tau > 0$  [Fefferman-Weinstein '13].

## Linear Dirac dynamics in $2d$

The coefficients  $\psi_j$  form a global-in-time solution to the following Dirac equation

$$\partial_t \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} 0 & -\bar{\lambda}(\partial_1 + i\partial_2) \\ -\lambda(\partial_1 - i\partial_2) & 0 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad 0 \neq \lambda \in \mathbb{C}$$

with initial data  $\begin{pmatrix} \psi_1(0, x) \\ \psi_2(0, x) \end{pmatrix} = \begin{pmatrix} \psi_{1,0}(x) \\ \psi_{2,0}(x) \end{pmatrix} \in [\mathcal{S}(\mathbb{R}^2)]^2$ .

The parameter  $\lambda \in \mathbb{C} \setminus \{0\}$  depends on the potential  $V$ .

# Nonlinear Dirac dynamics in 2d

Consider the following *focusing* NLS:

$$i\partial_t v = (-\Delta + V)v - |v|^2 v$$

with  $V$  as before.

The effective equation around a Dirac point is  
([Fefferman-Weinstein'12], formal derivation):

$$\begin{cases} \partial_t \psi_1 + \bar{\lambda}(\partial_1 + i\partial_2)\psi_2 = i(\beta_1|\psi_1|^2 + 2\beta_2|\psi_2|^2)\psi_1 \\ \partial_t \psi_2 + \lambda(\partial_1 - i\partial_2)\psi_1 = i(\beta_1|\psi_2|^2 + 2\beta_2|\psi_1|^2)\psi_2 \end{cases} \quad (6)$$

with  $0 \neq \lambda \in \mathbb{C}$ ,  $0 < \beta_2(V) \leq \beta_1(V)$  and  $\psi = (\psi_1, \psi_2)^T$  is a  $\mathbb{C}^2$ -spinor.

## Nonlinear Dirac dynamics in $2d$

Idea of the derivation: make a multiscale ansatz

$$v(t, x) \approx \sqrt{\delta} e^{-it\mu_*} \sum_{n=0}^N \delta^n v_n(\delta t, \delta x, x). \quad (7)$$

The system (6) describes the leading order term. In this case the approximation is valid in  $H^s$ -sense,  $s > 1$ , on a time-scale  $T_\delta \sim \delta^{-1}$  [Arbunich-Sparber '18].

## Towards Dirac solitons for honeycomb NLS

We are interested in stationary solutions of the focusing NLS

$$v(t, x) = e^{-i\mu_* t} u(x)$$

where  $\mu_* \in \sigma(-\Delta + V)$  is the energy of a Dirac point.



# Towards Dirac solitons for honeycomb NLS

We are interested in stationary solutions of the focusing NLS

$$v(t, x) = e^{-i\mu_* t} u(x)$$

where  $\mu_* \in \sigma(-\Delta + V)$  is the energy of a Dirac point. Then  $u$  solves

$$(-\Delta + V - \mu_*)u = |u|^2 u.$$

# Towards Dirac solitons for honeycomb NLS

We are interested in stationary solutions of the focusing NLS

$$v(t, x) = e^{-i\mu_* t} u(x)$$

where  $\mu_* \in \sigma(-\Delta + V)$  is the energy of a Dirac point. Then  $u$  solves

$$(-\Delta + V - \mu_*)u = |u|^2 u.$$

In particular, we may look for solutions of the form

$$u^\delta(x) = \sqrt{\delta} \sum_{j=1}^2 \psi_j(\delta x) \Phi_j(x) + \sqrt{\delta} \eta^\delta(x), \quad \delta > 0.$$

## The 1d case

- $\mu_* \in \sigma(-\Delta + V)$  lies in the spectrum.
- One cannot expect to build stationary solutions for frequencies in the linear spectrum.

## The 1d case

- $\mu_* \in \sigma(-\Delta + V)$  lies in the spectrum.
- One cannot expect to build stationary solutions for frequencies in the linear spectrum.
- A spectral gap around  $\mu_*$  is needed. (Similar idea in [Fefferman-Lee-Weinstein '12] to build bound states for 1d Schrödinger operators with a domain wall (slowly) interpolating between two periodic structures).

## The 1d case

- $\mu_* \in \sigma(-\Delta + V)$  lies in the spectrum.
- One cannot expect to build stationary solutions for frequencies in the linear spectrum.
- A spectral gap around  $\mu_*$  is needed. (Similar idea in [Fefferman-Lee-Weinstein '12] to build bound states for 1d Schrödinger operators with a domain wall (slowly) interpolating between two periodic structures).

Consider the lattice  $\Lambda = \mathbb{Z}$ , with Brillouin zone  $\mathcal{B} = [0, 2\pi]$ , so  $V \in C^\infty(\mathbb{R})$  is 1-periodic.

- If  $V$  is even (plus an additional translational symmetry) it has a Dirac point at  $k_* = \pi$  [FLW '12].
- Breaking parity with a perturbation  $\delta W$  one can open a gap of size  $O(\delta)$ : if  $0 < \delta \ll 1$  one may look for stationary solutions with frequency in the gap.

## Dirac solitons in 1d NLS

We consider the following 1d NLS

$$\left( -\frac{d^2}{dx^2} + V(x) + \delta W(x) - \mu_\delta \right) u = |u|^2 u, \quad (8)$$

where  $V$  and  $W$  are smooth 1-periodic potentials as described before, and  $\mu_\delta \in \mathbb{R}$ .

## Dirac solitons in 1d NLS

We consider the following 1d NLS

$$\left( -\frac{d^2}{dx^2} + V(x) + \delta W(x) - \mu_\delta \right) u = |u|^2 u, \quad (8)$$

where  $V$  and  $W$  are smooth 1-periodic potentials as described before, and  $\mu_\delta \in \mathbb{R}$ .

- Assume there exists a spectral gap  $I_\delta := (\mu_* - c\delta, \mu_* + c\delta)$  around the Dirac energy  $\mu_*$ .
- Taking  $\mu_\delta \in I_\delta$  it should be possible to prove the existence of stationary (localized) solutions (of arbitrary form), for instance by variational methods.
- So it makes sense to also look for them with the “Dirac ansatz”.

## The effective Dirac equation in 1d

As in the 2d case, the effective equation can be found *via* a multiscale expansion.

$$u(x) = \sqrt{\delta} \mathcal{U}_\delta(x, y) = \sqrt{\delta} (U_0(x, y) + \delta U_1(x, y) + \dots),$$

with  $y = \delta x$ ,  $\mathcal{U}_\delta(x + 1, y) = e^{ik_*} \mathcal{U}_\delta(x, y)$  and

$$\mu_\delta = \mu_* + \delta \mu_1 + \delta^2 \mu_2 + \dots$$



Treating  $x$  and  $y$  as independent variables and substituting into the NLS:

$$\left[ -(\partial_x + \delta \partial_y)^2 + V(x) + \delta W(x) - \lambda_* - \delta \mu_1 - \dots \right] \mathcal{U}_\delta = \delta |\mathcal{U}_\delta|^2 \mathcal{U}_\delta.$$

We now solve the above equation order by order in powers of  $\delta$ .

One finds

$$U_0(x, y) = \psi(y) \Phi_1(x) + \psi_2(y) \Phi_2(x),$$

where  $\Phi_j$  are the Bloch waves at  $(k_*, \mu_*)$  and  $\psi = (\psi_1, \psi_-)^T$  solves a NLD.

# The effective Dirac equation in 1d

The effective Dirac equation is given by

$$(i\sigma_3\partial_y + \sigma_1 - \mu_1)\psi = \mathcal{G}_{\beta_1,\beta_2,\beta_3}(\psi) \quad (9)$$

where  $\psi = (\psi_1, \psi_2)^T$ , and  $\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ,  $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  are Pauli matrices and the nonlinearity  $\mathcal{G}_{\beta_1,\beta_2,\beta_3}$  is given by

$$\mathcal{G}_{\beta_1,\beta_2,\beta_3}(\psi) = \begin{pmatrix} \beta_1|\psi_1|^2 + 2\beta_2|\psi_2|^2 & \beta_3 \bar{\psi}_1\psi_2 \\ \beta_3 \psi_1\bar{\psi}_2 & \beta_1|\psi_2|^2 + 2\beta_2|\psi_1|^2 \end{pmatrix} \psi$$

Here  $\beta_1 = \int_0^1 |\Phi_1|^4 dx = \int_0^1 |\Phi_2|^4 dx$ ,  $\beta_2 = \int_0^1 |\Phi_1|^2 |\Phi_2|^2 dx$ ,  $\beta_3 := \int_0^1 \bar{\Phi}_1 \bar{\Phi}_1 \Phi_2 \Phi_2 dx$ .

The 1d Dirac operator  $\mathcal{D} = i\sigma_3\partial_y + \sigma_1$  has spectrum  $(-\infty, -1] \cup [1, +\infty)$ , so that we take  $|\mu_1| < 1$ .

# The main result in 1d

## Theorem [Borrelli-D.-Dovetta-Tentarelli]

Taking  $|\mu_1| < 1$ , the NLD admits a smooth solution  $\psi = (\psi_1, \psi_2)^T$ , such that  $|\psi(x)| \lesssim e^{-\sqrt{1-\mu^2}|x|}$  for  $|x| \gg 1$ .

Given  $0 < \delta \ll 1$ , for any  $\mu_\delta \in I_\delta$  in the linear spectral gap, the NLS

$$\left( -\frac{d^2}{dx^2} + V(x) + \delta W(x) - \mu_\delta \right) u = |u|^2 u$$

admits a smooth solution in  $H^1(\mathbb{R})$ , of the form

$$u^\delta(x) = \sqrt{\delta} \sum_{j=1}^2 \psi_j(\delta x) \Phi_j(x) + \eta^\delta(x),$$

with  $\|\eta^\delta\|_{H^1} = O(\delta)$ .

## Some ideas on the proof

First of all, one first proves the existence of a solution to the NLD and study its properties.

## Some ideas on the proof

First of all, one first proves the existence of a solution to the NLD and study its properties.

Then, one uses the solution of the effective NLD to construct a solution of the NLS:

- Plugging the ansatz

$$u^\delta(x) = \sqrt{\delta} \sum_{j=1}^2 \psi_j(\delta x) \Phi_j(x) + \sqrt{\delta} \eta^\delta(x)$$

into the NLS one gets an equation for  $\eta^\delta$ .

- One gets a nonlinear equation

$$\left(-\frac{d^2}{dx^2} + V(x) + \delta W(x) - \mu_*\right) \eta^\delta = \mathcal{F}(\psi, \eta^\delta, \delta). \quad (10)$$

## Some ideas on the proof

Idea: the dominant contribution should come from  $(k, \mu) \approx (k_*, \mu_*)$ .

- One decomposes  $\eta^\delta = \eta_{\text{near}}^\delta + \eta_{\text{far}}^\delta$  according to momentum/energy near and far from  $(k_*, \mu_*)$ .
- Projecting onto Bloch waves one gets an equation for  $\eta_{\text{far}}^\delta = \eta_{\text{far}}^\delta[\eta_{\text{near}}^\delta, \delta]$  and then a closed equation for  $\eta_{\text{near}}^\delta$ .
- The nonlinear term mixes up the near/far part of the splitting.

## The 2d case

The 2d case is conceptually analogous, but technically more involved.

- As in the 1d case, one can open a gap in the linear spectrum adding a perturbation  $\delta W$  *breaking parity*.
- The effective equation is now a *massive* Dirac equation, for which smooth and exponentially localized solutions are known to exist [Borrelli '17,'22].
- Then we may consider an analogous “Dirac ansatz” for the NLS.

## The $2d$ case

However

- when projecting onto Bloch waves we use the fact that in  $1d$  we can parametrize crossing bands *in a smooth way*. Not possible in  $2d$ .
- In  $2d$  Bloch bands are only Lipschitz continuous and Bloch waves are discontinuous at the conical point.
- At some point in  $1d$  we use ODE arguments.



Thank you for your attention!