

# Spectral properties of electromagnetic waveguides

Thomas OURMIÈRES-BONAFOS

Joint work with

- Philippe BRIET, Université de Toulon.
- Maxence CASSIER, Aix-Marseille Université.
- Michele ZACCARRON, Aix-Marseille Université.

AAMP

Prague

27, August 2025

Institut de Mathématiques de Marseille



## 1 Introduction

- What is a waveguide ?
- The Maxwell operator

## 2 Main results

- Essential spectrum
- Discrete spectrum

## 3 Geometric transforms and Birman-Schwinger principle

- The Piola transform
- Essential spectrum of constantly twisted waveguide
- A Resolvent identity

## 4 On the discrete spectrum

- A Poincaré inequality
- On the existence of discrete spectrum

## 5 Geometric condition $X \neq 0$

## 1 Introduction

- What is a waveguide ?
- The Maxwell operator

## 2 Main results

- Essential spectrum
- Discrete spectrum

## 3 Geometric transforms and Birman-Schwinger principle

- The Piola transform
- Essential spectrum of constantly twisted waveguide
- A Resolvent identity

## 4 On the discrete spectrum

- A Poincaré inequality
- On the existence of discrete spectrum

## 5 Geometric condition $X \neq 0$

# A curve in $\mathbb{R}^3$

Let us consider a curve  $\Gamma \subset \mathbb{R}^3$  parametrized by  $\gamma : \mathbb{R} \rightarrow \mathbb{R}^3$  such that:

- $\gamma : \mathbb{R} \rightarrow \mathbb{R}^3$  is of class  $\mathcal{C}^2(\mathbb{R})$ ,
- $\gamma$  is an arc-length parametrization :  $\forall s \in \mathbb{R}, \quad |\gamma'(s)| = 1$ ,

**Assumption 1:** The curvature of  $\Gamma$  at a point  $\gamma(s)$  is defined as  $\kappa(s) = |\gamma''(s)|$  and verifies

$$\lim_{|s| \rightarrow +\infty} \kappa(s) = 0.$$

To construct the waveguide we need:

- a frame adapted to the curve  $\Gamma$ ,
- a bounded, simply connected, Lipschitz domain  $\omega \subset \mathbb{R}^2$ .

# There is more than one way to frame a curve ! (1/2)

A standard choice is the Frenet frame defined at a point  $\gamma(s) \in \Gamma$  as

$$\begin{cases} e_1(s) &= \gamma'(s), \\ e_2(s) &= \frac{1}{\kappa(s)} \gamma''(s), \\ e_3(s) &= e_1(s) \times e_2(s). \end{cases}$$

The frame is adapted because each derivative  $\frac{d}{ds} e_j \in \text{span}(e_k, e_p)$  with  $p, k \neq j$ .

$$\begin{cases} \frac{d}{ds} e_1(s) &= \kappa(s) e_2(s), \\ \frac{d}{ds} e_2(s) &= -\kappa(s) e_1(s) + \tau(s) e_3(s), \\ \frac{d}{ds} e_3(s) &= -\tau(s) e_2(s). \end{cases}$$

Here  $\tau(s)$  is the torsion of the curve  $\Gamma$  at the point  $\gamma(s)$ .

# There is more than one way to frame a curve ! (1/2)

A standard choice is the Frenet frame defined at a point  $\gamma(s) \in \Gamma$  as

$$\begin{cases} e_1(s) &= \gamma'(s), \\ e_2(s) &= \frac{1}{\kappa(s)} \gamma''(s), \\ e_3(s) &= e_1(s) \times e_2(s). \end{cases}$$

The frame is adapted because each derivative  $\frac{d}{ds} e_j \in \text{span}(e_k, e_p)$  with  $p, k \neq j$ .

$$\begin{cases} \frac{d}{ds} e_1(s) &= \kappa(s) e_2(s), \\ \frac{d}{ds} e_2(s) &= -\kappa(s) e_1(s) + \tau(s) e_3(s), \\ \frac{d}{ds} e_3(s) &= -\tau(s) e_2(s). \end{cases}$$

Here  $\tau(s)$  is the torsion of the curve  $\Gamma$  at the point  $\gamma(s)$ .

**Problem** : when  $\kappa(s) = 0$  this is ill-defined.

In particular if the curve contains part of a straight line one cannot define this frame.

## There is more than one way to frame a curve ! (2/2)

Instead, we fix  $e_1(s) = \gamma'(s)$  and construct a **relatively adapted parallel frame**  $(e_1, e_2, e_3)$  which always exists and verify

$$\begin{cases} \frac{d}{ds} e_1(s) &= k_1(s) e_2(s) + k_2(s) e_3(s), \\ \frac{d}{ds} e_2(s) &= -k_1(s) e_1(s), \\ \frac{d}{ds} e_3(s) &= -k_2(s) e_1(s). \end{cases}$$

with  $k_1, k_2 \in \mathcal{C}^0(\mathbb{R})$  such that  $k_1(s)^2 + k_2(s)^2 = \kappa(s)^2$ .

## There is more than one way to frame a curve ! (2/2)

Instead, we fix  $e_1(s) = \gamma'(s)$  and construct a **relatively adapted parallel frame**  $(e_1, e_2, e_3)$  which always exists and verify

$$\begin{cases} \frac{d}{ds} e_1(s) &= k_1(s)e_2(s) + k_2(s)e_3(s), \\ \frac{d}{ds} e_2(s) &= -k_1(s)e_1(s), \\ \frac{d}{ds} e_3(s) &= -k_2(s)e_1(s). \end{cases}$$

with  $k_1, k_2 \in \mathcal{C}^0(\mathbb{R})$  such that  $k_1(s)^2 + k_2(s)^2 = \kappa(s)^2$ .

**Relatively parallel** :

the derivatives of  $e_2$  and  $e_3$  have a component only on  $e_1$ .



# There is more than one way to frame a curve ! (2/2)

Instead, we fix  $e_1(s) = \gamma'(s)$  and construct a **relatively adapted parallel frame**  $(e_1, e_2, e_3)$  which always exists and verify

$$\begin{cases} \frac{d}{ds} e_1(s) &= k_1(s) e_2(s) + k_2(s) e_3(s), \\ \frac{d}{ds} e_2(s) &= -k_1(s) e_1(s), \\ \frac{d}{ds} e_3(s) &= -k_2(s) e_1(s). \end{cases}$$

with  $k_1, k_2 \in \mathcal{C}^0(\mathbb{R})$  such that  $k_1(s)^2 + k_2(s)^2 = \kappa(s)^2$ .

**Relatively parallel** :

the derivatives of  $e_2$  and  $e_3$  have a component only on  $e_1$ .

The existence of such a frame can be found in



R. L. BISHOP

AM. MATH. MON. (1975)

Such a frame is unique up to fixed rotations in the plane  $\text{span}(e_2, e_3)$ .

# Construction of the waveguide (1/2)

Take  $\omega \subset \mathbb{R}^2$  a Lipschitz and simply connected domain and consider the map

$$\Phi: \begin{cases} \Omega_0 := \mathbb{R} \times \omega & \rightarrow \mathbb{R}^3 \\ (s, y_2, y_3) & \mapsto \gamma(s) + y_2 e_2(s) + y_3 e_3(s). \end{cases}$$

# Construction of the waveguide (1/2)

Take  $\omega \subset \mathbb{R}^2$  a Lipschitz and simply connected domain and consider the map

$$\Phi: \begin{cases} \Omega_0 := \mathbb{R} \times \omega & \rightarrow \mathbb{R}^3 \\ (s, y_2, y_3) & \mapsto \gamma(s) + y_2 e_2(s) + y_3 e_3(s). \end{cases}$$

Assumption 2:

$$\left( \sup_{y \in \omega} |y| \right) \|\kappa\|_{L^\infty(\mathbb{R})} < 1.$$

Under this condition  $\Phi$  is a  $\mathcal{C}^1$ -diffeomorphism from  $\Omega_0$  to its image.

# Construction of the waveguide (1/2)

Take  $\omega \subset \mathbb{R}^2$  a Lipschitz and simply connected domain and consider the map

$$\Phi : \begin{cases} \Omega_0 := \mathbb{R} \times \omega & \rightarrow \mathbb{R}^3 \\ (s, y_2, y_3) & \mapsto \gamma(s) + y_2 e_2(s) + y_3 e_3(s). \end{cases}$$

Assumption 2:

$$\left( \sup_{y \in \omega} |y| \right) \|\kappa\|_{L^\infty(\mathbb{R})} < 1.$$

Under this condition  $\Phi$  is a  $\mathcal{C}^1$ -diffeomorphism from  $\Omega_0$  to its image.

**Problem** : as constructed the cross-section  $\omega$  is not allowed to turn around the curve.

## Construction of the waveguide (2/2)

To remedy to this problem, let  $\theta : \mathbb{R} \rightarrow \mathbb{R}$  be of class  $\mathcal{C}^1(\mathbb{R})$  and define for  $p \in \{2, 3\}$ :

$$e_p^\theta(s) := \begin{pmatrix} \cos(\theta(s)) & -\sin(\theta(s)) \\ \sin(\theta(s)) & \cos(\theta(s)) \end{pmatrix} e_p(s).$$

## Construction of the waveguide (2/2)

To remedy to this problem, let  $\theta : \mathbb{R} \rightarrow \mathbb{R}$  be of class  $\mathcal{C}^1(\mathbb{R})$  and define for  $p \in \{2, 3\}$ :

$$e_p^\theta(s) := \begin{pmatrix} \cos(\theta(s)) & -\sin(\theta(s)) \\ \sin(\theta(s)) & \cos(\theta(s)) \end{pmatrix} e_p(s).$$

- $\theta'$  is called the **twist**: it is the rotation speed of  $(e_2^\theta, e_3^\theta)$  in the plane  $\text{span}(e_1)^\perp$ .

## Construction of the waveguide (2/2)

To remedy to this problem, let  $\theta : \mathbb{R} \rightarrow \mathbb{R}$  be of class  $\mathcal{C}^1(\mathbb{R})$  and define for  $p \in \{2, 3\}$ :

$$e_p^\theta(s) := \begin{pmatrix} \cos(\theta(s)) & -\sin(\theta(s)) \\ \sin(\theta(s)) & \cos(\theta(s)) \end{pmatrix} e_p(s).$$

Consider

$$\Phi : \begin{cases} \Omega_0 := \mathbb{R} \times \omega & \rightarrow \mathbb{R}^3 \\ (s, y_2, y_3) & \mapsto \gamma(s) + y_2 e_2^\theta(s) + y_3 e_3^\theta(s). \end{cases}$$

$\Phi$  is a  $\mathcal{C}^1(\mathbb{R})$  diffeomorphism and we set

$$\Omega := \Phi(\mathbb{R} \times \omega).$$

## Construction of the waveguide (2/2)

To remedy to this problem, let  $\theta : \mathbb{R} \rightarrow \mathbb{R}$  be of class  $\mathcal{C}^1(\mathbb{R})$  and define for  $p \in \{2, 3\}$ :

$$e_p^\theta(s) := \begin{pmatrix} \cos(\theta(s)) & -\sin(\theta(s)) \\ \sin(\theta(s)) & \cos(\theta(s)) \end{pmatrix} e_p(s).$$

Consider

$$\Phi : \begin{cases} \Omega_0 := \mathbb{R} \times \omega & \rightarrow \mathbb{R}^3 \\ (s, y_2, y_3) & \mapsto \gamma(s) + y_2 e_2^\theta(s) + y_3 e_3^\theta(s). \end{cases}$$

$\Phi$  is a  $\mathcal{C}^1(\mathbb{R})$  diffeomorphism and we set

$$\Omega := \Phi(\mathbb{R} \times \omega).$$

Assumption 3:

$$\lim_{|s| \rightarrow +\infty} \theta'(s) = \beta, \quad \beta \in \mathbb{R} \quad (\text{constant twist at infinity}).$$



# Some pictures !




Figure: A straight waveguide  $\Omega_0 := \mathbb{R} \times \omega$ .

# Some pictures !




Figure: A constantly twisted waveguide.

# Some pictures !




Figure: A curved and twisted waveguide.

# Physical context and Mathematical challenges

**Goal:** Define the Maxwell operator in a waveguide  $\Omega$  built with a homogeneous and isotropic material and embedded in a perfect conductor.

- $\varepsilon_0 > 0$ : permittivity of the material,
- $\mu_0 > 0$ : permeability of the material,
- $E$  : electric field,
- $H$  : magnetic field
- $E \times n = 0$  and  $H \cdot n = 0$  on  $\partial\Omega$  (perfectly conducting boundary condition).

# Physical context and Mathematical challenges

**Goal:** Define the Maxwell operator in a waveguide  $\Omega$  built with a homogeneous and isotropic material and embedded in a perfect conductor.

- $\varepsilon_0 > 0$ : permittivity of the material,
- $\mu_0 > 0$ : permeability of the material,
- $E$  : electric field,
- $H$  : magnetic field
- $E \times n = 0$  and  $H \cdot n = 0$  on  $\partial\Omega$  (perfectly conducting boundary condition).

The waveguide  $\Omega \subset \mathbb{R}^3$  is an unbounded Lipschitz domain:

**Problem**: how to define (and in which sense) the boundary condition to obtain a self-adjoint Maxwell operator.

# Minimal and maximal curl operators

- 1 Consider the curl operator acting in  $L^2(\Omega, \mathbb{C}^3)$  defined as

$$S_0 := \text{curl}, \quad \text{Dom}(S_0) = \mathcal{C}_0^\infty(\Omega, \mathbb{C}^3).$$

- 2 As  $S_0$  is symmetric (thus closable), one can define the minimal curl operator  $S = (S_0^*)^* = \overline{S_0}$  acting as

$$S = \text{curl}, \quad \text{Dom}(S) := H_0(\text{curl}, \Omega).$$

By definition  $H_0(\text{curl}, \Omega)$  is the closure of  $\mathcal{C}_0^\infty(\Omega, \mathbb{C}^3)$  with respect to the graph norm for the curl operator.

- 3 Remark that the maximal curl operator is  $S^*$  and acts as

$$S^* = \text{curl}, \quad \text{Dom}(S^*) = H(\text{curl}, \Omega) := \left\{ F \in L^2(\Omega, \mathbb{C}^3) : \text{curl} F \in L^2(\Omega, \mathbb{C}^3) \right\}.$$

- 4 The operator with domain  $H_0(\text{curl}, \Omega) \times H(\text{curl}, \Omega) \subset L^2(\Omega, \mathbb{C}^3 \times \mathbb{C}^3)$  acting as

$$\begin{pmatrix} 0 & \text{curl} \\ \text{curl} & 0 \end{pmatrix} = \begin{pmatrix} 0 & S^* \\ S & 0 \end{pmatrix},$$

is self-adjoint by construction.

# The boundary condition (electric field)

- If  $F_1, F_2 \in \mathcal{C}^\infty(\overline{\Omega}, \mathbb{R}^3)$ :

$$\int_{\Omega} ((\operatorname{curl} F_1) \cdot F_2) dx = \int_{\Omega} (F_1 \cdot (\operatorname{curl} F_2)) dx + \int_{\partial\Omega} (F_1 \times n) \cdot F_2 d\sigma$$

- $F_1 \in H_0(\operatorname{curl}, \Omega)$  if and only if for all  $F_2 \in H(\operatorname{curl}, \Omega)$  there holds

$$\int_{\Omega} ((\operatorname{curl} F_1) \cdot F_2) dx = \int_{\Omega} (F_1 \cdot (\operatorname{curl} F_2)) dx.$$

Hence, by  $F_1 \times n = 0$  on  $\partial\Omega$ , we mean that  $F_1 \in H_0(\operatorname{curl}, \Omega)$ .



R. DAUTRAY, J. -L. LIONS

MATH. ANALYSIS AND NUMERICAL METHODS FOR SCIENCE AND  
TECHNOLOGY: VOL. 3 (1990)



V. GIRAULT, P. -A. RAVIART

FINITE ELEMENT APPROXIMATION OF THE NAVIER-STOKES EQUATIONS  
(1981)

# Physical motivations

When there is an (exterior) current density  $\mathcal{J}$  the time-dependent Maxwell's equations in the waveguide  $\Omega$  read

$$\left\{ \begin{array}{ll} \varepsilon_0 \partial_t E(x, t) - \operatorname{curl} H(x, t) = -\mathcal{J}(x, t) & \text{for } (x, t) \in \Omega \times \mathbb{R}_+^*, \\ \mu_0 \partial_t H(x, t) + \operatorname{curl} E(x, t) = 0 & \text{for } (x, t) \in \Omega \times \mathbb{R}_+^*, \\ \operatorname{div}(\varepsilon_0 E)(x, t) = \operatorname{div}(\mu_0 H)(x, t) = 0 & \text{for } (x, t) \in \Omega \times \mathbb{R}_+^*, \\ E(x, t) \times n(x) = 0 & \text{for } (x, t) \in \partial\Omega \times \mathbb{R}_+^*, \\ H(x, t) \cdot n(x) = 0 & \text{for } (x, t) \in \partial\Omega \times \mathbb{R}_+^*, \\ (E(\cdot, 0), H(\cdot, 0)) = (E_0, H_0) & \text{with } (E_0, H_0) \in L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3). \end{array} \right.$$



# Physical motivations

When there is an (exterior) current density  $\mathcal{J}$  the time-dependent Maxwell's equations in the waveguide  $\Omega$  read

$$\left\{ \begin{array}{ll} \varepsilon_0 \partial_t E(x, t) - \operatorname{curl} H(x, t) = -\mathcal{J}(x, t) & \text{for } (x, t) \in \Omega \times \mathbb{R}_+^*, \\ \mu_0 \partial_t H(x, t) + \operatorname{curl} E(x, t) = 0 & \text{for } (x, t) \in \Omega \times \mathbb{R}_+^*, \\ \operatorname{div}(\varepsilon_0 E)(x, t) = \rho(x, t) \quad \operatorname{div}(\mu_0 H)(x, t) = 0 & \text{for } (x, t) \in \Omega \times \mathbb{R}_+^*, \\ E(x, t) \times n(x) = 0 & \text{for } (x, t) \in \partial\Omega \times \mathbb{R}_+^*, \\ H(x, t) \cdot n(x) = 0 & \text{for } (x, t) \in \partial\Omega \times \mathbb{R}_+^*, \\ (E(\cdot, 0), H(\cdot, 0)) = (E_0, H_0) & \text{with } (E_0, H_0) \in L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3). \end{array} \right.$$

# Physical motivations

It can be rewritten as a Schrödinger type equation

$$\begin{cases} \partial_t \psi + i\mathcal{A}\psi = f & \text{for all } t > 0, \\ \psi(0) = \psi_0 & \text{initial data in } L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3). \end{cases}$$

Here  $\psi = (E, H)^\top \in H_0(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ ,

- $\mathcal{A}$  is the self-adjoint Maxwell operator

$$\mathcal{A} := i \begin{pmatrix} 0 & \varepsilon_0^{-1} \text{curl} \\ -\mu_0^{-1} \text{curl} & 0 \end{pmatrix}, \quad \text{Dom}(\mathcal{A}) = H_0(\text{curl}, \Omega) \times H(\text{curl}, \Omega).$$

# Physical motivations

It can be rewritten as a Schrödinger type equation

$$\begin{cases} \partial_t \psi + i\mathcal{A}\psi = f & \text{for all } t > 0, \\ \psi(0) = \psi_0 & \text{initial data in } L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3). \end{cases}$$

Here  $\psi = (E, H)^\top \in H_0(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ ,

- $\mathcal{A}$  is the self-adjoint Maxwell operator

$$\mathcal{A} := i \begin{pmatrix} 0 & \varepsilon_0^{-1} \text{curl} \\ -\mu_0^{-1} \text{curl} & 0 \end{pmatrix}, \quad \text{Dom}(\mathcal{A}) = H_0(\text{curl}, \Omega) \times H(\text{curl}, \Omega).$$

- It acts in  $L^2_{\varepsilon_0, \mu_0}(\Omega, \mathbb{R}^3 \times \mathbb{R}^3)$  the space of square integrable vector fields. The scalar product is defined as

$$\left\langle \begin{pmatrix} E_1 \\ H_1 \end{pmatrix}, \begin{pmatrix} E_2 \\ H_2 \end{pmatrix} \right\rangle_{L^2_{\varepsilon_0, \mu_0}(\Omega, \mathbb{R}^3 \times \mathbb{R}^3)} = \int_{\Omega} (\varepsilon_0 E_1 \cdot E_2 + \mu_0 H_1 \cdot H_2) dx.$$

# Physical motivations

It can be rewritten as a Schrödinger type equation

$$\begin{cases} \partial_t \psi + i\mathcal{A}\psi = f & \text{for all } t > 0, \\ \psi(0) = \psi_0 & \text{initial data in } L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3). \end{cases}$$

Here  $\psi = (E, H)^\top \in H_0(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ ,

- $\mathcal{A}$  is the self-adjoint Maxwell operator

$$\mathcal{A} := i \begin{pmatrix} 0 & \varepsilon_0^{-1} \text{curl} \\ -\mu_0^{-1} \text{curl} & 0 \end{pmatrix}, \quad \text{Dom}(\mathcal{A}) = H_0(\text{curl}, \Omega) \times H(\text{curl}, \Omega).$$

**Motivation:**

The spectrum of  $\mathcal{A}$  gives the dynamics of the Maxwell system.

# Divergence free fields and the magnetic boundary condition

To look for divergence-free fields we introduce the solenoidal subspace

$$\mathcal{J}(\Omega) := \left\{ (E, H) \in L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3) : \operatorname{div}(\varepsilon_0 E) = \operatorname{div}(\mu_0 H) = 0, (H \cdot n)|_{\partial\Omega} = 0 \right\}.$$

# Divergence free fields and the magnetic boundary condition

To look for divergence-free fields we introduce the solenoidal subspace

$$\mathcal{J}(\Omega) := \left\{ (E, H) \in L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3) : \operatorname{div}(\varepsilon_0 E) = \operatorname{div}(\mu_0 H) = 0, (H \cdot n)|_{\partial\Omega} = 0 \right\}.$$

Here  $(H \cdot n)|_{\partial\Omega} = 0$  means that for all  $\varphi \in H^1(\Omega)$  there holds

$$0 = \int_{\Omega} \operatorname{div}(H) \varphi \, dx = - \int_{\Omega} H \cdot (\nabla \varphi) \, dx.$$

# Divergence free fields and the magnetic boundary condition

To look for divergence-free fields we introduce the solenoidal subspace

$$\mathcal{J}(\Omega) := \left\{ (E, H) \in L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3) : \operatorname{div}(\varepsilon_0 E) = \operatorname{div}(\mu_0 H) = 0, (H \cdot n)|_{\partial\Omega} = 0 \right\}.$$

## Lemma

There holds  $\ker(\mathcal{A})^\perp = \mathcal{J}(\Omega)$ . In particular,  $\mathcal{J}(\Omega)$  is a closed subspace left stable by  $\mathcal{A}$ .

# Divergence free fields and the magnetic boundary condition

To look for divergence-free fields we introduce the solenoidal subspace

$$\mathcal{J}(\Omega) := \left\{ (E, H) \in L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3) : \operatorname{div}(\varepsilon_0 E) = \operatorname{div}(\mu_0 H) = 0, (H \cdot n)|_{\partial\Omega} = 0 \right\}.$$

## Lemma

There holds  $\ker(\mathcal{A})^\perp = \mathcal{J}(\Omega)$ . In particular,  $\mathcal{J}(\Omega)$  is a closed subspace left stable by  $\mathcal{A}$ .

Physically, the Maxwell operator is  $\mathcal{A}_\perp := \mathcal{A}|_{\mathcal{J}(\Omega)}$ .



# Divergence free fields and the magnetic boundary condition

To look for divergence-free fields we introduce the solenoidal subspace

$$\mathcal{J}(\Omega) := \left\{ (E, H) \in L^2(\Omega, \mathbb{R}^3 \times \mathbb{R}^3) : \operatorname{div}(\varepsilon_0 E) = \operatorname{div}(\mu_0 H) = 0, (H \cdot n)|_{\partial\Omega} = 0 \right\}.$$

## Lemma

There holds  $\ker(\mathcal{A})^\perp = \mathcal{J}(\Omega)$ . In particular,  $\mathcal{J}(\Omega)$  is a closed subspace left stable by  $\mathcal{A}$ .

Physically, the Maxwell operator is  $\mathcal{A}_\perp := \mathcal{A}|_{\mathcal{J}(\Omega)}$ .



# Some bibliography on waveguides

## Quantum waveguides:



G. BOUCHITTÉ, M. L. MASCARENHAS, L. TRABUCHO,  
ESAIM: COCV, (2007)



P. BRIET, H. KOVAŘÍK, G. RAIKOV, E. SOCCORSI,  
COMM. IN PDE (2009)



P. DUCLOS, P. EXNER,  
REV. MATH. PHYS. (1995)



J. GOLDSTONE, R. L. JAFFE,  
PHYS. REV. B (1992)

## Relativistic quantum waveguides:



W. BORRELLI, P. BRIET, D. KREJČÍŘÍK, T. O.-B.,  
ANN. HENRI POINCARÉ, (2022)



L. LE TREUST, T. O.-B., N. RAYMOND,  
PREPRINT, (2024).

## Electromagnetic waveguides:



N. FILONOV,  
ST. PETERSBURG MATH. J., (2019) + (2021)

## 1 Introduction

- What is a waveguide ?
- The Maxwell operator

## 2 Main results

- Essential spectrum
- Discrete spectrum

## 3 Geometric transforms and Birman-Schwinger principle

- The Piola transform
- Essential spectrum of constantly twisted waveguide
- A Resolvent identity

## 4 On the discrete spectrum

- A Poincaré inequality
- On the existence of discrete spectrum

## 5 Geometric condition $X \neq 0$

# Essential spectrum

Recall that we have assumed

$$\lim_{|s| \rightarrow +\infty} \kappa(s) = 0, \quad \lim_{|s| \rightarrow +\infty} \theta'(s) = \beta, \quad \beta \in \mathbb{R}.$$

## Theorem (structure of the essential spectrum)

There exists  $a_\beta > 0$  such that

$$Sp_{\text{ess}}(\mathcal{A}) = (-\infty, -a_\beta] \cup \{0\} \cup [a_\beta, +\infty).$$

If  $\beta = 0$ , one has  $a_\beta = \sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}}$ .

# Essential spectrum

Recall that we have assumed

$$\lim_{|s| \rightarrow +\infty} \kappa(s) = 0, \quad \lim_{|s| \rightarrow +\infty} \theta'(s) = \beta, \quad \beta \in \mathbb{R}.$$

## Theorem (structure of the essential spectrum)

There exists  $a_\beta > 0$  such that

$$Sp_{\text{ess}}(\mathcal{A}) = (-\infty, -a_\beta] \cup \{0\} \cup [a_\beta, +\infty).$$

If  $\beta = 0$ , one has  $a_\beta = \sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}}$ .

**Problem**: Focus on  $\mathcal{A}_\perp$  and we need to get rid of 0 !

# Essential spectrum (2/2)

Assume further

$$\theta' = 0 \quad (\text{no twist}), \quad \left( \sup_{y \in \omega} |y| \right) \|\kappa\|_{L^\infty(\mathbb{R})} < \frac{1}{2}.$$

# Essential spectrum (2/2)

Assume further

$$\theta' = 0 \quad (\text{no twist}), \quad \left( \sup_{y \in \omega} |y| \right) \|\kappa\|_{L^\infty(\mathbb{R})} < \frac{1}{2}.$$

## Proposition

There holds

$$\sigma_{\text{ess}}(\mathcal{A}_\perp) = \left( -\infty, -\sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}} \right] \cup \left[ -\sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}}, +\infty \right).$$

# Essential spectrum (2/2)

Assume further

$$\theta' = 0 \quad (\text{no twist}), \quad \left( \sup_{y \in \omega} |y| \right) \|\kappa\|_{L^\infty(\mathbb{R})} < \frac{1}{2}.$$

## Proposition

There holds

$$\sigma_{\text{ess}}(\mathcal{A}_\perp) = \left( -\infty, -\sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}} \right] \cup \left[ -\sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}}, +\infty \right).$$

The key-point is a Poincaré-type inequality of the form:

$$\|E\|_{L^2_{\varepsilon_0}(\Omega)} + \|H\|_{L^2_{\mu_0}(\Omega)} \leq C(\|\mu_0^{-1} \operatorname{curl} E\|_{L^2_{\mu_0}(\Omega)} + \|\varepsilon_0^{-1} \operatorname{curl} H\|_{L^2_{\varepsilon_0}(\Omega)}).$$

for some constant  $C > 0$  uniform in  $(E, H)^\top \in \mathcal{J}(\Omega) \cap \operatorname{Dom}(\mathcal{A})$ .



# Essential spectrum (2/2)

Assume further

$$\theta' = 0 \quad (\text{no twist}), \quad \left( \sup_{y \in \omega} |y| \right) \|\kappa\|_{L^\infty(\mathbb{R})} < \frac{1}{2}.$$

## Proposition

There holds

$$\sigma_{\text{ess}}(\mathcal{A}_\perp) = \left( -\infty, -\sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}} \right] \cup \left[ -\sqrt{\frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0}}, +\infty \right).$$

Should be true with **non-zero twist** and a condition involving  $\|\theta' - \beta\|_{L^\infty(\mathbb{R})}$  but ask for tedious computations...

# Discrete spectrum

- $\psi$  is a normalized eigenmode associated with  $\lambda_2^N(\omega)$ , the first non-trivial Neumann eigenvalue in  $\omega$ .

# Discrete spectrum

- $\psi$  is a normalized eigenmode associated with  $\lambda_2^N(\omega)$ , the first non-trivial Neumann eigenvalue in  $\omega$ .
- Set

$$k^\theta := \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} k_1 \\ k_2 \end{pmatrix}, \quad X := \int_{\partial\omega} n(y) |\psi(y)|^2 d\sigma(y).$$

# Discrete spectrum

- $\psi$  is a normalized eigenmode associated with  $\lambda_2^N(\omega)$ , the first non-trivial Neumann eigenvalue in  $\omega$ .
- Set

$$k^\theta := \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} k_1 \\ k_2 \end{pmatrix}, \quad X := \int_{\partial\omega} n(y) |\psi(y)|^2 d\sigma(y).$$

## Theorem (existence of discrete spectrum)

Assume  $\theta' = 0$  (no twist) and that the Poincaré inequality holds. Let  $\kappa \in L^1(\mathbb{R})$  be such that

$$X \cdot \left( \int_{\mathbb{R}} k^\theta(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

then

$$\sigma_{\text{dis}}(\mathcal{A}) \neq \emptyset.$$

Here  $b := \sup_{y \in \Omega} (|y|)$ .

# Some remarks on the geometric condition

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \| \kappa \|_{L^\infty(\mathbb{R})}}{1 - b \| \kappa \|_{L^\infty(\mathbb{R})}} \| \kappa \|_{L^1(\mathbb{R})}$$

- It is necessary that  $X \neq 0$  and  $\left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) \neq 0$  for this inequality to hold.

# Some remarks on the geometric condition

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \| \kappa \|_{L^\infty(\mathbb{R})}}{1 - b \| \kappa \|_{L^\infty(\mathbb{R})}} \| \kappa \|_{L^1(\mathbb{R})}$$

- It is necessary that  $X \neq 0$  and  $\left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) \neq 0$  for this inequality to hold.
- In this case, as  $\theta$  is a constant, there is an optimal choice of  $\theta_{\star}$  maximize the quantity

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta_{\star}}(s) ds \right)$$

# Some remarks on the geometric condition

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \| \kappa \|_{L^\infty(\mathbb{R})}}{1 - b \| \kappa \|_{L^\infty(\mathbb{R})}} \| \kappa \|_{L^1(\mathbb{R})}$$

- It is necessary that  $X \neq 0$  and  $\left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) \neq 0$  for this inequality to hold.
- In this case, as  $\theta$  is a constant, there is an optimal choice of  $\theta_{\star}$  maximize the quantity

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta_{\star}}(s) ds \right) > 0 \quad \text{for } \theta = \theta_{\star}.$$

# Some remarks on the geometric condition

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \| \kappa \|_{L^\infty(\mathbb{R})}}{1 - b \| \kappa \|_{L^\infty(\mathbb{R})}} \| \kappa \|_{L^1(\mathbb{R})}$$

- It is necessary that  $X \neq 0$  and  $\left( \int_{\mathbb{R}} k^{\theta}(s) ds \right) \neq 0$  for this inequality to hold.
- In this case, as  $\theta$  is a constant, there is an optimal choice of  $\theta_{\star}$  maximize the quantity

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta_{\star}}(s) ds \right)$$

- It means that there is an optimal way to attach the cross-section  $\omega$  to the curve  $\Gamma$ .



# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta*}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta*}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

- Assume  $X \neq 0$  and  $\int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds \neq 0$ .

# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta*}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

- Assume  $X \neq 0$  and  $\int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds \neq 0$ .
- For  $\delta > 0$ , one can define

$$\gamma_\delta : \mathbb{R} \rightarrow \mathbb{R}^3, \quad \gamma_\delta(s) = \frac{1}{\delta} \gamma(\delta s).$$

# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta*}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

- Assume  $X \neq 0$  and  $\int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds \neq 0$ .
- For  $\delta > 0$ , one can define

$$\gamma_\delta : \mathbb{R} \rightarrow \mathbb{R}^3, \quad \gamma_\delta(s) = \frac{1}{\delta} \gamma(\delta s).$$

- The associated curvatures verify  $k_1^\delta(s) = \delta k_1(\delta s)$ ,  $k_2^\delta(s) = \delta k_2(\delta s)$  but

$$\int_{\mathbb{R}} \begin{pmatrix} k_1^\delta(s) \\ k_2^\delta(s) \end{pmatrix} ds = \int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds, \quad \|\kappa_\delta\|_{L^1(\mathbb{R})} = \|\kappa\|_{L^1(\mathbb{R})}.$$

# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k_{\delta}^{\theta*}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \|\kappa_{\delta}\|_{L^{\infty}(\mathbb{R})}}{1 - b \|\kappa_{\delta}\|_{L^{\infty}(\mathbb{R})}} \|\kappa_{\delta}\|_{L^1(\mathbb{R})}$$

- Assume  $X \neq 0$  and  $\int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds \neq 0$ .
- For  $\delta > 0$ , one can define

$$\gamma_{\delta} : \mathbb{R} \rightarrow \mathbb{R}^3, \quad \gamma_{\delta}(s) = \frac{1}{\delta} \gamma(\delta s).$$

- The associated curvatures verify  $k_1^{\delta}(s) = \delta k_1(\delta s)$ ,  $k_2^{\delta}(s) = \delta k_2(\delta s)$  but

$$\int_{\mathbb{R}} \begin{pmatrix} k_1^{\delta}(s) \\ k_2^{\delta}(s) \end{pmatrix} ds = \int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds, \quad \|\kappa_{\delta}\|_{L^1(\mathbb{R})} = \|\kappa\|_{L^1(\mathbb{R})}.$$

# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta \star}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \delta \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b\delta \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

- Assume  $X \neq 0$  and  $\int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds \neq 0$ .
- For  $\delta > 0$ , one can define

$$\gamma_\delta : \mathbb{R} \rightarrow \mathbb{R}^3, \quad \gamma_\delta(s) = \frac{1}{\delta} \gamma(\delta s).$$

- The associated curvatures verify  $k_1^\delta(s) = \delta k_1(\delta s)$ ,  $k_2^\delta(s) = \delta k_2(\delta s)$  but

$$\int_{\mathbb{R}} \begin{pmatrix} k_1^\delta(s) \\ k_2^\delta(s) \end{pmatrix} ds = \int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds, \quad \|\kappa_\delta\|_{L^1(\mathbb{R})} = \|\kappa\|_{L^1(\mathbb{R})}.$$

# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta \star}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \delta \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b\delta \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

When  $\delta \rightarrow 0$ , the condition is satisfied.

# Slightly curved waveguides

Geometric condition:

$$X \cdot \left( \int_{\mathbb{R}} k^{\theta \star}(s) ds \right) > \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \delta \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b\delta \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}$$

When  $\delta \rightarrow 0$ , the condition is satisfied.

Given a curve for which

- $\int_{\mathbb{R}} \begin{pmatrix} k_1(s) \\ k_2(s) \end{pmatrix} ds \neq 0$ ,
- a cross-section  $\omega$  for which  $X \neq 0$ ,

one can always build a waveguide which traps electromagnetic modes.



## 1 Introduction

- What is a waveguide ?
- The Maxwell operator

## 2 Main results

- Essential spectrum
- Discrete spectrum

## 3 Geometric transforms and Birman-Schwinger principle

- The Piola transform
- Essential spectrum of constantly twisted waveguide
- A Resolvent identity

## 4 On the discrete spectrum

- A Poincaré inequality
- On the existence of discrete spectrum

## 5 Geometric condition $X \neq 0$

# (Unitary) Piola transform

**Key-point**: Rewrite the problem in a straight waveguide  $\Omega_0 = \mathbb{R} \times \omega$ .

$$\mathbb{U} : L^2_{\varepsilon_0, \mu_0}(\Omega) \rightarrow L^2_{\varepsilon, \mu}(\Omega_0), \quad \begin{pmatrix} E \\ H \end{pmatrix} \mapsto \begin{pmatrix} J_{\Phi}^{\top}(E \circ \Phi) \\ J_{\Phi}^{\top}(H \circ \Phi) \end{pmatrix}$$

$$\text{where } \varepsilon = \varepsilon_0 \mathbb{G}^{-1}, \quad \mu = \mu_0 \mathbb{G}^{-1} \text{ with } \mathbb{G} = \frac{J_{\Phi} J_{\Phi}^{\top}}{\det J_{\Phi}}.$$

# (Unitary) Piola transform

**Key-point**: Rewrite the problem in a straight waveguide  $\Omega_0 = \mathbb{R} \times \omega$ .

$$\mathbb{U} : L^2_{\varepsilon_0, \mu_0}(\Omega) \rightarrow L^2_{\varepsilon, \mu}(\Omega_0), \quad \begin{pmatrix} E \\ H \end{pmatrix} \mapsto \begin{pmatrix} J_{\Phi}^{\top}(E \circ \Phi) \\ J_{\Phi}^{\top}(H \circ \Phi) \end{pmatrix}$$

$$\text{where } \varepsilon = \varepsilon_0 \mathbb{G}^{-1}, \quad \mu = \mu_0 \mathbb{G}^{-1} \text{ with } \mathbb{G} = \frac{J_{\Phi} J_{\Phi}^{\top}}{\det J_{\Phi}}.$$

Unitary equivalent operator  $\widehat{\mathcal{A}} := \mathbb{U} \mathcal{A} \mathbb{U}^*$  acting in  $L^2_{\varepsilon, \mu}(\Omega_0) \rightarrow L^2_{\varepsilon, \mu}(\Omega_0)$ .

$$\mathcal{A} = \begin{pmatrix} 0 & i\varepsilon_0^{-1} \text{curl} \\ -i\mu_0^{-1} \text{curl} & 0 \end{pmatrix} \quad \widehat{\mathcal{A}} = \begin{pmatrix} 0 & i\varepsilon^{-1} \text{curl} \\ -i\mu^{-1} \text{curl} & 0 \end{pmatrix}$$

# (Unitary) Piola transform

**Key-point**: Rewrite the problem in a straight waveguide  $\Omega_0 = \mathbb{R} \times \omega$ .

$$\mathbb{U} : L^2_{\varepsilon_0, \mu_0}(\Omega) \rightarrow L^2_{\varepsilon, \mu}(\Omega_0), \quad \begin{pmatrix} E \\ H \end{pmatrix} \mapsto \begin{pmatrix} J_{\Phi}^{\top}(E \circ \Phi) \\ J_{\Phi}^{\top}(H \circ \Phi) \end{pmatrix}$$

$$\text{where } \varepsilon = \varepsilon_0 \mathbb{G}^{-1}, \quad \mu = \mu_0 \mathbb{G}^{-1} \text{ with } \mathbb{G} = \frac{J_{\Phi} J_{\Phi}^{\top}}{\det J_{\Phi}}.$$

Unitary equivalent operator  $\widehat{\mathcal{A}} := \mathbb{U} \mathcal{A} \mathbb{U}^*$  acting in  $L^2_{\varepsilon, \mu}(\Omega_0) \rightarrow L^2_{\varepsilon, \mu}(\Omega_0)$ .

$$\mathcal{A} = \begin{pmatrix} 0 & i\varepsilon_0^{-1} \text{curl} \\ -i\mu_0^{-1} \text{curl} & 0 \end{pmatrix} \quad \widehat{\mathcal{A}} = \begin{pmatrix} 0 & i\varepsilon^{-1} \text{curl} \\ -i\mu^{-1} \text{curl} & 0 \end{pmatrix}$$

The Piola transform preserves the Maxwell structure:

- $\text{Dom}(\widehat{A}) = H_0(\text{curl}, \Omega_0) \times H(\text{curl}, \Omega_0)$ ,
- $\mathbb{U} \mathcal{J}(\Omega) = \mathcal{J}_{\varepsilon, \mu}(\Omega_0)$  where

$$\mathcal{J}_{\varepsilon, \mu}(\Omega_0) := \{(E, H)^{\top} \in L^2_{\varepsilon, \mu}(\Omega_0, \mathbb{R}^3 \times \mathbb{R}^3) : \text{div}(\varepsilon E) = \text{div}(\mu H) = 0, \\ (\mu H) \cdot n = 0\}.$$

# (Unitary) Piola transform

**Key-point**: Rewrite the problem in a straight waveguide  $\Omega_0 = \mathbb{R} \times \omega$ .


$$\mathbb{U} : L^2_{\varepsilon_0, \mu_0}(\Omega) \rightarrow L^2_{\varepsilon, \mu}(\Omega_0), \quad \begin{pmatrix} \mathbf{E} \\ \mathbf{H} \end{pmatrix} \mapsto \begin{pmatrix} J_{\Phi}^{\top}(\mathbf{E} \circ \Phi) \\ J_{\Phi}^{\top}(\mathbf{H} \circ \Phi) \end{pmatrix}$$

where  $\varepsilon = \varepsilon_0 \mathbb{G}^{-1}$ ,  $\mu = \mu_0 \mathbb{G}^{-1}$  with  $\mathbb{G} = \frac{J_{\Phi} J_{\Phi}^{\top}}{\det J_{\Phi}}$ .

Unitary equivalent operator  $\widehat{\mathcal{A}} := \mathbb{U} \mathcal{A} \mathbb{U}^*$  acting in  $L^2_{\varepsilon, \mu}(\Omega_0) \rightarrow L^2_{\varepsilon, \mu}(\Omega_0)$ .

$$\mathcal{A} = \begin{pmatrix} 0 & i\varepsilon_0^{-1} \text{curl} \\ -i\mu_0^{-1} \text{curl} & 0 \end{pmatrix} \quad \widehat{\mathcal{A}} = \begin{pmatrix} 0 & i\varepsilon^{-1} \text{curl} \\ -i\mu^{-1} \text{curl} & 0 \end{pmatrix}$$

Look for electromagnetic waves with  
non-constant permittivity and permeability.



# Essential spectrum - constant twist $\beta$

Assume  $\kappa = 0$ ,  $\theta' = \beta$ .

Observe that

$$\mathbb{G}_\beta(s, y_2, y_3) = \mathbb{G}_\beta(y_2, y_3) := \begin{pmatrix} 1 + \beta^2 |y|^2 & -y_3 \beta & y_2 \beta \\ -y_3 \beta & 1 & 0 \\ y_2 \beta & 0 & 1 \end{pmatrix}$$

depends only on  $(y_2, y_3) \in \omega$ .

# Essential spectrum - constant twist $\beta$

Assume  $\kappa = 0$ ,  $\theta' = \beta$ .

Observe that

$$\mathbb{G}_\beta(s, y_2, y_3) = \mathbb{G}_\beta(y_2, y_3) := \begin{pmatrix} 1 + \beta^2 |y|^2 & -y_3 \beta & y_2 \beta \\ -y_3 \beta & 1 & 0 \\ y_2 \beta & 0 & 1 \end{pmatrix}$$

depends only on  $(y_2, y_3) \in \omega$ . There exists  $a_\beta > 0$  such that

$$\sigma(\mathcal{A}_\beta) = \sigma_{\text{ess}}(\mathcal{A}_\beta) = (-\infty, -a_\beta] \cup \{0\} \cup [a_\beta, +\infty)$$

# Essential spectrum - constant twist $\beta$

Assume  $\kappa = 0$ ,  $\theta' = \beta$ .

Observe that

$$\mathbb{G}_\beta(s, y_2, y_3) = \mathbb{G}_\beta(y_2, y_3) := \begin{pmatrix} 1 + \beta^2 |y|^2 & -y_3 \beta & y_2 \beta \\ -y_3 \beta & 1 & 0 \\ y_2 \beta & 0 & 1 \end{pmatrix}$$

depends only on  $(y_2, y_3) \in \omega$ . There exists  $a_\beta > 0$  such that

$$\sigma(\mathcal{A}_\beta) = \sigma_{\text{ess}}(\mathcal{A}_\beta) = (-\infty, -a_\beta] \cup \{0\} \cup [a_\beta, +\infty)$$





# Essential spectrum - constant twist $\beta$

Assume  $\kappa = 0$ ,  $\theta' = \beta$ .

Observe that

$$\mathbb{G}_\beta(s, y_2, y_3) = \mathbb{G}_\beta(y_2, y_3) := \begin{pmatrix} 1 + \beta^2 |y|^2 & -y_3 \beta & y_2 \beta \\ -y_3 \beta & 1 & 0 \\ y_2 \beta & 0 & 1 \end{pmatrix}$$

depends only on  $(y_2, y_3) \in \omega$ . There exists  $a_\beta > 0$  such that

$$\sigma(\mathcal{A}_\beta) = \sigma_{\text{ess}}(\mathcal{A}_\beta) = (-\infty, -a_\beta] \cup \{0\} \cup [a_\beta, +\infty)$$



N. FILONOV,

ST. PETERSBURG MATH. J., (2019)

## Comparison of $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{A}}_\beta$

We would like to compare the resolvents of  $\widehat{\mathcal{A}}$  and  $\widehat{\mathcal{A}}_\beta$ .

# Comparison of $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{A}}_\beta$

We would like to compare the resolvents of  $\widehat{\mathcal{A}}$  and  $\widehat{\mathcal{A}}_\beta$ .

**Problem**: They do not act in the same Hilbert space !

We adapt an approach developed by Weder



R. WEDER,

SPECTRAL AND SCATTERING THEORY FOR WAVE PROPAGATION IN PERTURBED  
STRATIFIED MEDIA. (1991)

# Comparison of $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{A}}_\beta$

We would like to compare the resolvents of  $\widehat{\mathcal{A}}$  and  $\widehat{\mathcal{A}}_\beta$ .

**Problem**: They do not act in the same Hilbert space !

We adapt an approach developed by Weder



R. WEDER,

SPECTRAL AND SCATTERING THEORY FOR WAVE PROPAGATION IN PERTURBED STRATIFIED MEDIA. (1991)

Another unitary transform

$$\mathbb{V} : L^2_{\varepsilon, \mu}(\Omega_0) \rightarrow L^2_{\varepsilon_0, \mu_0}(\Omega_0), \quad \begin{pmatrix} E \\ H \end{pmatrix} \mapsto \begin{pmatrix} \mathbb{G}^{-1/2} E \\ \mathbb{G}^{-1/2} H \end{pmatrix}$$

# Comparison of $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{A}}_\beta$

We would like to compare the resolvents of  $\widehat{\mathcal{A}}$  and  $\widehat{\mathcal{A}}_\beta$ .

**Problem**: They do not act in the same Hilbert space !

We adapt an approach developed by Weder



R. WEDER,

SPECTRAL AND SCATTERING THEORY FOR WAVE PROPAGATION IN PERTURBED STRATIFIED MEDIA. (1991)

Another unitary transform

$$\mathbb{V}_\beta : L^2_{\varepsilon_\beta, \mu_\beta}(\Omega_0) \rightarrow L^2_{\varepsilon_0, \mu_0}(\Omega_0), \quad \begin{pmatrix} E \\ H \end{pmatrix} \mapsto \begin{pmatrix} \mathbb{G}_\beta^{-1/2} E \\ \mathbb{G}_\beta^{-1/2} H \end{pmatrix}$$

# Comparison of $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{A}}_\beta$

We would like to compare the resolvents of  $\widehat{\mathcal{A}}$  and  $\widehat{\mathcal{A}}_\beta$ .

**Problem**: They do not act in the same Hilbert space !

We adapt an approach developed by Weder



R. WEDER,

SPECTRAL AND SCATTERING THEORY FOR WAVE PROPAGATION IN PERTURBED STRATIFIED MEDIA. (1991)


Another unitary transform

$$\mathbb{V}_\beta : L^2_{\varepsilon_\beta, \mu_\beta}(\Omega_0) \rightarrow L^2_{\varepsilon_0, \mu_0}(\Omega_0), \quad \begin{pmatrix} E \\ H \end{pmatrix} \mapsto \begin{pmatrix} \mathbb{G}_\beta^{-1/2} E \\ \mathbb{G}_\beta^{-1/2} H \end{pmatrix}$$

Define the unitary equivalent operators acting in  $L^2_{\varepsilon_0, \mu_0}(\Omega_0)$ :

- $\widetilde{\mathcal{A}} := \mathbb{V} \widehat{\mathcal{A}} \mathbb{V}^*,$
- $\widetilde{\mathcal{A}}_\beta := \mathbb{V}_\beta \widehat{\mathcal{A}}_\beta \mathbb{V}_\beta^*$

# Comparison of operators: a picture !



# Resolvent identity

The rest of the proof is by investigating, for  $\lambda \in \rho(\widetilde{\mathcal{A}}) \cap \rho(\widetilde{\mathcal{A}}_\beta)$ , the resolvents

$$\widetilde{R}(\lambda) := (\widetilde{\mathcal{A}} - \lambda)^{-1}, \quad \widetilde{R}_\beta(\lambda) := (\widetilde{\mathcal{A}}_\beta - \lambda)^{-1}$$

and remarking that



# Resolvent identity

The rest of the proof is by investigating, for  $\lambda \in \rho(\widetilde{\mathcal{A}}) \cap \rho(\widetilde{\mathcal{A}}_\beta)$ , the resolvents

$$\tilde{R}(\lambda) := (\widetilde{\mathcal{A}} - \lambda)^{-1}, \quad \tilde{R}_\beta(\lambda) := (\widetilde{\mathcal{A}}_\beta - \lambda)^{-1}$$

and remarking that

$$\tilde{R}(\lambda) = \mathbb{V} \mathcal{I}_\mathbb{G}^{-1} \mathcal{I}_\beta \mathbb{V}_\beta^* \tilde{R}_\beta(\lambda) \mathcal{Q}_\beta(\lambda)^{-1} \mathcal{I}_\beta \mathbb{V}_\beta^* \mathbb{V} \mathcal{I}_\mathbb{G}^{-1}$$

# Resolvent identity

The rest of the proof is by investigating, for  $\lambda \in \rho(\widetilde{\mathcal{A}}) \cap \rho(\widetilde{\mathcal{A}}_\beta)$ , the resolvents

$$\widetilde{R}(\lambda) := (\widetilde{\mathcal{A}} - \lambda)^{-1}, \quad \widetilde{R}_\beta(\lambda) := (\widetilde{\mathcal{A}}_\beta - \lambda)^{-1}$$

and remarking that

$$\widetilde{R}_\beta(\lambda) = \mathbb{V}_\beta \mathcal{I}_\beta^{-1} \mathcal{I}_G \mathbb{V}^* \widetilde{R}(\lambda) \mathcal{Q}(\lambda)^{-1} \mathcal{I}_G \mathbb{V}_G^* \mathbb{V}_\beta \mathcal{I}_\beta^{-1}.$$

# Resolvent identity

The rest of the proof is by investigating, for  $\lambda \in \rho(\widetilde{\mathcal{A}}) \cap \rho(\widetilde{\mathcal{A}}_\beta)$ , the resolvents

$$\widetilde{R}(\lambda) := (\widetilde{\mathcal{A}} - \lambda)^{-1}, \quad \widetilde{R}_\beta(\lambda) := (\widetilde{\mathcal{A}}_\beta - \lambda)^{-1}$$

and remarking that

$$\widetilde{R}_\beta(\lambda) = \mathbb{V}_\beta \mathcal{I}_\beta^{-1} \mathcal{I}_G \mathbb{V}^* \widetilde{R}(\lambda) \mathcal{Q}(\lambda)^{-1} \mathcal{I}_G \mathbb{V}_G^* \mathbb{V}_\beta \mathcal{I}_\beta^{-1}.$$

- $\mathcal{Q}(\lambda) = \mathcal{Q}(\lambda_0)(Id + \mathcal{K}(\lambda))$  with  $\mathcal{K}(\lambda)$  is a compact operator.
- $\mathcal{Q}_\beta(\lambda) = \mathcal{Q}_\beta(\lambda_0)(Id + \mathcal{K}_\beta(\lambda))$  with  $\mathcal{K}_\beta(\lambda)$  is a compact operator.
- It yields a Birman-Schwinger principle.
- We conclude studying the analytic family of compact operators  $(\lambda \mapsto \mathcal{K}_\beta(\lambda))$  and the finitely meromorphic family of compact operators  $(\lambda \mapsto \mathcal{K}(\lambda))$ .

## 1 Introduction

- What is a waveguide ?
- The Maxwell operator

## 2 Main results

- Essential spectrum
- Discrete spectrum

## 3 Geometric transforms and Birman-Schwinger principle

- The Piola transform
- Essential spectrum of constantly twisted waveguide
- A Resolvent identity

## 4 On the discrete spectrum

- A Poincaré inequality
- On the existence of discrete spectrum

## 5 Geometric condition $X \neq 0$

# Poincaré inequality

$$\text{Assume } \theta' = 0, \quad b \| \kappa \|_{L^\infty(\mathbb{R})} < \frac{1}{2}.$$

There exists  $C > 0$  such that for all  
 $(E, H)^\top \in \mathcal{J}_{\varepsilon, \mu}(\Omega_0) \cap (H_0(\text{curl}, \Omega_0) \times H(\text{curl}, \Omega_0))$ :

$$\|E\|_{L_\varepsilon^2(\Omega_0)} + \|H\|_{L_\mu^2(\Omega_0)} \leq C \left( \|\mu^{-1} \text{curl} E\|_{L_\mu^2(\Omega_0)} + \|\varepsilon^{-1} \text{curl} H\|_{L_\varepsilon^2(\Omega_0)} \right)$$

# Poincaré inequality

$$\text{Assume } \theta' = 0, \quad b \| \kappa \|_{L^\infty(\mathbb{R})} < \frac{1}{2}.$$

There exists  $C > 0$  such that for all  $(E, H)^\top \in \mathcal{J}_{\varepsilon, \mu}(\Omega_0) \cap (H_0(\text{curl}, \Omega_0) \times H(\text{curl}, \Omega_0))$ :

$$\|E\|_{L_\varepsilon^2(\Omega_0)} + \|H\|_{L_\mu^2(\Omega_0)} \leq C \left( \|\mu^{-1} \text{curl} E\|_{L_\mu^2(\Omega_0)} + \|\varepsilon^{-1} \text{curl} H\|_{L_\varepsilon^2(\Omega_0)} \right)$$

- It is a consequence of the Birman-Schwinger principle.
- A consequence is that  $0 \notin \widehat{Sp}(\mathcal{A}_\perp)$ .

# Strategy of the proof (discrete spectrum)

- Work with the quadratic form associated with  $(\widehat{\mathcal{A}}_{\perp})^2$

$$q[E, H] := \int_{\Omega_0} \left( \mu^{-1} \operatorname{curl} E \cdot \operatorname{curl} E + \varepsilon^{-1} \operatorname{curl} H \cdot H \right) dx,$$
$$(E, H) \in \operatorname{Dom}(q) := \operatorname{Dom}(\widehat{\mathcal{A}}) \cap \mathcal{J}_{\varepsilon, \mu}(\Omega_0).$$

# Strategy of the proof (discrete spectrum)

- Work with the quadratic form associated with  $(\widehat{\mathcal{A}}_{\perp})^2$

$$q[E, H] := \int_{\Omega_0} \left( \mu^{-1} \operatorname{curl} E \cdot \operatorname{curl} E + \varepsilon^{-1} \operatorname{curl} H \cdot H \right) dx,$$
$$(E, H) \in \operatorname{Dom}(q) := \operatorname{Dom}(\widehat{\mathcal{A}}) \cap \mathcal{J}_{\varepsilon, \mu}(\Omega_0).$$

- $\beta = \theta' = 0$  gives  $\varepsilon, \mu$  are diagonal !



# Strategy of the proof (discrete spectrum)

- Work with the quadratic form associated with  $(\widehat{\mathcal{A}}_{\perp})^2$

$$q[E, H] := \int_{\Omega_0} \left( \mu^{-1} \operatorname{curl} E \cdot \operatorname{curl} E + \varepsilon^{-1} \operatorname{curl} H \cdot H \right) dx,$$

$$(E, H) \in \operatorname{Dom}(q) := \operatorname{Dom}(\widehat{\mathcal{A}}) \cap \mathcal{J}_{\varepsilon, \mu}(\Omega_0).$$

- $\beta = \theta' = 0$  gives  $\varepsilon, \mu$  are diagonal !
- Look for a test function of the form  $(E, 0)^{\top} \neq (0, 0)^{\top}$  verifying:

$$q[E, 0] - \frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0} \|E\|_{L_{\varepsilon}^2(\Omega_0)}^2 < 0.$$

# Strategy of the proof (discrete spectrum)

- Work with the quadratic form associated with  $(\widehat{\mathcal{A}}_{\perp})^2$

$$q[E, H] := \int_{\Omega_0} \left( \mu^{-1} \operatorname{curl} E \cdot \operatorname{curl} E + \varepsilon^{-1} \operatorname{curl} H \cdot H \right) dx,$$

$$(E, H) \in \operatorname{Dom}(q) := \operatorname{Dom}(\widehat{\mathcal{A}}) \cap \mathcal{J}_{\varepsilon, \mu}(\Omega_0).$$

- $\beta = \theta' = 0$  gives  $\varepsilon, \mu$  are diagonal !
- Look for a test function of the form  $(E, 0)^{\top} \neq (0, 0)^{\top}$  verifying:

$$q[E, 0] - \frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0} \|E\|_{L^2_{\varepsilon}(\Omega_0)}^2 < 0.$$

- Conclude with the min-max principle.

# Choice of the trial function

Set  $F$  of the form

$$F(s, y) = \varphi(s) \begin{pmatrix} 0 \\ \partial_3 \psi(y) \\ -\partial_2 \psi(y) \end{pmatrix}.$$

**Problem**:  $(F, 0)^\top \notin \text{Dom}(q)$  because  $\text{div}(\varepsilon F) \neq 0$

We correct it as

$$E = F + \nabla u$$

with  $u \in H_0^1(\Omega)$  weak-solution of

$$\begin{cases} \text{div}(\varepsilon \nabla u) = -\text{div}(\varepsilon F) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

With this choice we control  $\|\nabla u\|_{L_\varepsilon^2(\Omega_0)}$  using  $\text{div}(F) = 0$  and compute

$$q[E, 0] - \frac{\lambda_2^N(\omega)}{\varepsilon_0 \mu_0} \|E\|_{L^2(\Omega_0)}^2 < -\chi \cdot \left( \int_{\mathbb{R}} k^\theta(s) ds \right) + \frac{2\lambda_2^N(\omega)}{\varepsilon_0} \frac{b^2 \|\kappa\|_{L^\infty(\mathbb{R})}}{1 - b \|\kappa\|_{L^\infty(\mathbb{R})}} \|\kappa\|_{L^1(\mathbb{R})}.$$

## 1 Introduction

- What is a waveguide ?
- The Maxwell operator

## 2 Main results

- Essential spectrum
- Discrete spectrum

## 3 Geometric transforms and Birman-Schwinger principle

- The Piola transform
- Essential spectrum of constantly twisted waveguide
- A Resolvent identity

## 4 On the discrete spectrum

- A Poincaré inequality
- On the existence of discrete spectrum

## 5 Geometric condition $X \neq 0$

- Most of the domains  $\omega$  for which one can compute explicitly  $\lambda_2^N(\omega)$  gives  $X = 0$  (disks, rectangles...).

- Most of the domains  $\omega$  for which one can compute explicitly  $\lambda_2^N(\omega)$  gives  $X = 0$  (disks, rectangles...).
- One can prove that  $X = 0$  as long as the domain has a rotational symmetry of angle  $\varphi \in (0, 2\pi)$ .

- Most of the domains  $\omega$  for which one can compute explicitly  $\lambda_2^N(\omega)$  gives  $X = 0$  (disks, rectangles...).
- One can prove that  $X = 0$  as long as the domain has a rotational symmetry of angle  $\varphi \in (0, 2\pi)$ .
- For  $\omega$  being an isosceles right triangle  $X \neq 0$  ! The modes can be computed by symmetry using the one of the square.

- Most of the domains  $\omega$  for which one can compute explicitly  $\lambda_2^N(\omega)$  gives  $X = 0$  (disks, rectangles...).
- One can prove that  $X = 0$  as long as the domain has a rotational symmetry of angle  $\varphi \in (0, 2\pi)$ .
- For  $\omega$  being an isosceles right triangle  $X \neq 0$  ! The modes can be computed by symmetry using the one of the square.
- Plenty many numerical examples of  $X \neq 0$ ...





Domains for which  $X \neq 0$ 


Figure: A Dumbbell domain.  $X_{\text{dumbbell}} = \begin{pmatrix} -0.1255 \\ 0 \end{pmatrix}$

Domains for which  $X \neq 0$ 

**Figure:** Rectangle of base  $2\pi$  and height  $\pi$  with bumps. The bumps are half-disks of varying radius and positions.

$$X_{\text{blue}} = \begin{pmatrix} 0.00315511 \\ 0.00126526 \end{pmatrix}, \quad X_{\text{yellow}} = \begin{pmatrix} -0.000798985 \\ 0.00301619 \end{pmatrix}$$

$$X_{\text{red}} = \begin{pmatrix} -0.0152116 \\ 0.00285829 \end{pmatrix}, \quad X_{\text{green}} = \begin{pmatrix} 0.0000322362 \\ 0.000374781 \end{pmatrix}$$

Domains for which  $X \neq 0$ 

**Figure:** Rectangle of base  $2\pi$  and height  $\pi$  with bumps. The bumps are half-disks of varying radius and positions.

- **Intuition**: Condition  $X = 0$  is not stable under perturbations of  $\omega$ .


Domains for which  $X \neq 0$ 

Figure: Rectangle of base  $2\pi$  and height  $\pi$  with bumps. The bumps are half-disks of varying radius and positions.

- **Intuition**: Condition  $X = 0$  is not stable under perturbations of  $\omega$ .
- Theoretical proof of this when  $\omega$  is a rectangle via shape derivatives.

## Conclusion and perspectives

- Structure of the essential spectrum of the Maxwell operator in a waveguide,

## Conclusion and perspectives

- Structure of the essential spectrum of the Maxwell operator in a waveguide,
- Geometric condition to have discrete spectrum: interplay between the geometry of the *curve* and the one of the cross-section  $\omega$ .

## Conclusion and perspectives

- Structure of the essential spectrum of the Maxwell operator in a waveguide,
- Geometric condition to have discrete spectrum: interplay between the geometry of the *curve* and the one of the cross-section  $\omega$ .
- To do : extension to any *twist* (Poincaré inequality, discrete spectrum), asymptotic analysis of thin-waveguides...

## Conclusion and perspectives

- Structure of the essential spectrum of the Maxwell operator in a waveguide,
- Geometric condition to have discrete spectrum: interplay between the geometry of the *curve* and the one of the cross-section  $\omega$ .
- To do : extension to any *twist* (Poincaré inequality, discrete spectrum), asymptotic analysis of thin-waveguides...

Thank you for your attention !