

Magnetic ground states and the conformal structure of a surface

Luigi Provenzano

Dipartimento di Scienze di Base e Applicate per l'Ingegneria
Sapienza Università di Roma

joint work with Bruno Colbois (Neuchâtel) and Alessandro Savo (Sapienza)



SAPIENZA
UNIVERSITÀ DI ROMA

Inverse spectral geometry

Let (M, h) be a compact Riemannian manifold and let $\{\lambda_n\}_{n=1}^{\infty}$ be the sequence of the **eigenvalues** of the Laplacian on (M, h) .

Inverse spectral geometry

Let (M, h) be a compact Riemannian manifold and let $\{\lambda_n\}_{n=1}^{\infty}$ be the sequence of the **eigenvalues** of the Laplacian on (M, h) .

What information on (M, h) is contained in $\{\lambda_n\}_{n=1}^{\infty}$? (*“can one hear the shape of a drum”?*)

Inverse spectral geometry

Let (M, h) be a compact Riemannian manifold and let $\{\lambda_n\}_{n=1}^{\infty}$ be the sequence of the **eigenvalues** of the Laplacian on (M, h) .

What information on (M, h) is contained in $\{\lambda_n\}_{n=1}^{\infty}$? (*“can one hear the shape of a drum”?*)

- $2d$ and $3d$ **isospectral** flat tori are isometric (Shienmann).

Inverse spectral geometry

Let (M, h) be a compact Riemannian manifold and let $\{\lambda_n\}_{n=1}^{\infty}$ be the sequence of the **eigenvalues** of the Laplacian on (M, h) .

What information on (M, h) is contained in $\{\lambda_n\}_{n=1}^{\infty}$? (*“can one hear the shape of a drum”?*)

- $2d$ and $3d$ **isospectral** flat tori are isometric (Shienmann).
- Examples of $\geq 4d$ isospectral non isometric flat tori (Milnor).

Inverse spectral geometry

Let (M, h) be a compact Riemannian manifold and let $\{\lambda_n\}_{n=1}^{\infty}$ be the sequence of the **eigenvalues** of the Laplacian on (M, h) .

What information on (M, h) is contained in $\{\lambda_n\}_{n=1}^{\infty}$? (*“can one hear the shape of a drum”*?)

- $2d$ and $3d$ **isospectral** flat tori are isometric (Shienmann).
- Examples of $\geq 4d$ isospectral non isometric flat tori (Milnor).
- Examples of isospectral non isometric hyperbolic surfaces of genus $g = 5$ or $g \geq 7$ (Buser).

Inverse spectral geometry

Let (M, h) be a compact Riemannian manifold and let $\{\lambda_n\}_{n=1}^{\infty}$ be the sequence of the **eigenvalues** of the Laplacian on (M, h) .

What information on (M, h) is contained in $\{\lambda_n\}_{n=1}^{\infty}$? (*“can one hear the shape of a drum”*?)

- $2d$ and $3d$ **isospectral** flat tori are isometric (Shienmann).
- Examples of $\geq 4d$ isospectral non isometric flat tori (Milnor).
- Examples of isospectral non isometric hyperbolic surfaces of genus $g = 5$ or $g \geq 7$ (Buser).
- Examples of isospectral, non-isometric planar domains (Gordon, Webb, Wolpert)
- ...

Inverse spectral geometry



You can't hear the shape of a drum!

Inverse spectral geometry



You can't hear the shape of a drum!

However the Laplacian spectrum tells us: the volume, the boundary measure, the dimension, the Euler characteristic for surfaces, etc.

The magnetic Laplacian

Let

- Σ_g be the compact, orientable surface of genus $g \geq 1$.
- A be a real **closed** 1-form on Σ_g (magnetic potential).
- h be a Riemannian metric on Σ_g .

The magnetic Laplacian

Let

- Σ_g be the compact, orientable surface of genus $g \geq 1$.
- A be a real **closed** 1-form on Σ_g (magnetic potential).
- h be a Riemannian metric on Σ_g .

For a smooth complex-valued u we define the **magnetic differential**:

$$d^A u = du - iuA$$

We set δ^A its **adjoint** .

The magnetic Laplacian

Let

- Σ_g be the compact, orientable surface of genus $g \geq 1$.
- A be a real **closed** 1-form on Σ_g (magnetic potential).
- h be a Riemannian metric on Σ_g .

For a smooth complex-valued u we define the **magnetic differential**:

$$d^A u = du - iuA$$

We set δ^A its **adjoint**. Then the **magnetic Laplacian** is defined as

$$\Delta_A u = \delta^A d^A u.$$

The magnetic Laplacian

Let

- Σ_g be the compact, orientable surface of genus $g \geq 1$.
- A be a real **closed** 1-form on Σ_g (magnetic potential).
- h be a Riemannian metric on Σ_g .

For a smooth complex-valued u we define the **magnetic differential**:

$$d^A u = du - iuA$$

We set δ^A its **adjoint**. Then the **magnetic Laplacian** is defined as

$$\Delta_A u = \delta^A d^A u.$$

Remark: if $A = 0$, $d^A = d$ and $\delta^A = \delta$, hence $\Delta_A = \Delta$ is the Laplacian (in \mathbb{R}^2 , $\delta = -\text{div}$ and hence $\Delta = -\partial_{xx}^2 - \partial_{yy}^2$).

The magnetic Laplacian: spectrum and gauge invariance

Fix g (hence Σ_g). Given a metric h and a closed 1-form A , we have that Δ_A has **discrete spectrum** $\{\lambda_n(h, A)\}_{n=1}^\infty$: $\Delta_A u_n = \lambda_n(h, A) u_n$.

The magnetic Laplacian: spectrum and gauge invariance

Fix g (hence Σ_g). Given a metric h and a closed 1-form A , we have that Δ_A has **discrete spectrum** $\{\lambda_n(h, A)\}_{n=1}^\infty$: $\Delta_A u_n = \lambda_n(h, A) u_n$.

We will be interested in the **ground state energy**:

$$\lambda_1(h, A).$$

The magnetic Laplacian: spectrum and gauge invariance

Fix g (hence Σ_g). Given a metric h and a closed 1-form A , we have that Δ_A has **discrete spectrum** $\{\lambda_n(h, A)\}_{n=1}^\infty$: $\Delta_A u_n = \lambda_n(h, A) u_n$.

We will be interested in the **ground state energy**:

$$\lambda_1(h, A).$$

The fundamental feature is the **gauge invariance**:

- If A, A' are 1-forms such that $A - A'$ is **exact**, then

$$\lambda_n(h, A) = \lambda_n(h, A'), \quad \forall n.$$

The magnetic Laplacian: spectrum and gauge invariance

Fix g (hence Σ_g). Given a metric h and a closed 1-form A , we have that Δ_A has **discrete spectrum** $\{\lambda_n(h, A)\}_{n=1}^\infty$: $\Delta_A u_n = \lambda_n(h, A) u_n$.

We will be interested in the **ground state energy**:

$$\lambda_1(h, A).$$

The fundamental feature is the **gauge invariance**:

- If A, A' are 1-forms such that $A - A'$ is **exact**, then

$$\lambda_n(h, A) = \lambda_n(h, A'), \quad \forall n.$$

- The same is true if $A - A'$ is **closed** and $\frac{1}{2\pi} \oint_\gamma (A - A') \in \mathbb{Z}$ for **any** closed curve γ .

The magnetic Laplacian: spectrum and gauge invariance

Fix g (hence Σ_g). Given a metric h and a closed 1-form A , we have that Δ_A has **discrete spectrum** $\{\lambda_n(h, A)\}_{n=1}^\infty$: $\Delta_A u_n = \lambda_n(h, A) u_n$.

We will be interested in the **ground state energy**:

$$\lambda_1(h, A).$$

The fundamental feature is the **gauge invariance**:

- If A, A' are 1-forms such that $A - A'$ is **exact**, then

$$\lambda_n(h, A) = \lambda_n(h, A'), \quad \forall n.$$

- The same is true if $A - A'$ is **closed** and $\frac{1}{2\pi} \oint_\gamma (A - A') \in \mathbb{Z}$ for **any** closed curve γ .

Therefore we can **confine** to **harmonic** 1-forms (for the metric h).

A first inequality

Let $Har(h)$ be the space of h -harmonic 1-forms on Σ_g : Euclidean space of dimension $2g$ with the L^2 -norm $\| \cdot \|_h$ on forms.

Note: $Har(h)$ does not depend on $[h]$.

A first inequality

Let $Har(h)$ be the space of h -harmonic 1-forms on Σ_g : Euclidean space of dimension $2g$ with the L^2 -norm $\|\cdot\|_h$ on forms.

Note: $Har(h)$ does not depend on $[h]$.

Let

$$L^* = \left\{ A \in Har(h) : \frac{1}{2\pi} \oint_{\gamma} A \in \mathbb{Z}, \forall \gamma \text{ closed curve} \right\}$$

be the **lattice** of harmonic 1-forms with **integral flux**.

A first inequality

Let $Har(h)$ be the space of h -harmonic 1-forms on Σ_g : Euclidean space of dimension $2g$ with the L^2 -norm $\|\cdot\|_h$ on forms.

Note: $Har(h)$ does not depend on $[h]$.

Let

$$L^* = \left\{ A \in Har(h) : \frac{1}{2\pi} \oint_{\gamma} A \in \mathbb{Z}, \forall \gamma \text{ closed curve} \right\}$$

be the **lattice** of harmonic 1-forms with **integral flux**. For $A \in Har(h)$ we define

$$d_h(A, L^*) := \min\{\|\omega - A\|_h : \omega \in L^*\}.$$

This is the L^2 -distance of A from the lattice L^* .

A first inequality

Theorem 1

$$|h|\lambda_1(h, A) \leq d_h(A, L^*)^2$$

where $|h|$ denotes the volume of Σ_g with the metric h .

Remark: $|h|\lambda_1(h, A)$ is the **normalized** eigenvalue, scale invariant.

Some conformal spectral invariants

Let h be a metric on Σ_g and A be a h -harmonic 1-form. We define

$$\Lambda_1([h], A) := \sup_{h' \in [h]} |h'| \lambda_1(h', A)$$

Some conformal spectral invariants

Let h be a metric on Σ_g and A be a h -harmonic 1-form. We define

$$\Lambda_1([h], A) := \sup_{h' \in [h]} |h'| \lambda_1(h', A)$$

We define next

$$\Lambda_1([h]) := \sup_{A \in \text{Har}(h)} \Lambda_1([h], A).$$

Some conformal spectral invariants

Let h be a metric on Σ_g and A be a h -harmonic 1-form. We define

$$\Lambda_1([h], A) := \sup_{h' \in [h]} |h'| \lambda_1(h', A)$$

We define next

$$\Lambda_1([h]) := \sup_{A \in \text{Har}(h)} \Lambda_1([h], A).$$

Finally, we define

$$\Lambda_1 = \inf_{[h]} \Lambda_1([h]).$$

This last invariant depends only on the genus g of the surface.

Conformal invariants and the Jacobian torus

All these invariants can be estimated by the geometry of the **Jacobian torus** of (Σ_g, h) :

$$Jac(h) = Har(h)/L^*.$$

Conformal invariants and the Jacobian torus

All these invariants can be estimated by the geometry of the **Jacobian torus** of (Σ_g, h) :

$$Jac(h) = Har(h)/L^*.$$

- It is a $2g$ -dimensional flat torus.
- It is **conformally invariant**: two conformal metrics h, h' on Σ_g have the same Jacobian torus.

Conformal invariants and the Jacobian torus

We define now the **inradius** of the lattice L^\star :

$$\mathcal{R}_h(L^\star) := \max_{A \in \text{Har}(h)} d_h(A, L^\star).$$

Conformal invariants and the Jacobian torus

We define now the **inradius** of the lattice L^\star :

$$\mathcal{R}_h(L^\star) := \max_{A \in \text{Har}(h)} d_h(A, L^\star).$$

It turns out that

$$\mathcal{R}_h(L^\star) = \text{diam}(\text{Jac}(h)).$$

Conformal invariants and the Jacobian torus

We define now the **inradius** of the lattice L^* :

$$\mathcal{R}_h(L^*) := \max_{A \in \text{Har}(h)} d_h(A, L^*).$$

It turns out that

$$\mathcal{R}_h(L^*) = \text{diam}(\text{Jac}(h)).$$

Combining Theorem 1 and the above definitions:

Theorem 2

- $\Lambda_1([h], A) \leq d_h(A, L^*)^2.$

Conformal invariants and the Jacobian torus

We define now the **inradius** of the lattice L^\star :

$$\mathcal{R}_h(L^\star) := \max_{A \in \text{Har}(h)} d_h(A, L^\star).$$

It turns out that

$$\mathcal{R}_h(L^\star) = \text{diam}(\text{Jac}(h)).$$

Combining Theorem 1 and the above definitions:

Theorem 2

- $\Lambda_1([h], A) \leq d_h(A, L^\star)^2.$
- $\Lambda_1([h]) \leq \mathcal{R}_h(L^\star)^2 = \text{diam}(\text{Jac}(h))^2.$

Conformal invariants and the Jacobian torus

We define now the **inradius** of the lattice L^* :

$$\mathcal{R}_h(L^*) := \max_{A \in \text{Har}(h)} d_h(A, L^*).$$

It turns out that

$$\mathcal{R}_h(L^*) = \text{diam}(\text{Jac}(h)).$$

Combining Theorem 1 and the above definitions:

Theorem 2

- $\Lambda_1([h], A) \leq d_h(A, L^*)^2.$
- $\Lambda_1([h]) \leq \mathcal{R}_h(L^*)^2 = \text{diam}(\text{Jac}(h))^2.$
- $\Lambda_1 \leq \inf_{[h]} \text{diam}(\text{Jac}(h))^2.$

Computations in genus $g = 1$

We consider now Σ_1 : the torus.

Computations in genus $g = 1$

We consider now Σ_1 : the torus.


By the uniformization theorem we have that any metric h on Σ_1 is conformal to a **unique flat metric** \hat{h} .

Computations in genus $g = 1$

We consider now Σ_1 : the torus.

By the uniformization theorem we have that any metric h on Σ_1 is conformal to a **unique flat metric** \hat{h} .

A flat torus is the quotient of \mathbb{R}^2/L by a lattice $L = \ell w_1 + m w_2$, $\ell, m \in \mathbb{Z}$, with the inherited Euclidean metric.



Computations in genus $g = 1$

Theorem 3

Given a metric h on Σ_1 we have

$$|h|\lambda_1(h, A) \leq |\hat{h}|\lambda_1(\hat{h}, A),$$

where \hat{h} is the unique flat metric in $[h]$. Equality holds iff $h = \hat{h}$.

Computations in genus $g = 1$

Theorem 3

Given a metric h on Σ_1 we have

$$|h|\lambda_1(h, A) \leq |\hat{h}|\lambda_1(\hat{h}, A),$$

*where \hat{h} is the unique flat metric in $[h]$. Equality holds iff $h = \hat{h}$.
In particular*

$$\Lambda_1([h], A) = |\hat{h}|\lambda_1(\hat{h}, A).$$

So we can compute the conformal invariants if we understand the spectrum of **flat tori**.

Computations in genus $g = 1$

Consider then the flat torus \mathbb{R}^2/L with basis (w_1, w_2) .

- The dual lattice L^* of L is

$$L^* = \{w^* \in \mathbb{R}^2 : \langle w^*, w \rangle \in 2\pi\mathbb{Z}, \quad \forall w \in L\}$$

Computations in genus $g = 1$

Consider then the flat torus \mathbb{R}^2/L with basis (w_1, w_2) .

- The **dual lattice** L^* of L is

$$L^* = \{w^* \in \mathbb{R}^2 : \langle w^*, w \rangle \in 2\pi\mathbb{Z}, \quad \forall w \in L\}$$

- It is easy to see that L^* is **homothetic** to L : $L^* \sim \frac{2\pi L}{|\hat{h}|}$

Computations in genus $g = 1$

Consider then the flat torus \mathbb{R}^2/L with basis (w_1, w_2) .

- The **dual lattice** L^* of L is

$$L^* = \{w^* \in \mathbb{R}^2 : \langle w^*, w \rangle \in 2\pi\mathbb{Z}, \quad \forall w \in L\}$$

- It is easy to see that L^* is **homothetic** to L : $L^* \sim \frac{2\pi L}{|\hat{h}|}$
- A harmonic 1-form A on \mathbb{R}^2/L descends from a parallel form on \mathbb{R}^2 :

$$A = a dx + b dy \quad \sim \quad P_A = (a, b) \in \mathbb{R}^2.$$

Computations in genus $g = 1$

Consider then the flat torus \mathbb{R}^2/L with basis (w_1, w_2) .

- The **dual lattice** L^* of L is

$$L^* = \{w^* \in \mathbb{R}^2 : \langle w^*, w \rangle \in 2\pi\mathbb{Z}, \quad \forall w \in L\}$$

- It is easy to see that L^* is **homothetic** to L : $L^* \sim \frac{2\pi L}{|\hat{h}|}$
- A harmonic 1-form A on \mathbb{R}^2/L descends from a parallel form on \mathbb{R}^2 :

$$A = a dx + b dy \quad \sim \quad P_A = (a, b) \in \mathbb{R}^2.$$

- $\{e^{i\langle \cdot, w^* \rangle} : w^* \in L^*\}$ are all the **eigenfunctions** of Δ_A .

Computations in genus $g = 1$

Consider then the flat torus \mathbb{R}^2/L with basis (w_1, w_2) .

- The **dual lattice** L^* of L is

$$L^* = \{w^* \in \mathbb{R}^2 : \langle w^*, w \rangle \in 2\pi\mathbb{Z}, \quad \forall w \in L\}$$

- It is easy to see that L^* is **homothetic** to L : $L^* \sim \frac{2\pi L}{|\hat{h}|}$
- A harmonic 1-form A on \mathbb{R}^2/L descends from a parallel form on \mathbb{R}^2 :

$$A = a dx + b dy \quad \sim \quad P_A = (a, b) \in \mathbb{R}^2.$$

- $\{e^{i\langle \cdot, w^* \rangle} : w^* \in L^*\}$ are all the **eigenfunctions** of Δ_A .
- The eigenvalue corresponding to $e^{i\langle \cdot, w^* \rangle}$ is $|P_A - w^*|^2$

Computations in genus $g = 1$

Consider then the flat torus \mathbb{R}^2/L with basis (w_1, w_2) .

- The **dual lattice** L^* of L is

$$L^* = \{w^* \in \mathbb{R}^2 : \langle w^*, w \rangle \in 2\pi\mathbb{Z}, \quad \forall w \in L\}$$


- It is easy to see that L^* is **homothetic** to L : $L^* \sim \frac{2\pi L}{|\hat{h}|}$
- A harmonic 1-form A on \mathbb{R}^2/L descends from a parallel form on \mathbb{R}^2 :

$$A = a dx + b dy \quad \sim \quad P_A = (a, b) \in \mathbb{R}^2.$$


- $\{e^{i\langle \cdot, w^* \rangle} : w^* \in L^*\}$ are all the **eigenfunctions** of Δ_A .
- The eigenvalue corresponding to $e^{i\langle \cdot, w^* \rangle}$ is $|P_A - w^*|^2$
- The first eigenvalue is

$$\lambda_1(\hat{h}, A) = \min_{w^* \in L^*} |P_A - w^*|^2.$$

Computations in genus $g = 1$



Computations in genus $g = 1$



Remark: The dual lattice L^* is identified with the lattice of harmonic 1-forms with **integral fluxes**: $\mathbb{R}^2/L^* = \text{Jac}(\hat{h})$ with a homotetic metric of factor $|\hat{h}|$.

Computations in genus $g = 1$

We go back to our **spectral invariants**:

$$\Lambda_1([h], A) = \sup_{h' \in [h]} |h'| \lambda_1(h', A)$$

- We have seen that

$$\Lambda_1([h], A) = |\hat{h}| \lambda_1(\hat{h}, A) = d_h(A, L^*)^2.$$

Computations in genus $g = 1$

We go back to our **spectral invariants**:

$$\Lambda_1([h], A) = \sup_{h' \in [h]} |h'| \lambda_1(h', A)$$

- We have seen that

$$\Lambda_1([h], A) = |\hat{h}| \lambda_1(\hat{h}, A) = d_h(A, L^*)^2.$$

- **Conclusion**: the first invariant “hears” the distance of a harmonic 1-form from the lattice of 1-forms with integral flux in a given conformal class $[h]$.

Computations in genus $g = 1$

The second invariant is:

$$\Lambda_1([h]) = \sup_{A \in \text{Har}(h)} \Lambda_1([h], A).$$

Computations in genus $g = 1$

The second invariant is:

$$\Lambda_1([h]) = \sup_{A \in \text{Har}(h)} \Lambda_1([h], A).$$

- The best potential corresponds to P_A at the **maximum distance** from L^* : we get the **inradius** $\mathcal{R}_h(L^*)$ of L^* :

$$\Lambda_1([h]) = \mathcal{R}_h(L^*)^2 = \text{diam}(\text{Jac}(h))^2$$

Computations in genus $g = 1$

The second invariant is:


$$\Lambda_1([h]) = \sup_{A \in \text{Har}(h)} \Lambda_1([h], A).$$

- The best potential corresponds to P_A at the **maximum distance** from L^* : we get the **inradius** $\mathcal{R}_h(L^*)$ of L^* :

$$\Lambda_1([h]) = \mathcal{R}_h(L^*)^2 = \text{diam}(\text{Jac}(h))^2$$

- **Conclusion:** the second invariant “hears” the diameter of the Jacobian torus of $[h]$.

Computations in genus $g = 1$



Computations in genus $g = 1$

The third invariant is:

$$\Lambda_1 = \inf_{[h]} \Lambda_1([h]).$$


- This problem is equivalent to finding the lattice of given area with the **smallest inradius**.

Computations in genus $g = 1$

The third invariant is:

$$\Lambda_1 = \inf_{[h]} \Lambda_1([h]).$$

- This problem is equivalent to finding the lattice of given area with the **smallest inradius**. This is the **equilateral torus**.



- Λ_1 identifies the conformal class of the equilateral flat torus.

Some questions in $g \geq 2$

In $g \geq 2$ explicit computations are not possible.

- What is the optimal metric for $\Lambda_1([h], A)$? In general it is not the hyperbolic metric.

Some questions in $g \geq 2$

In $g \geq 2$ explicit computations are not possible.

- What is the optimal metric for $\Lambda_1([h], A)$? In general it is not the hyperbolic metric.
- Analogous questions for $\Lambda_1([h])$.

Some questions in $g \geq 2$

In $g \geq 2$ explicit computations are not possible.

- What is the optimal metric for $\Lambda_1([h], A)$? In general it is not the hyperbolic metric.
- Analogous questions for $\Lambda_1([h])$.
- What is the optimal conformal class in the definition of Λ_1 ? Is Λ_1 always positive?

Some questions in $g \geq 2$

In $g \geq 2$ explicit computations are not possible.

- What is the optimal metric for $\Lambda_1([h], A)$? In general it is not the hyperbolic metric.
- Analogous questions for $\Lambda_1([h])$.
- What is the optimal conformal class in the definition of Λ_1 ? Is Λ_1 always positive?
- Do the three invariants “hear” the same information on the Jacobian torus as in $g = 1$?

Can we hear the conformal class of a surface?

We prove the following:

Theorem 4

Let ω be a closed 1-form on the surface Σ_g . Let h be a Riemannian metric on Σ_g .

Can we hear the conformal class of a surface?

We prove the following:

Theorem 4

Let ω be a closed 1-form on the surface Σ_g . Let h be a Riemannian metric on Σ_g . Then

$$\lim_{t \rightarrow 0} \frac{\lambda_1(h, t\omega)}{t^2}$$

Can we hear the conformal class of a surface?

We prove the following:

Theorem 4

Let ω be a closed 1-form on the surface Σ_g . Let h be a Riemannian metric on Σ_g . Then

$$\lim_{t \rightarrow 0} \frac{\lambda_1(h, t\omega)}{t^2} = \frac{\|A\|_h^2}{|h|}$$

where A is the unique harmonic form for h co-homologous to ω .

Can we hear the conformal class of a surface?

Our main result is the following

Theorem 5

Assume that the metrics h_1, h_2 on Σ_g are such that

$$\lambda_1(h_1, \omega) = \lambda_1(h_2, \omega)$$

for all closed 1-forms ω on Σ_g .

Can we hear the conformal class of a surface?

Our main result is the following

Theorem 5

Assume that the metrics h_1, h_2 on Σ_g are such that

$$\lambda_1(h_1, \omega) = \lambda_1(h_2, \omega)$$

for all closed 1-forms ω on Σ_g . Then they are conformal and have the same volume.

Can we hear the conformal class of a surface?

Our main result is the following

Theorem 5

Assume that the metrics h_1, h_2 on Σ_g are such that

$$\lambda_1(h_1, \omega) = \lambda_1(h_2, \omega)$$

for all closed 1-forms ω on Σ_g . Then they are conformal and have the same volume.

We deduce the following corollary

Corollary

*Assume that the metrics in Theorem 5 are **hyperbolic**. Then they are isometric. For $g = 1$, the statement holds for flat metrics.*


Can we hear the conformal class of a surface?

It seems that this notion of “isospectrality” requires the knowledge of **uncountable** information: $\lambda_1(h, \omega)$ for all closed 1-forms ω . This isn't quite the case.

Can we hear the conformal class of a surface?

It seems that this notion of “isospectrality” requires the knowledge of **uncountable** information: $\lambda_1(h, \omega)$ for all closed 1-forms ω . This isn't quite the case.


Fix a **canonical homology basis** $(\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g)$ and let be $(\omega_1, \dots, \omega_{2g})$ a **dual basis** of closed 1-forms.



Can we hear the conformal class of a surface?

It seems that this notion of “isospectrality” requires the knowledge of **uncountable** information: $\lambda_1(h, \omega)$ for all closed 1-forms ω . This isn't quite the case.

Fix a **canonical homology basis** $(\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g)$ and let be $(\omega_1, \dots, \omega_{2g})$ a **dual basis** of closed 1-forms.




$$\text{Let } \omega_{i,j}^n = \frac{\omega_i + \omega_j}{n}.$$

Can we hear the conformal class of a surface?

It seems that this notion of “isospectrality” requires the knowledge of **uncountable** information: $\lambda_1(h, \omega)$ for all closed 1-forms ω . This isn't quite the case.

Fix a **canonical homology basis** $(\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g)$ and let be $(\omega_1, \dots, \omega_{2g})$ a **dual basis** of closed 1-forms.



Let $\omega_{i,j}^n = \frac{\omega_i + \omega_j}{n}$. If two metrics h_1, h_2 on Σ_g satisfy

$$\lambda_1(h_1, \omega_{i,j}^n) = \lambda_1(h_2, \omega_{i,j}^n), \quad \forall n \geq 1, 1 \leq i \leq j \leq 2g,$$

then they are conformal and have the same volume.

A few words on the proof

- **Starting point:** given a closed 1-form ω , $\lim_{t \rightarrow 0} \frac{\lambda_1(h, t\omega)}{t^2} = \frac{\|A\|_h^2}{|h|}$, where A is h -harmonic co-homologous to ω :

“we hear all the L^2 -norms of h -harmonic 1-forms”.

A few words on the proof

- **Starting point:** given a closed 1-form ω , $\lim_{t \rightarrow 0} \frac{\lambda_1(h, t\omega)}{t^2} = \frac{\|A\|_h^2}{|h|}$, where A is h -harmonic co-homologous to ω :
“we hear all the L^2 -norms of h -harmonic 1-forms”.
- We can reconstruct the **Gram matrix** of the Jacobian torus.

A few words on the proof

- **Starting point:** given a closed 1-form ω , $\lim_{t \rightarrow 0} \frac{\lambda_1(h, t\omega)}{t^2} = \frac{\|A\|_h^2}{|h|}$, where A is h -harmonic co-homologous to ω :
“we hear all the L^2 -norms of h -harmonic 1-forms”.
- We can reconstruct the **Gram matrix** of the Jacobian torus.
- The homology basis is fixed and is the canonical one, hence the Gram matrix has determinant 1 (Riemann period relations)

A few words on the proof

- **Starting point:** given a closed 1-form ω , $\lim_{t \rightarrow 0} \frac{\lambda_1(h, t\omega)}{t^2} = \frac{\|A\|_h^2}{|h|}$, where A is h -harmonic co-homologous to ω :

“we hear all the L^2 -norms of h -harmonic 1-forms”.

- We can reconstruct the **Gram matrix** of the Jacobian torus.
- The homology basis is fixed and is the canonical one, hence the Gram matrix has determinant 1 (Riemann period relations)
- If two metrics are “ground state isospectral”, they have the same Gram matrix and the same volume by the three points above.

A few words on the proof

- **Starting point:** given a closed 1-form ω , $\lim_{t \rightarrow 0} \frac{\lambda_1(h, t\omega)}{t^2} = \frac{\|A\|_h^2}{|h|}$, where A is h -harmonic co-homologous to ω :

“we hear all the L^2 -norms of h -harmonic 1-forms”.

- We can reconstruct the **Gram matrix** of the Jacobian torus.
- The homology basis is fixed and is the canonical one, hence the Gram matrix has determinant 1 (Riemann period relations)
- If two metrics are “ground state isospectral”, they have the same Gram matrix and the same volume by the three points above.
- We prove that if h_1, h_2 have the same Gram matrix, they are conformal. Here we use a deep result in Riemann surfaces theory, the **Torelli's Theorem**.

Thank you for your attention



B. Colbois, L. Provenzano and A. Savo,
Magnetic ground states and the conformal class of a surface,
preprint (2025).