

# Eigenvalues of operator functions: A commutativity result with an application to 1D Dirac operators

**Nicolas Weber**

Graz University of Technology  
Joint work with J. Behrndt and P. Siegl

**Analytic and algebraic methods in physics XXII**  
**Prague, August 26-29, 2025**

# Content

1. Motivation
2. Eigenvalues of operator functions
3. Outlines of the proofs

# Motivation: Spectrum of perturbed 1D Dirac

Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

# Motivation: Spectrum of perturbed 1D Dirac

Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

Perturbed Dirac:  $H_0 + V$  with matrix potential

$$V : \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}, \quad \|V(\cdot)\| \in L^1(\mathbb{R}) \cap L^2(\mathbb{R}).$$

# Motivation: Spectrum of perturbed 1D Dirac

Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

Perturbed Dirac:  $H_0 + V$  with matrix potential

$$V : \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}, \quad \|V(\cdot)\| \in L^1(\mathbb{R}) \cap L^2(\mathbb{R}).$$

$$\rightsquigarrow \sigma(H_0) = \sigma_{\text{ess}}(H_0) = (-\infty, -m] \cup [m, \infty).$$

# Motivation: Spectrum of perturbed 1D Dirac


Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

Perturbed Dirac:  $H_0 + V$  with matrix potential

$$V : \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}, \quad \|V(\cdot)\| \in L^1(\mathbb{R}) \cap L^2(\mathbb{R}).$$

$$\rightsquigarrow \sigma(H_0) = \sigma_{\text{ess}}(H_0) = (-\infty, -m] \cup [m, \infty).$$



# Motivation: Spectrum of perturbed 1D Dirac


Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

Perturbed Dirac:  $H_0 + V$  with matrix potential

$$V : \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}, \quad \|V(\cdot)\| \in L^1(\mathbb{R}) \cap L^2(\mathbb{R}).$$

$$\rightsquigarrow \sigma_{\text{ess}}(H_0 + V) = \sigma_{\text{ess}}(H_0) = (-\infty, -m] \cup [m, \infty).$$



# Motivation: Spectrum of perturbed 1D Dirac


Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

Perturbed Dirac:  $H_0 + V$  with matrix potential

$$V : \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}, \quad \|V(\cdot)\| \in L^1(\mathbb{R}) \cap L^2(\mathbb{R}).$$

Non-embedded eigenvalues: [Cuenin-Laptev-Tretter '14]





# Motivation: Spectrum of perturbed 1D Dirac


Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

Perturbed Dirac:  $H_0 + V$  with matrix potential

$$V : \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}, \quad \|V(\cdot)\| \in L^1(\mathbb{R}) \cap L^2(\mathbb{R}).$$

Non-embedded eigenvalues: [Cuenin-Laptev-Tretter '14]



# Motivation: Spectrum of perturbed 1D Dirac


Consider for  $m \geq 0$  in  $L^2(\mathbb{R}; \mathbb{C}^2)$  the free Dirac operator

$$H_0 = \begin{pmatrix} m & -i\partial_x \\ -i\partial_x & -m \end{pmatrix}, \quad \text{dom } H_0 = H^1(\mathbb{R}; \mathbb{C}^2).$$

Perturbed Dirac:  $H_0 + V$  with matrix potential

$$V : \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}, \quad \|V(\cdot)\| \in L^1(\mathbb{R}) \cap L^2(\mathbb{R}).$$

Non-embedded eigenvalues: [Cuenin-Laptev-Tretter '14]



Does  $H_0 + V$  have eigenvalues?  $\rightsquigarrow H_\varepsilon = H_0 + \varepsilon V$ .

# Existence of eigenvalues in weak coupling

# Existence of eigenvalues in weak coupling

Theorem (Cuenin-Siegl 18' )

# Existence of eigenvalues in weak coupling

## Theorem (Cuenin-Siegl 18' )

*Set  $U := \int_{\mathbb{R}} V(x) \, dx \in \mathbb{C}^{2 \times 2}$ . If  $m > 0$ :*

# Existence of eigenvalues in weak coupling

## Theorem (Cuenin-Siegl 18' )

*Set  $U := \int_{\mathbb{R}} V(x) dx \in \mathbb{C}^{2 \times 2}$ . If  $m > 0$ :  $\forall 0 < \varepsilon < \varepsilon_0$ :*

# Existence of eigenvalues in weak coupling

## Theorem (Cuenin-Siegl 18' )

Set  $U := \int_{\mathbb{R}} V(x) dx \in \mathbb{C}^{2 \times 2}$ . If  $m > 0$ :  $\forall 0 < \varepsilon < \varepsilon_0$ :

1.  $\Re(U_{11}) < 0 \implies \exists z_+(\varepsilon) \in \sigma_p(H_\varepsilon)$

# Existence of eigenvalues in weak coupling

## Theorem (Cuenin-Siegl 18' )

Set  $U := \int_{\mathbb{R}} V(x) dx \in \mathbb{C}^{2 \times 2}$ . If  $m > 0$ :  $\forall 0 < \varepsilon < \varepsilon_0$ :

$$1. \Re(U_{11}) < 0 \implies \exists z_+(\varepsilon) \in \sigma_p(H_\varepsilon)$$

$$z_+(\varepsilon) = m - \frac{m}{2} U_{11}^2 \varepsilon^2 + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0+.$$



# Existence of eigenvalues in weak coupling

## Theorem (Cuenin-Siegl 18' )

Set  $U := \int_{\mathbb{R}} V(x) dx \in \mathbb{C}^{2 \times 2}$ . If  $m > 0$ :  $\forall 0 < \varepsilon < \varepsilon_0$ :

1.  $\Re(U_{11}) < 0 \implies \exists z_+(\varepsilon) \in \sigma_p(H_\varepsilon)$

$$z_+(\varepsilon) = m - \frac{m}{2} U_{11}^2 \varepsilon^2 + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0+.$$

2.  $\Re(U_{22}) > 0 \implies \exists z_-(\varepsilon) \in \sigma_p(H_\varepsilon)$

# Existence of eigenvalues in weak coupling

## Theorem (Cuenin-Siegl 18' )

Set  $U := \int_{\mathbb{R}} V(x) dx \in \mathbb{C}^{2 \times 2}$ . If  $m > 0$ :  $\forall 0 < \varepsilon < \varepsilon_0$ :

1.  $\Re(U_{11}) < 0 \implies \exists z_+(\varepsilon) \in \sigma_p(H_\varepsilon)$

$$z_+(\varepsilon) = m - \frac{m}{2} U_{11}^2 \varepsilon^2 + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0+.$$

2.  $\Re(U_{22}) > 0 \implies \exists z_-(\varepsilon) \in \sigma_p(H_\varepsilon)$

$$z_-(\varepsilon) = -m + \frac{m}{2} U_{22}^2 \varepsilon^2 + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0+.$$

# Existence of eigenvalues in weak coupling

## Theorem (Cuenin-Siegl 18')


Set  $U := \int_{\mathbb{R}} V(x) dx \in \mathbb{C}^{2 \times 2}$ . If  $m > 0$ :  $\forall 0 < \varepsilon < \varepsilon_0$ :

1.  $\Re(U_{11}) < 0 \implies \exists z_+(\varepsilon) \in \sigma_p(H_\varepsilon)$

$$z_+(\varepsilon) = m - \frac{m}{2} U_{11}^2 \varepsilon^2 + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0+.$$

2.  $\Re(U_{22}) > 0 \implies \exists z_-(\varepsilon) \in \sigma_p(H_\varepsilon)$

$$z_-(\varepsilon) = -m + \frac{m}{2} U_{22}^2 \varepsilon^2 + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0+.$$



What are the algebraic multiplicities of  $z_{\pm}(\varepsilon)$ ?

What are the algebraic multiplicities of  $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

What are the algebraic multiplicities of  $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ff}}(W) \iff$$

What are the algebraic multiplicities of  $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \left\{ \begin{array}{l} \ker W(z_0) \neq \{0\} \end{array} \right.$$

What are the algebraic multiplicities of  $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, W(z_0) \text{ Fredholm} \end{cases}$$



# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} W'(z) W(z)^{-1} dz \right)$$

# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} W'(z) W(z)^{-1} dz \right)$$

Special case:  $W(z) = zI - A$

# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} W'(z) W(z)^{-1} dz \right)$$

Special case:  $W(z) = zI - A$

$z_0 \in \sigma_p(A)$  isolated

# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, \ W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} W'(z) W(z)^{-1} dz \right)$$

Special case:  $W(z) = zI - A$

$z_0 \in \sigma_p(A)$  isolated

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} (zI - A)^{-1} dz \right)$$

# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} W'(z) W(z)^{-1} dz \right)$$

Special case:  $W(z) = zI - A$

$z_0 \in \sigma_p(A)$  isolated

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} \overbrace{(zI - A)^{-1}}^{\text{Riesz-projection } P_{z_0}} dz \right)$$

# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} W'(z) W(z)^{-1} dz \right)$$

Special case:  $W(z) = zI - A$

$z_0 \in \sigma_p(A)$  isolated

$$m_a(z_0; W) := \text{tr} \left( \overbrace{\frac{1}{2\pi i} \int_{\gamma} (zI - A)^{-1} dz}^{\text{Riesz-projection } P_{z_0}} \right) = \dim \text{ran } P_{z_0}$$

# What are the algebraic multiplicities of $z_{\pm}(\varepsilon)$ ?

Algebraic multiplicities for  $W : \Omega \rightarrow \mathcal{B}(\mathcal{H})$  analytic

$$z_0 \in \sigma_{\text{ft}}(W) \iff \begin{cases} \ker W(z_0) \neq \{0\}, \quad W(z_0) \text{ Fredholm} \\ W(z)^{-1} \in \mathcal{B}(\mathcal{H}) \quad \forall z \in D_r(z_0) \setminus \{z_0\} \end{cases}$$

$$m_a(z_0; W) := \text{tr} \left( \frac{1}{2\pi i} \int_{\gamma} W'(z) W(z)^{-1} dz \right)$$

Special case:  $W(z) = zI - A$

$z_0 \in \sigma_p(A)$  isolated

$$\begin{aligned} m_a(z_0; W) &:= \text{tr} \left( \overbrace{\frac{1}{2\pi i} \int_{\gamma} (zI - A)^{-1} dz}^{\text{Riesz-projection } P_{z_0}} \right) = \dim \text{ran } P_{z_0} \\ &= \text{alg. mult. of } z_0 \in \sigma_p(A). \end{aligned}$$



# Existence of eigenvalues: Idea of the proof

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle:

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$z \in \sigma_p(H_\varepsilon) \iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} |V|^{\frac{1}{2}})$$

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$z \in \sigma_p(H_\varepsilon) \iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} V^{\frac{1}{2}}) \quad m_g \checkmark$$

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$z \in \sigma_p(H_\varepsilon) \iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} V^{\frac{1}{2}}) \quad m_g \checkmark \quad m_a \checkmark$$

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$z \in \sigma_p(H_\varepsilon) \iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} |V|^{\frac{1}{2}}) \quad m_g \checkmark \quad m_a \checkmark$$

$\varepsilon \ll 1$   
 $\iff$

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) &\iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} |V|^{\frac{1}{2}}) & m_g \checkmark \quad m_a \checkmark \\ &\stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - A_\varepsilon(z)B) \end{aligned}$$

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$z \in \sigma_p(H_\varepsilon) \iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} |V|^{\frac{1}{2}}) \quad m_g \checkmark \quad m_a \checkmark$$
$$\stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - A_\varepsilon(z)B)$$

Here:  $A_\varepsilon(z) : \mathbb{C}^2 \rightarrow L^2(\mathbb{R}; \mathbb{C}^2)$ ,  $B : L^2(\mathbb{R}; \mathbb{C}^2) \rightarrow \mathbb{C}^2$ .



# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) &\iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} |V|^{\frac{1}{2}}) && m_g \checkmark \quad m_a \checkmark \\ &\stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - A_\varepsilon(z)B) && m_g \checkmark \quad m_a \checkmark \end{aligned}$$

Here:  $A_\varepsilon(z) : \mathbb{C}^2 \rightarrow L^2(\mathbb{R}; \mathbb{C}^2)$ ,  $B : L^2(\mathbb{R}; \mathbb{C}^2) \rightarrow \mathbb{C}^2$ .

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) &\iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} |V|^{\frac{1}{2}}) && m_g \checkmark \quad m_a \checkmark \\ &\stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - A_\varepsilon(z)B) && m_g \checkmark \quad m_a \checkmark \end{aligned}$$

Here:  $A_\varepsilon(z) : \mathbb{C}^2 \rightarrow L^2(\mathbb{R}; \mathbb{C}^2)$ ,  $B : L^2(\mathbb{R}; \mathbb{C}^2) \rightarrow \mathbb{C}^2$ .

Use:  $\ker(I - TS) \simeq \ker(I - ST) \quad [T \in \mathcal{B}(\mathcal{H}, \mathcal{G}), S \in \mathcal{B}(\mathcal{G}, \mathcal{H})]$ .

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) &\iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1}|V|^{\frac{1}{2}}) && m_g \checkmark \quad m_a \checkmark \\ &\stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - A_\varepsilon(z)B) && m_g \checkmark \quad m_a \checkmark \end{aligned}$$

Here:  $A_\varepsilon(z) : \mathbb{C}^2 \rightarrow L^2(\mathbb{R}; \mathbb{C}^2)$ ,  $B : L^2(\mathbb{R}; \mathbb{C}^2) \rightarrow \mathbb{C}^2$ .

Use:  $\ker(I - TS) \simeq \ker(I - ST) \quad [T \in \mathcal{B}(\mathcal{H}, \mathcal{G}), S \in \mathcal{B}(\mathcal{G}, \mathcal{H})]$ .

$$z \in \sigma_p(H_\varepsilon) \stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - BA_\varepsilon(z)) \quad m_g \checkmark$$

# Existence of eigenvalues: Idea of the proof

Birman-Schwinger principle: If  $z \in \rho(H_0)$ , then

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) &\iff 0 \in \sigma_p(I - \varepsilon V^{\frac{1}{2}}(H_0 - z)^{-1} |V|^{\frac{1}{2}}) && m_g \checkmark \quad m_a \checkmark \\ &\stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - A_\varepsilon(z)B) && m_g \checkmark \quad m_a \checkmark \end{aligned}$$

Here:  $A_\varepsilon(z) : \mathbb{C}^2 \rightarrow L^2(\mathbb{R}; \mathbb{C}^2)$ ,  $B : L^2(\mathbb{R}; \mathbb{C}^2) \rightarrow \mathbb{C}^2$ .

Use:  $\ker(I - TS) \simeq \ker(I - ST) \quad [T \in \mathcal{B}(\mathcal{H}, \mathcal{G}), S \in \mathcal{B}(\mathcal{G}, \mathcal{H})]$ .

$$z \in \sigma_p(H_\varepsilon) \stackrel{\varepsilon \ll 1}{\iff} 0 \in \sigma_p(I - BA_\varepsilon(z)) \quad m_g \checkmark \quad m_a ?$$

**Question:**  $m_a(z; I - TS) = m_a(z; I - ST)$ ?

# Eigenvalues of operator functions

## Theorem

# Eigenvalues of operator functions

## Theorem

*Suppose*

- $\Omega \subset \mathbb{C}$  *open*.

# Eigenvalues of operator functions

## Theorem

*Suppose*

- $\Omega \subset \mathbb{C}$  *open*.
- $T : \Omega \rightarrow \mathcal{L}(\mathcal{H}, \mathcal{G})$ ,  $S : \Omega \rightarrow \mathcal{L}(\mathcal{G}, \mathcal{H})$  *analytic*.

# Eigenvalues of operator functions

## Theorem

*Suppose*

- $\Omega \subset \mathbb{C}$  open.
- $T : \Omega \rightarrow \mathcal{L}(\mathcal{H}, \mathcal{G}), S : \Omega \rightarrow \mathcal{L}(\mathcal{G}, \mathcal{H})$  analytic.

*Then:*

$$\forall z \in \Omega : \quad z \in \sigma_{\text{ff}}(I - TS) \iff z \in \sigma_{\text{ff}}(I - ST),$$



# Eigenvalues of operator functions

## Theorem

*Suppose*

- $\Omega \subset \mathbb{C}$  open.
- $T : \Omega \rightarrow \mathcal{L}(\mathcal{H}, \mathcal{G}), S : \Omega \rightarrow \mathcal{L}(\mathcal{G}, \mathcal{H})$  analytic.

*Then:*

$$\forall z \in \Omega : \quad z \in \sigma_{\text{ft}}(I - TS) \iff z \in \sigma_{\text{ft}}(I - ST),$$

*in which case*

$$m_a(z_0; I - TS) = m_a(z_0; I - ST).$$

# Eigenvalues of operator functions

## Theorem

*Suppose*

- $\Omega \subset \mathbb{C}$  open.
- $T : \Omega \rightarrow \mathcal{L}(\mathcal{H}, \mathcal{G})$ ,  $S : \Omega \rightarrow \mathcal{L}(\mathcal{G}, \mathcal{H})$  analytic.

*Then:*

$$\forall z \in \Omega : \quad z \in \sigma_{\text{ft}}(I - TS) \iff z \in \sigma_{\text{ft}}(I - ST),$$

*in which case*

$$m_a(z_0; I - TS) = m_a(z_0; I - ST).$$

Applied to our situation:

$$m_a(z_0; I - H_\epsilon) = m_a(z_0; I - A_\epsilon B) = m_a(z_0; I - BA_\epsilon).$$

# Application to our problem

To summarize:

# Application to our problem

To summarize:

$$z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) \iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}})$$

# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0 \end{aligned}$$

# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0. \end{aligned}$$

Special structure of Dirac:

$$\det F(z) = 1 + \varepsilon \operatorname{tr}(U\Upsilon(z)) + o(1), \quad U, \Upsilon(z) \in \mathbb{C}^{2 \times 2}$$

# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0. \end{aligned}$$

Special structure of Dirac:

$$\det F(z) = 1 + \varepsilon \operatorname{tr}(U\Upsilon(z)) + o(1), \quad U, \Upsilon(z) \in \mathbb{C}^{2 \times 2}$$

$\xRightarrow{\text{Rouche}} \exists$  simple root  $z_0$  of  $\det F$ .

# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0. \end{aligned}$$

Special structure of Dirac:

$$\det F(z) = 1 + \varepsilon \operatorname{tr}(U\Upsilon(z)) + o(1), \quad U, \Upsilon(z) \in \mathbb{C}^{2 \times 2}$$

$\xRightarrow{\text{Rouche}} \exists$  simple root  $z_0$  of  $\det F$ . Moreover:

$$m_a(z_0; zI - H_\varepsilon)$$



# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0. \end{aligned}$$

Special structure of Dirac:

$$\det F(z) = 1 + \varepsilon \operatorname{tr}(U\Upsilon(z)) + o(1), \quad U, \Upsilon(z) \in \mathbb{C}^{2 \times 2}$$

$\xRightarrow{\text{Rouche}} \exists$  simple root  $z_0$  of  $\det F$ . Moreover:

$$m_a(z_0; zI - H_\varepsilon) = m_a(z_0; F)$$

# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0. \end{aligned}$$

Special structure of Dirac:

$$\det F(z) = 1 + \varepsilon \operatorname{tr}(U\Upsilon(z)) + o(1), \quad U, \Upsilon(z) \in \mathbb{C}^{2 \times 2}$$

$\xRightarrow{\text{Rouche}} \exists$  simple root  $z_0$  of  $\det F$ . Moreover:

$$\begin{aligned} m_a(z_0; zI - H_\varepsilon) &= m_a(z_0; F) \\ &= \frac{1}{2\pi i} \int_\gamma \operatorname{tr}\{F'F^{-1}\} dz \end{aligned}$$

# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0. \end{aligned}$$

Special structure of Dirac:

$$\det F(z) = 1 + \varepsilon \operatorname{tr}(U\Upsilon(z)) + o(1), \quad U, \Upsilon(z) \in \mathbb{C}^{2 \times 2}$$

$\xRightarrow{\text{Rouche}} \exists$  simple root  $z_0$  of  $\det F$ . Moreover:

$$\begin{aligned} m_a(z_0; zI - H_\varepsilon) &= m_a(z_0; F) \\ &= \frac{1}{2\pi i} \int_\gamma \operatorname{tr}\{F'F^{-1}\} dz \\ &= \frac{1}{2\pi i} \int_\gamma \frac{(\det F)'}{\det F} dz \end{aligned}$$

# Application to our problem

To summarize:

$$\begin{aligned} z \in \sigma_p(H_\varepsilon) \cap \rho(H_0) &\iff 0 \in \sigma_p(\overbrace{I - BA_\varepsilon(z)}^{=: F(z) \in \mathbb{C}^{2 \times 2}}) \\ &\iff \det F(z) = 0. \end{aligned}$$

Special structure of Dirac:

$$\det F(z) = 1 + \varepsilon \operatorname{tr}(U\Upsilon(z)) + o(1), \quad U, \Upsilon(z) \in \mathbb{C}^{2 \times 2}$$

$\xRightarrow{\text{Rouche}} \exists$  simple root  $z_0$  of  $\det F$ . Moreover:

$$\begin{aligned} m_a(z_0; zI - H_\varepsilon) &= m_a(z_0; F) \\ &= \frac{1}{2\pi i} \int_\gamma \operatorname{tr}\{F'F^{-1}\} dz \\ &= \frac{1}{2\pi i} \int_\gamma \frac{(\det F)'}{\det F} dz \\ &= \text{multiplicity of } z_0 \text{ as a root of } \det F. \end{aligned}$$

# Outlines of the proofs

## Theorem

*Suppose*

- $\Omega \subset \mathbb{C}$  *open*.
- $T : \Omega \rightarrow \mathcal{L}(\mathcal{H}, \mathcal{G}), S : \Omega \rightarrow \mathcal{L}(\mathcal{G}, \mathcal{H})$  *analytic*.

*Then:*

$$\forall z \in \Omega : \quad z \in \sigma_{\text{ft}}(I - TS) \iff z \in \sigma_{\text{ft}}(I - ST),$$

*in which case*

$$m_a(z_0; I - TS) = m_a(z_0; I - ST).$$

# Outlines of the proofs

Proof.

# Outlines of the proofs

Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

# Outlines of the proofs

Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .



# Outlines of the proofs

## Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .

$W_{TS}$  Fredholm / bdd. inv.  $\iff W_{ST}$  Fredholm / bdd. inv.

# Outlines of the proofs

## Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .

$W_{TS}$  Fredholm / bdd. inv.  $\iff W_{ST}$  Fredholm / bdd. inv.

$$W_{ST}^{-1} \in \mathcal{B}(\mathcal{H}) \implies W_{TS}^{-1} = I + TW_{ST}^{-1}S$$

# Outlines of the proofs

## Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .

$W_{TS}$  Fredholm / bdd. inv.  $\iff W_{ST}$  Fredholm / bdd. inv.

$$W_{ST}^{-1} \in \mathcal{B}(\mathcal{H}) \implies W_{TS}^{-1} = I + TW_{ST}^{-1}S$$

**Step 2:**  $m_a(z_0; I - TS) = m_a(z_0; I - ST)$

# Outlines of the proofs

## Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .

$W_{TS}$  Fredholm / bdd. inv.  $\iff W_{ST}$  Fredholm / bdd. inv.

$$W_{ST}^{-1} \in \mathcal{B}(\mathcal{H}) \implies W_{TS}^{-1} = I + TW_{ST}^{-1}S$$

**Step 2:**  $m_a(z_0; I - TS) = m_a(z_0; I - ST)$

$$W'_{TS} W_{TS}^{-1} = -(T'S + TS')(I + TW_{ST}^{-1}S)$$

# Outlines of the proofs

## Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .

$W_{TS}$  Fredholm / bdd. inv.  $\iff W_{ST}$  Fredholm / bdd. inv.

$$W_{ST}^{-1} \in \mathcal{B}(\mathcal{H}) \implies W_{TS}^{-1} = I + TW_{ST}^{-1}S$$

**Step 2:**  $m_a(z_0; I - TS) = m_a(z_0; I - ST)$

$$W'_{TS} W_{TS}^{-1} = -(T'S + TS')(I + TW_{ST}^{-1}S)$$

$$\rightsquigarrow m_a(z_0; W_{TS}) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\gamma} \text{tr}\{W'_{TS} W_{TS}^{-1}\}$$

# Outlines of the proofs

## Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .

$W_{TS}$  Fredholm / bdd. inv.  $\iff W_{ST}$  Fredholm / bdd. inv.

$$W_{ST}^{-1} \in \mathcal{B}(\mathcal{H}) \implies W_{TS}^{-1} = I + TW_{ST}^{-1}S$$

**Step 2:**  $m_a(z_0; I - TS) = m_a(z_0; I - ST)$

$$W'_{TS} W_{TS}^{-1} = -(T'S + TS')(I + TW_{ST}^{-1}S)$$

$$\rightsquigarrow m_a(z_0; W_{TS}) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\gamma} \text{tr}\{W'_{TS} W_{TS}^{-1}\} \stackrel{\text{cycl.}}{=} \frac{1}{2\pi i} \int_{\gamma} \text{tr}\{W'_{ST} W_{ST}^{-1}\}$$

# Outlines of the proofs

## Proof.

Set  $W_{TS} := I - TS$  and  $W_{ST} := I - ST$ .

**Step 1:**  $\sigma_{\text{ff}}(W_{TS}) = \sigma_{\text{ff}}(W_{ST})$ .

$W_{TS}$  Fredholm / bdd. inv.  $\iff W_{ST}$  Fredholm / bdd. inv.

$$W_{ST}^{-1} \in \mathcal{B}(\mathcal{H}) \implies W_{TS}^{-1} = I + TW_{ST}^{-1}S$$

**Step 2:**  $m_a(z_0; I - TS) = m_a(z_0; I - ST)$

$$W'_{TS} W_{TS}^{-1} = -(T'S + TS')(I + TW_{ST}^{-1}S)$$

$$\begin{aligned} \rightsquigarrow m_a(z_0; W_{TS}) &\stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\gamma} \text{tr}\{W'_{TS} W_{TS}^{-1}\} \stackrel{\text{cycl.}}{=} \frac{1}{2\pi i} \int_{\gamma} \text{tr}\{W'_{ST} W_{ST}^{-1}\} \\ &\stackrel{\text{def}}{=} m_a(z_0; W_{ST}) \end{aligned}$$



# Bibliography

- [1] J.-C. Cuenin, A. Laptev, and C. Tretter. Eigenvalue estimates for non-selfadjoint Dirac operators on the real line. *Ann. Henri Poincaré* **15.4** (2014), 707–736.
- [2] J.-C. Cuenin and P. Siegl. Eigenvalues of one-dimensional non-self-adjoint Dirac operators and applications. *Lett. Math. Phys.* **108** (2018), 1757–1778.
- [3] I. Gohberg, S. Goldberg, and M. A. Kaashoek. *Classes of Linear Operators Vol. I*. Birkhäuser Verlag, 1990.



Thank you for your attention!