

Explicit n -particle harmonic oscillator states

Hubert de Guise
Lakehead University

and

Noah Kakekaspan
now at McGill University



Context

- In constructing multiparticle states, the permutation group is often essential.

Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$ systems gives:

Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$ systems gives:
one *antisymmetric* singlet $S = 0$ and

Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$ systems gives:
one *antisymmetric* singlet $S = 0$ and
one *symmetric* triplet $S = 1$.

Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$ systems gives:
one *antisymmetric* singlet $S = 0$ and
one *symmetric* triplet $S = 1$.
- The permutation symmetry uniquely identifies S when taking multiple copies of spin- $1/2$ states.

Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$ systems gives:
one *antisymmetric* singlet $S = 0$ and
one *symmetric* triplet $S = 1$.
- The permutation symmetry uniquely identifies S when taking multiple copies of spin-1/2 states.
- In combining single-particle harmonic oscillator states of the same parity ($\mathfrak{su}(1, 1) \sim \mathfrak{sp}(2, \mathbb{R})$ states), the permutation group is **not enough**.

Promise

- Construct many-particle $\mathfrak{su}(1,1)$ states “easily”,

Promise

- Construct many-particle $\mathfrak{su}(1, 1)$ states “easily”,
- Construct *some* many-particle $\mathfrak{sp}(4, \mathbb{R})$ states “easily”

Promise

- Construct many-particle $\mathfrak{su}(1, 1)$ states “easily”,
- Construct *some* many-particle $\mathfrak{sp}(4, \mathbb{R})$ states “easily”
- have an idea of where the permutation symmetry is hiding.

$\mathfrak{su}(1,1)$ oscillator states

- The $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$ operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

$\mathfrak{su}(1,1)$ oscillator states

- The $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$ operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

- (One particle) oscillator representation:

$$K_+ = \frac{1}{2} \hat{a}^\dagger \hat{a}^\dagger, \quad K_- = \frac{1}{2} \hat{a} \hat{a}, \quad K_0 = \frac{1}{2} (\hat{a}^\dagger \hat{a} + \frac{1}{2})$$

$$\begin{array}{c} \vdots \\ \text{---} |6\rangle \\ \text{---} |4\rangle \\ \text{---} |2\rangle \\ \text{---} |0\rangle \end{array}$$

$\mathfrak{su}(1,1)$ oscillator states

- The $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$ operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

- (One particle) oscillator representation:

$$K_+ = \frac{1}{2} \hat{a}^\dagger \hat{a}^\dagger, \quad K_- = \frac{1}{2} \hat{a} \hat{a}, \quad K_0 = \frac{1}{2} (\hat{a}^\dagger \hat{a} + \frac{1}{2})$$

- The bottom state $|0\rangle$ satisfies:

$$K_- |0\rangle = 0$$

$$\begin{array}{c} \vdots \\ \text{---} |6\rangle \\ \text{---} |4\rangle \\ \text{---} |2\rangle \\ \text{---} |0\rangle \end{array}$$

$\mathfrak{su}(1,1)$ oscillator states

- The $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$ operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

- (One particle) oscillator representation:

$$K_+ = \frac{1}{2} \hat{a}^\dagger \hat{a}^\dagger, \quad K_- = \frac{1}{2} \hat{a} \hat{a}, \quad K_0 = \frac{1}{2} (\hat{a}^\dagger \hat{a} + \frac{1}{2})$$

- The bottom state $|0\rangle$ satisfies:

$$K_- |0\rangle = 0$$

- Natural for boson systems



Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$, $K_+ = K_-^\dagger$, $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$

Combining two $\text{su}(1,1)$ systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$, $K_+ = K_-^\dagger$, $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.

Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$, $K_+ = K_-^\dagger$, $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.
- There are infinitely many towers of symmetric states.

⋮

—

— $|4\rangle|0\rangle + \sqrt{\frac{2}{3}}|2\rangle|2\rangle + |0\rangle|4\rangle$

— $|2\rangle|0\rangle + |0\rangle|2\rangle$

— $|0\rangle|0\rangle$

⋮

—

— $|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle$

Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$, $K_+ = K_-^\dagger$, $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.
- There are infinitely many towers of symmetric states.

\vdots

—

— $|4\rangle|0\rangle - |0\rangle|4\rangle$

— $|2\rangle|0\rangle - |0\rangle|2\rangle$

- There are infinitely many towers of antisymmetric states.

Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$, $K_+ = K_-^\dagger$, $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.
- There are infinitely many towers of symmetric states.

\vdots	\vdots	\vdots
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle + 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle + 0\rangle 4\rangle$
— $ 2\rangle 0\rangle + 0\rangle 2\rangle$	— $ 2\rangle 0\rangle - 0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

- Clearly permutation symmetry is not enough to identify the tower of states.

The Laplacian approach

\vdots	\vdots	\vdots
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle + 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle + 0\rangle 4\rangle$
— $ 2\rangle 0\rangle + 0\rangle 2\rangle$	— $ 2\rangle 0\rangle - 0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

The Laplacian approach

\vdots	\vdots	\vdots
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle + 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle + 0\rangle 4\rangle$
— $ 2\rangle 0\rangle + 0\rangle 2\rangle$	— $ 2\rangle 0\rangle - 0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

- The 3 bottom states satisfy

The Laplacian approach

\vdots	\vdots	\vdots
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle + 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - 0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle + 0\rangle 4\rangle$
— $ 2\rangle 0\rangle + 0\rangle 2\rangle$	— $ 2\rangle 0\rangle - 0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

- The 3 bottom states satisfy

$$\begin{aligned}
 K_-|0\rangle|0\rangle &= 0, \\
 K_- (|2\rangle|0\rangle - |0\rangle|2\rangle) &= 0, \\
 K_- (|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle) &= 0
 \end{aligned}$$

The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

- The bottom states map to polynomials:

$$|0\rangle|0\rangle \mapsto f_0 = 1,$$

$$|2\rangle|0\rangle - |0\rangle|2\rangle \mapsto f_2 = x_1^2 - x_2^2,$$

$$|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle \mapsto f_4 = x_1^4 - 6x_1^2x_2^2 + x_2^4$$

The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

- The bottom states map to polynomials:

$$|0\rangle|0\rangle \mapsto f_0 = 1,$$

$$|2\rangle|0\rangle - |0\rangle|2\rangle \mapsto f_2 = x_1^2 - x_2^2,$$

$$|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle \mapsto f_4 = x_1^4 - 6x_1^2x_2^2 + x_2^4$$

- Now we have

$$K_- f_k \mapsto \nabla^2 f_k = 0.$$

The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

- The bottom states map to polynomials:

$$|0\rangle|0\rangle \mapsto f_0 = 1,$$

$$|2\rangle|0\rangle - |0\rangle|2\rangle \mapsto f_2 = x_1^2 - x_2^2,$$

$$|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle \mapsto f_4 = x_1^4 - 6x_1^2x_2^2 + x_2^4$$

- Now we have

$$K_- f_k \mapsto \nabla^2 f_k = 0.$$

- The f_k are two-dimensional harmonic functions.

The Laplacian approach

- Solutions f_k have a generating function:

The Laplacian approach

- Solutions f_k have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2}$$

The Laplacian approach

- Solutions f_k have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2} \approx 1 + h_2(x_1 + ix_2) + h_2^2 \frac{1}{2}(x_1 + ix_2)^2 + \dots$$

The Laplacian approach

- Solutions f_k have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2} \approx 1 + h_2(x_1 + ix_2) + h_2^2 \frac{1}{2}(x_1 + ix_2)^2 + \dots$$

- In the expansion of $H_2^+(x_1, x_2)$, distinct powers of h_2 give distinct functions f_k that satisfy $\nabla^2 f_k = 0$.

The Laplacian approach

- Solutions f_k have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2} \approx 1 + h_2(x_1 + ix_2) + h_2^2 \frac{1}{2}(x_1 + ix_2)^2 + \dots$$

- In the expansion of $H_2^+(x_1, x_2)$, distinct powers of h_2 give distinct functions f_k that satisfy $\nabla^2 f_k = 0$.
- Only even numbers of excitations are possible so even powers of x_1 or x_2 must be kept:

$$1 + \frac{1}{2}h_2^2(x_1^2 - x_2^2) + \frac{1}{24}h_2^4(x_1^4 - 6x_1^2x_2^2 + x_2^4) + \dots$$

The Laplacian approach

- For three particles:

$$\nabla^2 = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial x_3^2} = 0.$$

The Laplacian approach

- For three particles:

$$\nabla^2 = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial x_3^2} = 0.$$

- There's also a generating function for the solutions to this.



Connection with spherical harmonics

- Again keeping only terms in even powers of x_i :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

Connection with spherical harmonics

- Again keeping only terms in even powers of x_i :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

Connection with spherical harmonics

- Again keeping only terms in even powers of x_i :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

Connection with spherical harmonics

- Again keeping only terms in even powers of x_i :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

$$r^2(Y_2^{-2}(\theta, \varphi) + Y_2^2(\theta, \varphi)) \sim x_1^2 - x_2^2$$

Connection with spherical harmonics

- Again keeping only terms in even powers of x_i :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

$$\begin{aligned} r^2(Y_2^{-2}(\theta, \varphi) + Y_2^2(\theta, \varphi)) &\sim x_1^2 - x_2^2 \\ r^2 Y_2^0(\theta, \varphi) &\sim x_1^2 + x_2^2 - 2x_3^2 \end{aligned}$$

Connection with spherical harmonics

- Again keeping only terms in even powers of x_i :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

$$\begin{aligned} r^2(Y_2^{-2}(\theta, \varphi) + Y_2^2(\theta, \varphi)) &\sim x_1^2 - x_2^2 \\ r^2 Y_2^0(\theta, \varphi) &\sim x_1^2 + x_2^2 - 2x_3^2 \end{aligned}$$

- There is in fact a symmetry to the solutions: angular momentum plus parity.

Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$

Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$

- Introduce

$$\hat{L}_{12} = i \left(\hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1 \right) \rightarrow \hat{L}_z, \quad \hat{L}_{23} = i \left(\hat{a}_2^\dagger \hat{a}_3 - \hat{a}_3^\dagger \hat{a}_2 \right) \rightarrow \hat{L}_x,$$
$$\hat{L}_{31} = i \left(\hat{a}_3^\dagger \hat{a}_1 - \hat{a}_1^\dagger \hat{a}_3 \right) \rightarrow \hat{L}_y.$$

Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$

- Introduce

$$\hat{L}_{12} = i \left(\hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1 \right) \rightarrow \hat{L}_z, \quad \hat{L}_{23} = i \left(\hat{a}_2^\dagger \hat{a}_3 - \hat{a}_3^\dagger \hat{a}_2 \right) \rightarrow \hat{L}_x,$$
$$\hat{L}_{31} = i \left(\hat{a}_3^\dagger \hat{a}_1 - \hat{a}_1^\dagger \hat{a}_3 \right) \rightarrow \hat{L}_y.$$

- Of course $[\hat{L}_j, \hat{L}_k] = i\epsilon_{jkl} \hat{L}_l$

Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left(\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$

- Introduce

$$\hat{L}_{12} = i \left(\hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1 \right) \rightarrow \hat{L}_z, \quad \hat{L}_{23} = i \left(\hat{a}_2^\dagger \hat{a}_3 - \hat{a}_3^\dagger \hat{a}_2 \right) \rightarrow \hat{L}_x,$$
$$\hat{L}_{31} = i \left(\hat{a}_3^\dagger \hat{a}_1 - \hat{a}_1^\dagger \hat{a}_3 \right) \rightarrow \hat{L}_y.$$

- Of course $[\hat{L}_j, \hat{L}_k] = i\epsilon_{jkl} \hat{L}_l$
- Also $[\hat{L}_j, K_q] = 0$.

Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers ℓ and $|m|$.

Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers ℓ and $|m|$.
- Because of parity, states are combo of $|m|$ states with m even.

Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers ℓ and $|m|$.
- Because of parity, states are combo of $|m|$ states with m even.
- Recall

$$K_+ = a_1^\dagger a_1^\dagger + a_2^\dagger a_2^\dagger + a_3^\dagger a_3^\dagger$$

Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers ℓ and $|m|$.
- Because of parity, states are combo of $|m|$ states with m even.
- Recall

$$K_+ = a_1^\dagger a_1^\dagger + a_2^\dagger a_2^\dagger + a_3^\dagger a_3^\dagger \quad \mapsto x_1^2 + x_2^2 + x_3^2 = r^2$$

Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers ℓ and $|m|$.
- Because of parity, states are combo of $|m|$ states with m even.
- Recall

$$K_+ = a_1^\dagger a_1^\dagger + a_2^\dagger a_2^\dagger + a_3^\dagger a_3^\dagger \quad \mapsto x_1^2 + x_2^2 + x_3^2 = r^2$$

- The states $|\ell, m; n\rangle$ in $\ell = 0, m = 0$ tower:

$$Y_0^0(\theta, \varphi) = 1 \quad \sim |0\rangle|0\rangle|0\rangle \quad = |00; 0\rangle$$

$$r^2 Y_0^0(\theta, \varphi) = x_1^2 + x_2^2 + x_3^2 \quad \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle + |0\rangle|0\rangle|2\rangle = |00; 2\rangle$$

$$r^4 Y_0^0(\theta, \varphi) = (x_1^2 + x_2^2 + x_3^2)^2 \quad = |00; 4\rangle$$

Three particles and $O(3)$

- The $\ell = 4, m = 0$ tower are built on:

$$r^4 Y_4^0(\theta, \varphi) \sim |40; 4\rangle = |4\rangle|0\rangle|0\rangle + \sqrt{\frac{2}{3}}|2\rangle|2\rangle|0\rangle - 4\sqrt{\frac{2}{3}}|2\rangle|0\rangle|2\rangle + \frac{8}{3}|0\rangle|0\rangle|4\rangle$$

Three particles and $O(3)$

- The $\ell = 4, m = 0$ tower are built on:

$$r^4 Y_4^0(\theta, \varphi) \sim |40; 4\rangle = |4\rangle|0\rangle|0\rangle + \sqrt{\frac{2}{3}}|2\rangle|2\rangle|0\rangle - 4\sqrt{\frac{2}{3}}|2\rangle|0\rangle|2\rangle + \frac{8}{3}|0\rangle|0\rangle|4\rangle$$

- We also have $\ell = 4, |m| = 2$ and $|m| = 4$ towers built on:

$$\begin{aligned} r^4(Y_4^2(\theta, \varphi) + Y_4^{-2}(\theta, \varphi)) \sim |42; 4\rangle \sim & |4\rangle|0\rangle|0\rangle - \sqrt{6}|2\rangle|0\rangle|2\rangle \\ & - |0\rangle|4\rangle|0\rangle + \sqrt{6}|0\rangle|2\rangle|2\rangle \end{aligned}$$

$$r^4(Y_4^4(\theta, \varphi) + Y_4^{-4}(\theta, \varphi)) \sim |44; 4\rangle \sim |4\rangle|0\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle|0\rangle + |0\rangle|4\rangle|0\rangle$$

Where did the permutation symmetry go?

Where did the permutation symmetry go?

- The combination $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$ is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}} (|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the $|00; 0\rangle$ state.

Where did the permutation symmetry go?

- The combination $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$ is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the $|00; 0\rangle$ state.

- There are two remaining states with are partially symmetric.

Where did the permutation symmetry go?

- The combination $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$ is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}} (|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the $|00; 0\rangle$ state.

- There are two remaining states which are partially symmetric.
One goes to a linear combination of the other two under permutation.

Where did the permutation symmetry go?

- The combination $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$ is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the $|00; 0\rangle$ state.

- There are two remaining states which are partially symmetric. One goes to a linear combination of the other two under permutation.
- The permutation group appears as a subgroup of $O(3)$:

Where did the permutation symmetry go?

- The combination $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$ is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the $|00; 0\rangle$ state.

- There are two remaining states with are partially symmetric.
One goes to a linear combination of the other two under permutation.
- The permutation group appears as a subgroup of $O(3)$:

$$P_{12} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{is an orthogonal matrix: } P_{12}^\top = P_{12}^{-1}$$

More than 3 particles

- The strategy works for $d \geq 3$ particles but...

More than 3 particles

- The strategy works for $d \geq 3$ particles but...
 - The generating function becomes complicated.

More than 3 particles

- The strategy works for $d \geq 3$ particles but...
 - The generating function becomes complicated.
 - Other methods use hyperspherical coordinates.

More than 3 particles

- The strategy works for $d \geq 3$ particles but...
 - The generating function becomes complicated.
 - Other methods use hyperspherical coordinates.
 - Tensor operator methods

Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a $u(2)$ subalgebra.

Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a $u(2)$ subalgebra.

- The operators $\{A_{11}, A_{12}, A_{22}\}$ transform under the $u(2)$ as states in the $u(2)$ irrep $(2, 0)$.

Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a $u(2)$ subalgebra.

- The operators $\{A_{11}, A_{12}, A_{22}\}$ transform under the $u(2)$ as states in the $u(2)$ irrep $(2, 0)$.
- As per $su(1, 1)$, the idea is to start from the bottom of the tower and repeated act with powers of the A_{ij} :

$$A_{ij} A_{kl} \in [(2, 0) \otimes (2, 0)]_{\text{symm}}$$

Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left(a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a $u(2)$ subalgebra.

- The operators $\{A_{11}, A_{12}, A_{22}\}$ transform under the $u(2)$ as states in the $u(2)$ irrep $(2, 0)$.
- As per $su(1, 1)$, the idea is to start from the bottom of the tower and repeated act with powers of the A_{ij} :

$$A_{ij} A_{kl} \in [(2, 0) \otimes (2, 0)]_{\text{symm}}$$

- Because A_{ij} 's commute only the symmetric part of the tensor product is needed.

Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of $(2, 0)$ irreps contains states in the $u(2)$ irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of $(2, 0)$ irreps contains states in the $u(2)$ irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

- the sum is over partitions of even integers having even-only parts.

J. Phys. A: Math. Gen. 18 (1985) 939-953. Printed in Great Britain

Unitary representations, branching rules and matrix elements for the non-compact symplectic groups

D J Rowe†, B G Wybourne and P H Butler

Physics Department, University of Canterbury, Christchurch 1, New Zealand

Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of $(2, 0)$ irreps contains states in the $u(2)$ irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

- the sum is over partitions of even integers having even-only parts.
- These must in turn be coupled to the bottom state of each tower.

J. Phys. A: Math. Gen. 18 (1985) 939-953. Printed in Great Britain

Unitary representations, branching rules and matrix elements for the non-compact symplectic groups

D J Rowe†, B G Wybourne and P H Butler

Physics Department, University of Canterbury, Christchurch 1, New Zealand

Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of $(2, 0)$ irreps contains states in the $u(2)$ irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

- the sum is over partitions of even integers having even-only parts.
- These must in turn be coupled to the bottom state of each tower.
- For single particles irrep of $sp(4, \mathbb{R})$ this bottom state is carries the $(0, 0)$ irrep so D lists the $u(2)$ contents of the single particle irrep.

J. Phys. A: Math. Gen. 18 (1985) 939-953. Printed in Great Britain

Unitary representations, branching rules and matrix elements for the non-compact symplectic groups

D J Rowe†, B G Wybourne and P H Butler

Physics Department, University of Canterbury, Christchurch 1, New Zealand

Two-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^2 a_{ki}^{\dagger} a_{kj}^{\dagger},$$

$$C_{ij} = \sum_{k=1}^2 \frac{1}{2} \left(a_{ki}^{\dagger} a_{kj} + a_{kj} a_{ki}^{\dagger} \right),$$

$$B_{ij} = A_{ij}^{\dagger}.$$

Two-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^2 a_{ki}^{\dagger} a_{kj}^{\dagger},$$

$$C_{ij} = \sum_{k=1}^2 \frac{1}{2} \left(a_{ki}^{\dagger} a_{kj} + a_{kj} a_{ki}^{\dagger} \right),$$

$$B_{ij} = A_{ij}^{\dagger}.$$

- The lowering operators

$$B_{11} \mapsto \frac{\partial^2}{\partial x_{11}^2} + \frac{\partial^2}{\partial x_{21}^2},$$

$$B_{12} \mapsto \frac{\partial^2}{\partial x_{11} \partial x_{12}} + \frac{\partial^2}{\partial x_{21} \partial x_{22}},$$

$$B_{22} \mapsto \frac{\partial^2}{\partial x_{12}^2} + \frac{\partial^2}{\partial x_{22}^2}.$$

Two-particle irreps of $sp(4, \mathbb{R})$

- Not so easy to find the functions f *simultaneously* solutions to

$$\begin{aligned}\left(\frac{\partial^2}{\partial x_{11}^2} + \frac{\partial^2}{\partial x_{21}^2}\right) f &= 0, \\ \left(\frac{\partial^2}{\partial x_{11} \partial x_{12}} + \frac{\partial^2}{\partial x_{21} \partial x_{22}}\right) f &= 0 \\ \left(\frac{\partial^2}{\partial x_{12}^2} + \frac{\partial^2}{\partial x_{22}^2}\right) f &= 0.\end{aligned}$$

Two-particle irreps of $sp(4, \mathbb{R})$

- Use $O(2)$ by constructing the tensors

$$T^1 = \frac{1}{2} \left(a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left(a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$: particle number i , mode number α .

- Note that

Clarify what is L12

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

Two-particle irreps of $sp(4, \mathbb{R})$

- Use $O(2)$ by constructing the tensors

$$T^1 = \frac{1}{2} \left(a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left(a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$: particle number i , mode number α .

- Note that

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

- It appears all lowest $sp(4, \mathbb{R})$ states for 2 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle \sim V^{2n}|00\rangle|00\rangle \sim V^{2k}T^{2(n-k)}|00\rangle|00\rangle$$

Two-particle irreps of $sp(4, \mathbb{R})$

- Use $O(2)$ by constructing the tensors

$$T^1 = \frac{1}{2} \left(a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left(a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$: particle number i , mode number α .

- Note that

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

- It appears all lowest $sp(4, \mathbb{R})$ states for 2 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle \sim V^{2n}|00\rangle|00\rangle \sim V^{2k}T^{2(n-k)}|00\rangle|00\rangle$$

- Example: $|m| = 4$ irrep has lowest state

$$x_{11}^4 - 6x_{11}^2x_{21}^2 + x_{21}^4 \sim |40\rangle|00\rangle - \sqrt{6}|20\rangle|20\rangle + |00\rangle|40\rangle$$

Two-particle irreps of $sp(4, \mathbb{R})$

- Use $O(2)$ by constructing the tensors

$$T^1 = \frac{1}{2} \left(a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left(a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$: particle number i , mode number α .

- Note that

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

- It appears all lowest $sp(4, \mathbb{R})$ states for 2 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle \sim V^{2n}|00\rangle|00\rangle \sim V^{2k}T^{2(n-k)}|00\rangle|00\rangle$$

- Example: $|m| = 4$ irrep has lowest state

$$x_{11}^4 - 6x_{11}^2x_{21}^2 + x_{21}^4 \sim |40\rangle|00\rangle - \sqrt{6}|20\rangle|20\rangle + |00\rangle|40\rangle$$

- Compare the $su(1, 1)$ states at the bottom of $|m| = 4$:

$$x_1^4 - 6x_1^2x_2^2 + x_2^4 \sim |4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle$$

Three-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^3 a_{ki}^{\dagger} a_{kj}^{\dagger},$$

$$C_{ij} = \sum_{k=1}^3 \frac{1}{2} \left(a_{ki}^{\dagger} a_{kj} + a_{kj} a_{ki}^{\dagger} \right),$$

$$B_{ij} = A_{ij}^{\dagger}.$$

Three-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^3 a_{ki}^\dagger a_{kj}^\dagger,$$

$$C_{ij} = \sum_{k=1}^3 \frac{1}{2} \left(a_{ki}^\dagger a_{kj} + a_{kj} a_{ki}^\dagger \right),$$

$$B_{ij} = A_{ij}^\dagger.$$

- Use $O(3)$ by constructing the tensors T_M^L and V_M^L :

$$T_{-1}^1 = -\frac{1}{2} \left(a_{11}^\dagger + i a_{21}^\dagger \right), \quad T_0^1 = -\frac{1}{\sqrt{2}} a_{31}^\dagger, \quad T_1^1 = \frac{1}{2} \left(a_{11}^\dagger - i a_{21}^\dagger \right),$$

$$V_{-1}^1 = -\frac{1}{2} \left(a_{12}^\dagger + i a_{22}^\dagger \right), \quad V_0^1 = -\frac{1}{\sqrt{2}} a_{32}^\dagger, \quad V_1^1 = \frac{1}{2} \left(a_{12}^\dagger - i a_{22}^\dagger \right).$$

Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest $sp(4, \mathbb{R})$ states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest $sp(4, \mathbb{R})$ states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

- Some are: $L = 2, |m| = 0$ irrep has lowest state

$$x_{11}^2 + x_{21}^2 - 2x_{31}^2 \sim |20\rangle|00\rangle|00\rangle + |00\rangle|20\rangle|00\rangle - 2|00\rangle|00\rangle|20\rangle$$

- Compare the $su(1, 1)$ states at the bottom of $L = 2, |m| = 0$:

$$x_1^2 + x_2^2 - 2x_3^2 \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest $sp(4, \mathbb{R})$ states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

- Some are: $L = 2, |m| = 0$ irrep has lowest state

$$x_{11}^2 + x_{21}^2 - 2x_{31}^2 \sim |20\rangle|00\rangle|00\rangle + |00\rangle|20\rangle|00\rangle - 2|00\rangle|00\rangle|20\rangle$$

- Compare the $su(1, 1)$ states at the bottom of $L = 2, |m| = 0$:

$$x_1^2 + x_2^2 - 2x_3^2 \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

- Some aren't:

$$\sum_{mm'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \middle| \begin{matrix} 3 \\ 2 \end{matrix} \right\rangle T_m^2 V_{m'}^2 |00\rangle|00\rangle|00\rangle$$

Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest $sp(4, \mathbb{R})$ states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

- Some are: $L = 2, |m| = 0$ irrep has lowest state

$$x_{11}^2 + x_{21}^2 - 2x_{31}^2 \sim |20\rangle|00\rangle|00\rangle + |00\rangle|20\rangle|00\rangle - 2|00\rangle|00\rangle|20\rangle$$

- Compare the $su(1, 1)$ states at the bottom of $L = 2, |m| = 0$:

$$x_1^2 + x_2^2 - 2x_3^2 \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

- Some aren't:

$$\sum_{mm'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \middle| \begin{matrix} 3 \\ 2 \end{matrix} \right\rangle T_m^2 V_{m'}^2 |00\rangle|00\rangle|00\rangle$$

- T^2 and V^2 are $L = 2$ tensors under $O(3)$, and the combo

$\sum_{mm'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \middle| \begin{matrix} 3 \\ 2 \end{matrix} \right\rangle T_m^2 V_{m'}^2$ is an “axial tensor” in the sense that it lives in the antisymmetric space of the decomposition $(L = 2) \otimes (L = 2)$.

Three-particle irreps of $sp(4, \mathbb{R})$

- Some don't exist:

$$\sum_{m m'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \mid \begin{matrix} 1 \\ 0 \end{matrix} \right\rangle T_m^2 V_{m'}^2 |00\rangle |00\rangle |00\rangle$$

does not yield product states with only even number of excitations for each particles.

- This is because $m = 0$ irreps of $O(2)$ (as a subgroup of $O(3)$) must have even parity but the coupling is antisymmetric in T and V .

Conclusion

- Angular momentum helps in constructing multiple states when combining degenerate representations of $\mathfrak{su}(1, 1)$.

Conclusion

- Angular momentum helps in constructing multiple states when combining degenerate representations of $\mathfrak{su}(1, 1)$.
- It is also helpful for 2-particle irreps of $\mathfrak{sp}(2k, \mathbb{R})$.

Conclusion

- Angular momentum helps in constructing multiple states when combining degenerate representations of $\mathfrak{su}(1, 1)$.
- It is also helpful for 2-particle irreps of $\mathfrak{sp}(2k, \mathbb{R})$.
- Some extra care required for $n \geq 3$ particles

